Lower bounds for geometric diameter problems

H. Fournier, Antoine Vigneron

► To cite this version:

H. Fournier, Antoine Vigneron. Lower bounds for geometric diameter problems. 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Mar 2006, Valdivia, Chile. hal-02753352

HAL Id: hal-02753352

<https://hal.inrae.fr/hal-02753352>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lower Bounds for Geometric Diameter Problems

Hervé Fournier

University of Versailles St-Quentin en Yvelines

Antoine Vigneron

INRA Jouy-en-Josas

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Outline

- Review of previous work on the 2D and 3D diameter problems.
- $\Omega(n \log n)$ lower bound for computing the diameter of a 3D convex polytope.
- Reduction from Hopcroft's problem to the diameter problem for point sets in $\mathbb{R}^7$.

Previous work

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

The diameter problem

- INPUT: a set P of n points in $\mathbb{R}^d$.
- OUTPUT: $\text{diam}(P) := \max\{d(x, y) \mid x, y \in P\}$.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

The diameter problem

- $\text{diam}(P) = d(p, p')$.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Decision problem

- We will give lower bounds for the *decision problem* associated with the diameter problem.
- INPUT: a set P of n points in $\mathbb{R}^d$.
- OUTPUT:
 - YES if $\text{diam}(P) < 1$
 - NO if $\text{diam}(P) \geq 1$

Observation

- P lies in the intersection of the two balls with radius $d(p, p')$ centered at p and p' .

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887. Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

The diameter problem

- P lies between two parallel hyperplanes through p and p' . We say that (p, p') is an *antipodal pair*.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

The diameter problem

- Any antipodal pair (and therefore any diametral pair) lies on the convex hull $CH(P)$ of P .

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887. Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Finding the antipodal pairs

- The rotating calipers technique.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Finding the antipodal pairs

- The rotating calipers technique.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887. Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Finding the antipodal pairs

- The rotating calipers technique.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Finding the antipodal pairs

- The rotating calipers technique.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Finding the antipodal pairs

- The rotating calipers technique.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Computing the diameter of a 2D-point set

- Compute the convex hull $CH(P)$ of P .
 - $O(n \log n)$ time.
- Find all the antipodal pairs on $CH(P)$.
 - There are at most n such pairs in non-degenerate cases.
 - $O(n)$ time using the rotating calipers technique.
- Find the diametral pairs among the antipodal pairs.
 - $O(n)$ time by brute force.
- Conclusion:
 - The diameter of a 2D-point set can be found in $O(n \log n)$ time
 - The diameter of a convex polygon can be found in $O(n)$ time.

3D-diameter problem

- Randomized $O(n \log n)$ time algorithm (Clarkson and Shor, 1988).
 - Randomized incremental construction of an intersection of balls and decimation.
- Deterministic $O(n \log n)$ time algorithm (E. Ramos, 2000).
- These two algorithms compute the diameter of an n -point set in $\mathbb{R}^3$.
- Can we compute the diameter of a convex 3D-polytope in linear time?
 - No, we give an $\Omega(n \log n)$ lower bound.

Model of computation

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Real-RAM

- Real Random Access Machine.
- Each registers stores a *real* number.
- Access to registers in unit time.
- Arithmetic operation $(+, -, \times, /)$ in unit time.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Algebraic computation tree: definition

- Input: $\bar{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$.
- Output: YES or NO
- It is a binary tree with 3 types of nodes
 - **Leaves**: YES or NO
 - Degree-1 nodes: **computation nodes**. Perform $\{+, -, \times, /, \sqrt{\cdot}\}$ on two operands. An operand is
 - a real constant,
 - some input number x_i , or
 - the value obtained by computation nodes that are ancestors of the current node.
 - Degree-2 nodes: **branching nodes**. Compares with 0 the value obtained at a computation node that is an ancestor of the current node.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887. Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Algebraic computation tree (ACT)

- We say that an ACT *decides* $S \subset \mathbb{R}^n$ if
 - $\forall (x_1, \dots, x_n) \in S$, it reaches a leaf labeled YES, and
 - $\forall (x_1, \dots, x_n) \notin S$, it reaches a leaf labeled NO.
- The ACT model is stronger than the real-RAM model.
- To get a lower bound on the worst-case running time of a real-RAM that decides S , it suffices to have a lower bound on the *depth* of all the ACTs that decide S

Theorem 1 (Ben-Or) *Any ACT that decides S has depth*

$$\Omega(\log(\text{number of connected components of } S)).$$

Lower bound for 3D convex polytopes

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Problem statement

- We are given a convex 3-polytope P with n vertices.
- P is given by the coordinates of its vertices and its *combinatorial structure*:
 - All the inclusion relations between its vertices, edges and faces.
 - The cyclic ordering of the edges of each face.
- Example: P is given in a doubly-connected edge list.
- Problem: we want to decide whether $\text{diam}(P) < 1$.
- We show an $\Omega(n \log n)$ lower bound. Our approach:
 - We define a family of convex polytopes.
 - We show that the sub-family with diameter < 1 has $n^{\Omega(n)}$ connected components.
 - We apply Ben-Or's bound.

Polytopes $P(\bar{\beta})$

- The family of polytopes is parametrized by $\bar{\beta} \in \mathbb{R}^{2n-1}$.
- When n is fixed, only the $2n - 1$ blue points change with $\bar{\beta}$.

Polytopes $P(\bar{\beta})$

- The family of polytopes is parametrized by $\bar{\beta} \in \mathbb{R}^{2n-1}$.
- When n is fixed, only the $2n - 1$ blue points change with $\bar{\beta}$.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887. Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Notation

- Example where $n = 3$.

Comment citer ce document :

Notation

- $\bar{a} := (a_{-n}, a_{-n+1} \dots, a_n)$.
- $A := \{a_{-n}, a_{-n+1}, \dots, a_n\}$.
- $\bar{\beta} := (\beta_{-n+1}, \dots, \beta_{n-1})$.
- $B(\bar{\beta}) := \{b_{-n+1}(\beta_{-n+1}), \dots, b_{n-1}(\beta_{n-1})\}$.
- $\bar{b}(\bar{\beta}) := (b_{-n+1}(\beta_{-n+1}), \dots, b_{n-1}(\beta_{n-1}))$.
- $\bar{c} := (c_{-n}^{-1}, c_{-n+1}^{-1}, \dots, c_{n-1}^{-1}, c_{-n}^1, c_{-n+1}^1, \dots, c_{n-1}^1)$.
- $C := \{c_{-n}^{-1}, c_{-n+1}^{-1}, \dots, c_{n-1}^{-1}, c_{-n}^1, c_{-n+1}^1, \dots, c_{n-1}^1\}$.
- $P(\bar{\beta}) := \text{CH}(A \cup B(\bar{\beta}) \cup C)$.

Point sets A and C

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Point $b_j(\beta_j)$

- The blue region is parallel to Oxz .
- $\beta \in [-\alpha, \alpha]$

Comment citer ce document :

Coordinates of points in A , $B(\bar{\beta})$ and C

$$a_i := \begin{pmatrix} \frac{1}{2}(1 - \cos(i\gamma)) \\ 0 \\ \frac{1}{2} \sin(i\gamma) \end{pmatrix}$$

$$c_i^s := \begin{pmatrix} r \cos \left(\left(i + \frac{1}{2} \right) \psi \right) \\ r \sin \left(\left(i + \frac{1}{2} \right) \psi \right) \\ \frac{1}{2} s \alpha \end{pmatrix}$$

$$b_j(\beta) := \begin{pmatrix} \cos(j\psi) - \frac{1}{2}(1 - \cos \beta) \\ \sin(j\psi) \\ \frac{1}{2} \sin(\beta) \end{pmatrix}$$

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Parameters

- $\varphi = 1/4n$
- α is small.
- $\psi = \varphi/n$
- $\gamma = \alpha/n$

Proof

- Notation: $\text{diam}(E, F) := \max\{d(e, f) \mid (e, f) \in E \times F\}$.

Lemma 1 *The set*

$$\{b_j(\beta) \mid \beta \in [-\alpha, \alpha] \text{ and } \text{diam}(A, \{b_j(\beta)\}) < 1\}$$

has at least $2n$ connected components.

Comment citer ce document :

Proof

- Proof of Lemma 1: Calculations, until the second-order terms.

Lemma 2 *The combinatorial structure of $\text{CH}(A \cup B(\bar{\beta}) \cup C)$ is independent of $\bar{\beta}$.*

- We denote $P(\bar{\beta}) = \text{CH}(A \cup B(\bar{\beta}) \cup C)$.

Lemma 3 $\text{diam}(A \cup B(\bar{\beta}) \cup C) = \text{diam}(A, B(\bar{\beta}))$.

Proof

- Definitions:

$$\mathcal{S}_n = \{(\bar{a}, \bar{b}(\bar{\beta}), \bar{c}) \mid \bar{\beta} \in [-\alpha, \alpha]^{2n-1}\}$$

$$\mathcal{E}_n = \{(\bar{a}, \bar{b}(\bar{\beta}), \bar{c}) \mid \bar{\beta} \in [-\alpha, \alpha]^{2n-1} \text{ and } \text{diam}(P(\bar{\beta})) < 1\}$$

- Notice that $\mathcal{E}_n \subset \mathcal{S}_n \subset \mathbb{R}^{24n}$.
- **Lemma 4** *The set $\mathcal{S}_n$ can be decided by an ACT with depth $O(n)$.*
- **Lemma 5** *Any ACT that decides $\mathcal{E}_n$ has depth $\Omega(n \log n)$.*

Proof: By lemmas 1 and 3, $\mathcal{E}_n$ has at least $(2n)^{2n-1}$ connected components. Apply Ben-Or's bound (Theorem 1).

End of the proof

Theorem 2 *Assume that an algebraic computation tree T_n decides whether the diameter of a 3-polytope is smaller than 1. Then T_n has depth $\Omega(n \log n)$.*

- Proof: Take as input graph of T_n the the graph of $P(\bar{\beta})$. Denote by U_n the ACT of Lemma 4. Plug U_n into the accepting leaves of T_{8n} . It gives an ACT with depth the depth of $T_{8n} + O(n)$ that decides $\mathcal{E}_n$. Apply Lemma 5.

Related work

- (Chazelle) The convex hull of two 3-polytopes can be computed in linear time.
- (Chazelle et al.) It is not known whether the convex hull of a subset of the vertices of a 3-polytope can be computed in linear time.
- (Chazelle et al.) However, we can compute in linear time the Delaunay triangulation of a subset of the vertices of a Delaunay triangulation.

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Diameter is harder than Hopcroft's problem

Comment citer ce document :

Fournier, H., Vigneron, A. (2006). Lower bounds for geometric diameter problems. In: Latin 2006 : theoretical informatics (p. 467-478). Lecture Notes in Computer Science, 3887.
Presented at 7. Latin American Symposium on Theoretical Informatics (LATIN 2006), Valdivia, CHL (2006-03-20 - 2006-03-24). Heidelberg, DEU : Springer-Verlag.

Hopcroft's problem

- P is a set of n points in $\mathbb{R}^2$.
- L is a set of n lines in $\mathbb{R}^2$.
- Problem: decide whether

$$\exists (p, \ell) \in P \times L : p \in \ell.$$

- An $o(n^{4/3} \log n)$ algorithm is known. (Matoušek).
- No $o(n^{4/3})$ algorithm is known.
- We give a linear-time reduction from Hopcroft's problem to the diameter problem in $\mathbb{R}^7$.
- We first give a reduction to the *red-blue diameter* problem in $\mathbb{R}^6$: compute $\text{diam}(E, F)$ when E and F are n -point sets in $\mathbb{R}^6$.

Proof

- $\theta(x, y, z) := \frac{1}{x^2 + y^2 + z^2} (x^2, y^2, y^2, \sqrt{2}xy, \sqrt{2}yz, \sqrt{2}zx).$
- Note that $\|\theta(x, y, z)\| = 1.$
- For $1 \leq i \leq n$
 - $p_i = (x_i, y_i, 1)$
 - $\ell_i = (u_i, v_i, w_i)$ is the line $\ell_i : u_i x + v_i y + w_i = 0.$
- Let $p'_i := \theta(p_i)$ and $\ell'_j = \theta(\ell_j).$
- We get

$$\begin{aligned} \|p'_i - \ell'_j\|^2 &= \|p'_i\|^2 + \|\ell'_j\|^2 - 2 \langle p'_i, \ell'_j \rangle \\ &= 2 - 2 \frac{\langle p_i, \ell_j \rangle^2}{\|p_i\|^2 \|\ell_j\|^2} \end{aligned}$$

Comment citer ce document :

Proof

- Note that $p_i \in \ell_j$ iff $\langle p_i, \ell_j \rangle = 0$.
- Thus, there exists i, j such that $p_i \in \ell_j$ if and only if $\text{diam}(\theta(P), \theta(L)) = 2$.
- $\theta(P)$ and $\theta(L)$ are n -point sets in $\mathbb{R}^6$.
- Similarly, we can get a reduction from Hopcroft's problem to the diameter problem in $\mathbb{R}^7$, using this linearization:

$$\theta(x, y, z) := \frac{1}{x^2 + y^2 + z^2} (x^2, y^2, z^2, \sqrt{2}xy, \sqrt{2}yz, \sqrt{2}zx)$$

Related work

- The red-blue diameter in $\mathbb{R}^4$ can be computed in $O(n^{4/3} \text{polylog } n)$ (Matoušek and Scharzkopf). It would be interesting to get a reduction from Hopcroft's problem.
- Erickson gave reduction from Hopcroft problem to other computational geometry problems.