



**HAL**  
open science

## Proteomic and morphological characterization of isolated rainbow trout adipocytes originating from various fat tissues

Claudine Weil, Nathalie Sabin, Jérôme Bugeon, Gilles Paboeuf, Florence Lefèvre

### ► To cite this version:

Claudine Weil, Nathalie Sabin, Jérôme Bugeon, Gilles Paboeuf, Florence Lefèvre. Proteomic and morphological characterization of isolated rainbow trout adipocytes originating from various fat tissues. Congrès de la Société Française d'Electrophorèse et d'Analyse Protéomique, Oct 2006, Saint Malo, France. 49 p. hal-02753790

**HAL Id: hal-02753790**

**<https://hal.inrae.fr/hal-02753790v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Proteomic and morphological characterization of isolated rainbow trout adipocytes originating from various fat tissues**

**Claudine Weil, N. Sabin, J. Bugeon, G. Paboeuf, F. Lefèvre**

*Equipe Croissance et Qualité de la Chair des Poissons, INRA-SCRIBE, IFR 140, Campus de Beaulieu, 35000 Rennes*

E-mail : [Claudine.Weil@rennes.inra.fr](mailto:Claudine.Weil@rennes.inra.fr)

In fish, like in other edible species, the repartition of the different fat deposits plays an important role in the determination of carcass quality and flesh organoleptic properties. In mammals, different works report regional differences in fat tissue with respect to cell size, metabolic and secretory functions. So far, in fish, data on adipose cell characteristics, in relation with the origin of fat deposit, are very scarce and concern principally their size distribution. The aim of the present work is to compare, in rainbow trout, the cellularity and proteomic profiles of isolated adipose cells originating from various fat deposits to better understand their functions. The following fat depots were studied : (i) perivisceral fat which constitutes the more important part of non edible residues, subcutaneous (ii) ventral and (iii) dorsal fat deposits which are eliminated when fish fillets are trimmed.


This work has necessitated the setting up of various methods : adipose cell dispersion, extraction and solubilisation of their proteins followed by their separation by bidimensional electrophoresis. Different conditions were tested in order to maximize each previous cited step and the following protocol was chosen. The different adipose tissues (12 g) were collected on fish with a mean body weight of 1.5 kilos and were minced before being submitted to collagenase (2mg/ml) dispersion for 1.5 hr at 12°C. After filtration of the digests through a 200 µm mesh fabric, adipocytes were collected by centrifugation (5 mn, 50g). The collected adipocytes were washed 3 times (5 mn, 50g) in DMEM medium and then, were submitted either to direct microscopy observation to determine cell diameter (with an image analyzer system (Visilog 5.4) or to protein extraction/solubilisation. The latter was performed at 18°C in a glass-glass homogeniser with 500 µl of buffer (40mM TRIS, 8M urea, 2M thiourea, CHAPS 4%, 50 mM DTT, 250 mM EDTA, 1% IPG buffer, protease inhibitors) followed by 2 centrifugations (15 000 g, 30 mn and 100 000 g, 1hr). 100 µg of protein was submitted to bidimensional electrophoresis. IEF was performed by using IPG DryStrips (24 cm, pH, 4-7 L) on IPG-Phor. The 2<sup>nd</sup> electrophoresis, SDS-PAGE 12.5 % in the format 240 mm x180 mm, was performed on EttanDALT six, followed by silver staining. Scanning of the gels allowed to detect approximately 1500 spots and work is in progress to detect spots differentially expressed in relation with the fat tissue origin of adipocytes. Preliminary results on morphological characteristics of collagenase dispersed adipose cells originating from the different fat tissues indicate that perivisceral adipocytes have a larger mean diameter compared to subcutaneous dorsal and ventral adipocytes.

In conclusion, the whole work would allow, in rainbow trout, to identify specific proteic markers of adipose cells from different fat deposits and would conduct to a better knowledge on the regulation of the regional lipid distribution.

**Acknowledgements** to N. Guittou and C. Pineau (High-Throughput Proteomics Platform OUEST-genopole®) for helpful assistance and guidance.



# Congrès de la Société Française d'Electrophorèse et d'Analyse Protéomique



Marqueurs biologiques: stratégies de recherche et challenges technologiques

**Saint-Malo**

**16-18 octobre 2006**

**Palais du Grand Large**

