



HAL
open science

Characterization of historical demographic expansions from linked microsatellite data

Miguel Navascués, Olivier Hardy

► **To cite this version:**

Miguel Navascués, Olivier Hardy. Characterization of historical demographic expansions from linked microsatellite data. Jacques Monod Conference "Evolutionary Genomics", May 2007, Roscoff, France. 2007. hal-02754333

HAL Id: hal-02754333

<https://hal.inrae.fr/hal-02754333v1>

Submitted on 1 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The mismatch distribution expanded to linked microsatellites

Miguel Navascués* & Olivier Hardy



Laboratoire Eco-Ethologie Évolutive, Faculté des Sciences, Université Libre de Bruxelles, Avenue F.D. Roosevelt 50, Campus du Solbosch, 1050 Brussels, Belgium

* Current address: Laboratoire d'Ecologie, École Normale Supérieure, 46 rue d'Ulm 75230 Paris Cedex 05, France. E-mail: m.navascues@gmail.com

The use of genetic data to make inferences on demographic dynamics is one of the common applications of population genetics. Many statistical methods employed on this task are based on coalescent theory which describes the genealogical relationships among a group of genes as a function of population demography (fig 1).

The theoretical background provided by the coalescent describes the distribution of mutations among the sample of genes under different demographic scenarios. However, the applicability of these predictions to empirical genetic data will be hampered by the mutational mechanisms that can erase or distort some of the useful information with back or parallel mutations. The inclusion of mutation models in theoretical predictions can improve the performance of statistical methods.

Li (1977) described the probability distribution for the number of mutations (j) between a pair of haplotypes evolving under a demographic expansion. Schneider & Excoffier (1999) introduced a correction on that equation that allowed describing the distribution of the number of genetic differences (i) for mutation model for DNA sequence polymorphisms (see equation).

Polymorphism on linked microsatellites (repetitive DNA sequences of short repeat motif, fig 2) is sensitive to demographic expansions (Navascués *et al.* 2006). However large number of homoplasious mutations are expected on these markers which affects the estimation of demographic parameters. In order to improve the accuracy on these estimates we have developed a correction similar to that of Schneider & Excoffier (1999) but using the stepwise mutation model to describe microsatellite evolution.

$$P(i; \theta_0, \theta_1, \tau) = \sum_{j=i}^{\infty} P(j; \theta_0, \theta_1, \tau) \times P(i; j)$$

probability distribution of the number genetic differences (i) given a demographic model $(\theta_0, \theta_1, \tau)$

probability distribution of the number of genetic differences (i) given a number of mutations (j) under some mutational model

probability distribution of the number of mutations (j) given a demographic model $(\theta_0, \theta_1, \tau)$

Data was simulated for a set of linked loci evolving under a stepwise mutation model in a population going through a demographic expansion. Four different ages (τ) for the expansion were simulated and maximum likelihood estimates were obtained for the time of expansion using a model without homoplasy [using only Li's $P(j; \theta_0, \theta_1, \tau)$] and a model accounting for homoplasy [using $P(i; \theta_0, \theta_1, \tau)$ with the correction based on the stepwise mutation model]. Table 1 shows how bias and error decrease when using the model corrected for homoplasy, specially for older expansions.

	Maximum likelihood estimator, model without homoplasy			Maximum likelihood estimator, model with homoplasy		
	mean among estimates	mean squared error	bias	mean among estimates	mean squared error	bias
Sim $\tau = 1$	0.885	0.083	-0.115	1.024	0.154	0.025
Sim $\tau = 3$	2.346	0.564	-0.654	2.856	0.493	-0.144
Sim $\tau = 5$	3.502	2.398	-1.498	4.900	0.872	-0.100
Sim $\tau = 7$	4.361	7.328	-2.639	6.784	2.725	-0.216

REFERENCES

- Li (1977) *Genetics* **85**:331–337
- Navascués *et al.* (2006) *Molecular Ecology* **15**:2691–2698
- Schneider & Excoffier (1999) *Genetics* **152**:1079–1089

ACKNOWLEDGMENTS

I am grateful to Brent Emerson & Concetta Burgarella for their support and discussion. This study was partially funded by the European Science Foundation ConGen Exchange Grant 1142




Fig 2 - WHERE TO FIND LINKED MICROSATELLITES

Linked microsatellites are typically found on the Y chromosome and on the chloroplast genome.

