



**HAL**  
open science

## The virtual fruit: towards a tool to design ideotypes for sustainable production systems

Bénédicte Quilot, Mohamed Mahmoud Ould Sidi, Abdeslam Kadrani, Michel M. Génard, Françoise F. Lescourret, Nadine Hilgert

### ► To cite this version:

Bénédicte Quilot, Mohamed Mahmoud Ould Sidi, Abdeslam Kadrani, Michel M. Génard, Françoise F. Lescourret, et al.. The virtual fruit: towards a tool to design ideotypes for sustainable production systems. 11. ESA Congress, Aug 2010, Montpellier, France. hal-02754522

**HAL Id: hal-02754522**

**<https://hal.inrae.fr/hal-02754522v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




# The virtual fruit: towards a tool to design ideotypes for sustainable production systems

Bénédicte Quilot



Mohamed Ould Sidi  
Abdeslam Kadrani  
Michel Génard  
Françoise Lescourret




Nadine Hilgert

MISTEA


Controlling storage disease is a priority  
to reduce fruit residues and increase safety

Critical question : finding the best combinations of  
genetic resources and cultural practices  
adapted to, and respectful of specific environments


- brown rot caused by *Monilinia* species, can provoke as much as 30 to 40% of crop losses
- no other alternative than chemical treatment is available : fungicide applications are generalized and occur till pre-harvest
- all cultivated peaches are more or less sensitive to brown rot
- mechanisms of resistance may be complex and largely linked to fruit characteristics including fruit craking




*Monilinia  
laxa*




**Fruit mass (g)**



**S<sub>cracking</sub> (cm<sup>2</sup>)**




Days after bloom





# Virtual fruit




Adapted from Bertin et al, 2010

Comment citer ce document :




Virtual fruit

Fruit growth variations between tree load




Cuticular cracking for different scenarii

$S_{\text{cracking}}$  (cm<sup>2</sup>)




Sucrose accumulation for different tree loads





# Sensibility analysis

55 parameters → 6 genotypic parameters involved in different processes



PARAMETERS	DEFINITION	PROCESSES	REFERENCE	MINI	MAXI
coeff.transfo1	Part of the leaves in the structural part of the leafy shoots	Vegetative structure	0.766	0.672	0.842*
MuF	Initial relative fruit growth rate	Fruit demand	0.0035	<b>0.001</b>	<b>0.01**</b>
ca1	Proportion of carbon as sucrose in phloemic sap	Sugar metabolism	0.347	<b>0.23</b>	<b>0.54***</b>
FSsat	Parameter of the relationship between stone dry mass and total fruit dry mass	Stone-pulp partition	6.78	<b>6.10</b>	<b>7.46**</b>
cut.be	Parameters of the equation of the relative elongation rate of the cuticle	Transpiration	1.26	1.134	1.386*
Lx	Conductivity of the composite membrane for water transport	Water fluxes	0.23328	<b>0.0769</b>	<b>0.667**</b>

\*  $\pm 10\%$  variations

\*\* Observed variability within BC2 population (Quilot et al 2005)

\*\*\* Extreme values from literature

Comment citer ce document :



## Criteria to optimize


CRITERIA	OPTIMISATION	MINI	MAXI
Fruit mass (g)	maximiser	50	300
Sweetness (%)	maximiser	4	20
Crack density (%)	minimiser	0	20




# 8 scenarii

CLIMATE	IRRIGATION	FRUIT LOAD
Avignon	Well irrigated (WI)	4 fruits/stem LC
		20 fruits/stem HC
	Water deficit (WD)	4 fruits/stem LC
		20 fruits/stem HC

same for Bordeaux



# NSGA-II : a multiobjective evolutionary algorithm, a Pareto approach




**Dominance relation between solutions**

**A large choice of solutions:**

**40 optimum genotypes over 4000 tested per scenario**


- AV\_WI\_LC
- AV\_WI\_HC
- AV\_WD\_LC
- AV\_WD\_HC



Comment citer ce document :



# Parameter correlations




Optimum are concentrated in the space  
No correlations between parameters

- AV\_WI\_LC
- AV\_WI\_HC
- AV\_WD\_LC
- AV\_WD\_HC



- AV\_WI\_LC    ■ AV\_WD\_LC
- AV\_WI\_HC    ■ AV\_WD\_HC



Complex interactions  
Direct influence of Lx





# Conclusions

**Specific genotypes for high crop load**  
**High variability of ideotypes for WD\_LC**



Comment citer ce document :

- explore genotypes selected in Bordeaux climate
- test the best genotypes of one scenario in the other scenarii:
  - are their genotypes good in different scenarii ?
- try to develop methods from Operations Research and compare results with those from Artificial Intelligence methods (NSGA)



Thank you for your attention