



Quantitative review and meta-analysis of the determinants of ruminant production in the tropics

Nadège Edouard, Maryline Boval

► To cite this version:

Nadège Edouard, Maryline Boval. Quantitative review and meta-analysis of the determinants of ruminant production in the tropics. International Symposium on Sustainable Animal Production in the Tropics: Farming in a Changing World (SAPT), Nov 2010, Gosier, France. Cambridge University Press, Advances in Animal Biosciences, 1 (2), 2010, International Sustainable Animal Production in the Tropics: Farming in a Changing World. hal-02755105

HAL Id: hal-02755105

<https://hal.inrae.fr/hal-02755105>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantitative review and meta-analysis of the determinants of ruminant production in the tropics

N. Edouard*, M. Boval*, D. Sauvant†

* INRA Antilles-Guyane, UR0143 URZ Unité de Recherches Zootechniques, 97170 Petit-Bourg, F.W.I.


† INRA AgroParisTech, UMR0791 MoSAR Modélisation Systémique Appliquée aux Ruminants, 75231 Paris, France.

contact: Maryline Boval, Maryline.Boval@antilles.inra.fr

Introduction and objective



Environmental issues



Ruminant production systems



Multi-function of grazing systems
⇒ Sustainable production



Food demand (meat, milk...)



In the tropics: grazing = major ruminant breeding system

However:

- Limited knowledge about the variability of production in different tropical contexts compared to temperate conditions
- What is the maximum productivity that can be achieved and in which feeding context ?



Quantitative review of the bibliography and meta-analysis

Feeding
(intake, digestibility)



Performances
(average daily gain)

animal - ressource -
environnement

Material and methods

Bibliographic search:

- First step: several species of ruminants

Keywords: tropical, grazing, average daily gain (ADG), intake/digestibility

- Second step: focused on growing cattle

(>50% of the papers) ⇒ Data set construction (9 papers), N = 68

Data set description:

ADG in g/d (mean +/- SE) / variables

Variable	N	Mean	SE	p	R ²	
Latitude	North	21	406	91	<0.001	0.15
	South	47	707	43		
Climate	Tropical Humid	19	628	65	<0.001	0.35
	Sub-Trop Humid	6	924	85		
	Semi Arid	33	696	55		
	Warm Arid	10	132	96		
Season	Rainy	35	746	52	<0.01	0.14
	Dry	33	474	64		
Breed	Mixed	55	715	41	<0.001	0.33
	Indian Zebu	13	188	80		
Forage	Grass	29	448	70	<0.001	0.16
	Grass + Legume	39	738	49		

Statistical analyses (ANOVAs):

- Intra-study variability determinants: experimental conditions (year, season)

- Inter-studies variability determinants (on the previous model residuals)


=> animal, resource, environment factors

Results

Focus on ADG ~ digestible Dry Matter Intake (dDMI)

- Intra-study variability determinants: experimental conditions (CodeExpe)

Quadratic Model (GLM): $ADG = f(dDMI + dDMI^2 + CodeExpe)$



Predict ADG = -1207.1 + 184.16 dDMI - 4.196 dDMI², p<0.001, R² = 0.93, RSE = 124g


⇒ dDMI & experimental cond. explain a large part of intra-study ADG variations

- Inter-studies variability determinants (on the previous model residuals)

Best model: (Quadratic Model - GLM)

$ADG_{obs-pred} = f(\% \text{legume in the diet} + \% \text{legume in the diet}^2 + \text{Season} + \text{Climate})$
p<0.01, R² = 0.71, RSE = 156g

⇒ Forage and environmental factors seem to be major determinants of inter-studies dDMI~ADG variations



⚠ Some factors (eg season) appear both among intra and inter-studies determinants making model interpretations tricky

Conclusion

The great variability of ADG observed in tropical conditions illustrates the flexibility to improve animal production from tropical pastures: young cattle having same digestible intakes can grow 4 times more than others, in response to climatic and resource variations.

This work will contribute to better quantify the relationship between ADG and feeding, even if few data are available, and should offer the opportunity to identify appropriate strategies for animal management in tropical pastures.

