



**HAL**  
open science

## **Increased levels of serum IGF-1 can rescue the skeletal impairment of female IGF-1 null mice but are insufficient for the rescue of the male skeleton**

Sébastien Elis, Hayden-William Courtland, Vingjie Wu, Hui Sun, Valerie Williams, Karl Jepsen, Shoshana Yakar

### ► **To cite this version:**

Sébastien Elis, Hayden-William Courtland, Vingjie Wu, Hui Sun, Valerie Williams, et al.. Increased levels of serum IGF-1 can rescue the skeletal impairment of female IGF-1 null mice but are insufficient for the rescue of the male skeleton. 31. Annual Meeting of the American Society for Bone and Mineral Research, Sep 2009, Denver, United States. , Journal of Bone and Mineral Research, 24 (Supplement 1), 2009, Thirty-first Annual Meeting of the American Society for Bone and Mineral Research. hal-02755337

**HAL Id: hal-02755337**

**<https://hal.inrae.fr/hal-02755337v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Increased levels of serum IGF-1 can rescue the skeletal impairment of both male and female IGF-1 null mice.

Sebastien Elis, Yingjie Wu, Hayden-William Courtland, Hui Sun, Valerie Williams, Karl Jepsen and Shoshana Yakar

Division of Endocrinology, Diabetes and Bone disease, Mount Sinai School of Medicine, New York, NY 10029


## BACKGROUND

IGF-1 acts on the growing skeleton in an endocrine and autocrine/paracrine manner. IGF-1 null mice exhibit significant impairment of skeletal growth and development. Here, we studied whether increased levels of endocrine (serum) IGF-1 can rescue the severe skeletal phenotype of both male and female IGF-1 null mice. We performed skeletal analyses of three mouse models: 1) control mice, which express normal levels of autocrine/paracrine and endocrine IGF-1, 2) mice which express autocrine/paracrine IGF-1 as in control, but also overexpress Hepatic IGF-1 transgene (HIT), and 3) IGF-1 null mice that overexpress the Hepatic IGF-1 transgene (KO-HIT) and, thus, overexpress endocrine IGF-1.

## AIM


This study aimed at defining the skeletal response to increased levels of serum IGF-1 in the presence or absence of tissue IGF-1.

## MOUSE MODELS




We performed longitudinal analyses of three mouse models: 1) control mice, which express normal levels of autocrine/paracrine and endocrine IGF-1; 2) Hepatic IGF-1 transgenic (HIT) mice, which express normal levels of autocrine/paracrine IGF-1 but overexpress endocrine IGF-1 and 3) IGF-1 null mice, which do not express autocrine/paracrine IGF-1 but overexpress endocrine IGF-1 (KO-HIT)

## RESULTS




IGF-1 null mice have severe growth retardation and exhibit 70% reduction in body weight throughout growth. However, elevations in serum IGF-1 levels in IGF-1 null mice where tissue IGF-1 expression is blunted (KO-HIT mice) restore body weight to controls in both genders. Furthermore, when tissue IGF-1 expression is normal, elevations in serum IGF-1 levels (HIT mice) increase in body weight of both genders.




# : HIT significantly different from Control \* : KO-HIT significantly different from Control

Increases in serum IGF-1 levels enhance skeletal growth (HIT mice) in both genders, indicated by increase in total cross-sectional area, cortical area, cortical thickness and relative cortical area (RCA). When tissue IGF-1 expression is blunted, elevations in serum IGF-1 (KO-HIT mice) fail to restore those skeletal properties early postnatally, evident by decreases in total cross-sectional area, cortical area and thickness at 4 weeks of age. Nonetheless, at 8 weeks of age KO-HIT mice show catch up growth and their skeletal traits are similar to controls. Of note, trabecular architecture, assessed at the distal femur, did not reveal differences between the groups at 4, 8 or 16 weeks of age in both males and females.

Control mice: local IGF1 normal + endocrine IGF1 normal  
HIT mice : local IGF1 normal + endocrine IGF1 increased  
KO-HIT mice : no local IGF1 + endocrine IGF1 increased



Elevations in serum IGF-1 lead to increased transversal bone growth rate (total cross-sectional area and cortical area) during puberty (4-8 weeks of age) in both HIT and KO-HIT male and female mice. However, while control and HIT mice "cease" transversal bone growth after puberty (8-16 weeks), KO-HIT females show prolongation of transversal bone growth.



In the absence of tissue IGF-1, elevations in serum IGF-1 levels (KO-HIT mice) diminish sex-related differences in skeletal properties. At 16 weeks of age total cross-sectional area, cortical area and marrow area in control and HIT mice differ between males and females. In contrast, femoral parameters of KO-HIT male and female at 16 weeks of age are indistinguishable.

## CONCLUSIONS

\*The KO-HIT model show that elevations in serum IGF-1 are insufficient to restore body weight and skeletal size early postnatally.  
\*During puberty and adult age, elevations in serum IGF-1 result in increased transversal bone growth rate leading to restoration of body weight and skeletal size, despite blunted tissue IGF-1 expression.  
\* When tissue IGF-1 expression is blunted, elevations in serum IGF-1 (KO-HIT mice) diminish sex-related differences in skeletal properties.

## GRANTS

National Institutes of Health : R01AR055141- R01AR054919