



**HAL**  
open science

## Genetic and physiological determinants controlling cadmium accumulation in lettuce (*Lactuca sativa*)

Walid Zorrig, C. Sarrobert, Aïda Rouached, Brigitte B. Maisonneuve, Jean-Claude Davidian, Chedly Abdelly, Pierre Berthomieu

### ► To cite this version:

Walid Zorrig, C. Sarrobert, Aïda Rouached, Brigitte B. Maisonneuve, Jean-Claude Davidian, et al.. Genetic and physiological determinants controlling cadmium accumulation in lettuce (*Lactuca sativa*). 8. Colloque National de la SFBV, Jul 2009, Strasbourg, France. 2009, 8ème Colloque National de la SFBV8th SFBV Meeting. hal-02755969

**HAL Id: hal-02755969**

**<https://hal.inrae.fr/hal-02755969v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Société Française de Biologie Végétale SFBV**  
<http://sfbv.snv.jussieu.fr/>



# **8ème Colloque National de la SFBV 8th SFBV Meeting**

## **STRASBOURG**

### **8-10 July 2009**

#### **Invited speakers**

**Harro Bouwmeester, Lucia Colombo,  
Yves Gibon, José Gualberto, Claudia Köhler,  
Alison Mallory, Lionel Navarro,  
Karl Oparka, Holger Puchta,  
Catherine Rameau**



# 8th National Meeting of the SFBV STRASBOURG 8-10 July 2009

## National Scientific Committee

**Anis Limami**, Professeur à l'Université d'Angers, Président de la SFBV

**Thierry Langin**, Directeur de l'Institut de Biotechnologie des Plantes d'Orsay, Vice-président de la SFBV

**Pascal Genschik**, Directeur de l'Institut de Biologie Moléculaire des Plantes de Strasbourg

**Jean Masson**, Président du Centre INRA de Colmar

**Michel Rohmer**, Directeur du Département Synthèse, biosynthèse et activité des biomolécules du CNRS à Strasbourg, Membre de l'Académie des Sciences

## Local Scientific Committee

Thomas Bach

Andréa Hemmerlin

Géraldine Bonnard

Michel Legrand

Marie-Edith Chabouté

Jean Molinier

Laurence Drouard

Christophe Ritzenthaler

Manfred Heinlein

Arp Schnittger

Thierry Heitz

Véronique Ziegler-Graff

## Organizing Committee

Danièle Werck-Reichhart, Chair

Lydie Thiriet, Secrétaire de l'IBMP

Christine Guibert, Cellule Congrès de l'Université de Strasbourg

Gisèle Cordillot, Secrétaire administrative de la SFBV

Martine Miquel, Trésorière de la SFBV

Comment citer ce document :

Zorrig, W., Sarrobert, C., Rouached, A., Maisonneuve, B., Davidian, J.-C., Abdelly, C., Berthomieu, P.(2009). Genetic and physiological determinants controlling cadmium accumulation in lettuce (*Lactuca sativa*). In: 8ème Colloque National de la SFBV8th SFBV Meeting (p. 2.3). Presented at 8. Colloque National de la SFBV. Strasbourg. FRA (2009-07-08 - 2009-07-10).

## Genetic and Physiological determinants controlling cadmium accumulation in lettuce (*Lactuca sativa*)

Zorrig W.<sup>(1,2)</sup>, Sarrobert C.<sup>(3)</sup>, Rouached A.<sup>(1)</sup>, Maisonneuve B.<sup>(4)</sup>, Davidian J.C.<sup>(1)</sup>, Abdelly C.<sup>(2)</sup>, **Berthomieu P.**<sup>(1)</sup>

(1) *Biochimie & Physiologie Moléculaire des Plantes, UMR CNRS-INRA-Université Montpellier II-Montpellier SupAgro, Place Viala, 34060 Montpellier Cédex 1, France*

(2) *Laboratoire d'Adaptation Des Plantes aux Stress Abiotiques, Centre de Biotechnologie, BP 901, Hammam-Lif 2050, Tunisie*

(3) *Groupe de Recherches Appliquées en Phytotechnologie, DEVM, CEN Cadarache, 13108 St Paul les Durance, France*

(4) *Génétique et Amélioration des Fruits et Légumes, UR INRA, 1052 Domaine St Maurice, BP 94, 84143 Montfavet Cédex, France*

correspondence : [berthomieu@supagro.inra.fr](mailto:berthomieu@supagro.inra.fr)

Cadmium is an extremely toxic pollutant causing a great number of diseases. It is a widespread metal whose concentration rises in cultivated soils. Food or feed crops are thus exposed to it, which ultimately causes potential sanitary problems as a consequence of the entry of cadmium in the food chain. Among the vegetable species used for food, lettuce shows one of the highest capacities to accumulate cadmium. Our objectives are to characterize mechanisms controlling cadmium accumulation in this species and identify the corresponding genetic determinants. The long-term goal is the breeding for cadmium under-accumulation in lettuce.

A phenotypical analysis of cadmium accumulation was carried out on 18 lettuce genotypes representing the genetic diversity of the species. A great variability of response was revealed, both with respect to cadmium accumulation and to cadmium translocation from roots to shoot. The lettuce genotypes displayed independent variations for both traits, and also between these traits and cadmium tolerance. In contrast, a very strong positive correlation linked cadmium and zinc accumulation.

Increasing calcium and iron concentration in the culture medium had a protective effect against cadmium accumulation, however this characteristic did not discriminate the lettuce genotypes presenting extreme cadmium accumulation phenotypes. Interestingly, experiments measuring <sup>109</sup>Cd influx in roots, <sup>109</sup>Cd efflux from roots and <sup>109</sup>Cd translocation from roots to shoots revealed that the genotype displaying the least cadmium accumulation could be discriminated from the genotype displaying the highest one by its markedly increased ability to efflux cadmium from the roots to the culture medium.

Progeny analysis from crosses between the genotypes displaying extreme performances for cadmium accumulation, cadmium translocation from roots to shoots and cadmium tolerance revealed that none of these traits was supported by a single genetic determinism. QTL mapping is under development. Surprisingly, the genetic analysis demonstrated that the ability to limit cadmium accumulation (and thus probably to increase cadmium efflux from the roots) was recessive.

In conclusion, phenotypic analysis of cadmium accumulation in a large set of lettuce genotypes revealed that a major determinant limiting cadmium accumulation in this species is the ability to efflux cadmium from the root to the culture medium.