



HAL
open science

What do interception data really tell us about actual insect invasions in Europe?

Alain Roques, Marc Kenis, Wolfgang Rabitsch, Marie-Anne Auger-Rozenberg

► **To cite this version:**

Alain Roques, Marc Kenis, Wolfgang Rabitsch, Marie-Anne Auger-Rozenberg. What do interception data really tell us about actual insect invasions in Europe?. 11. European Ecological Congress; 38. Annual Conference of the Ecological Society of Germany, Austria, and Switzerland; EURECO-GFOE 2008 "Biodiversity in an Ecosystem Context", Sep 2008, Leipzig, Germany. hal-02756126

HAL Id: hal-02756126

<https://hal.inrae.fr/hal-02756126>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

What do interception data really tell us about actual insect invasions in Europe ?

Alain Roques¹, Marc Kenis², Wolfgang Rabitsch³, Marie-Anne Auger-Rozenberg¹

¹INRA UR0633, Station de Zoologie Forestière, 45075 Orléans, France ; alain.roques@orleans.inra.fr.

²CABI Europe-Switzerland, 2800 Delémont, Switzerland

³Biodiversity and Nature Conservation, 1090 Wien, Austria

The 2005-2008 European project DAISIE (*Delivering Alien Invasive Species Inventory in Europe*) aimed at delivering the first continental inventory of the alien species of animals and plants already established in Europe. It revealed that alien insect species are establishing in Europe at a mean rate of 15.9 species per year since 2000 whereas this rate was ca. twice less 25 years ago (9.5 species/ year). A major part of these species have phytophagous habits and arrived accidentally with human activities, less than 10 % being parasites/ predators deliberately introduced for biological control. It was therefore interesting to compare these data with the interceptions reported by the national quarantine services of the European countries, which essentially concern phytophagous pests registered on quarantine lists (A1 and A2).

A total of 238 alien species were identified at species level in the notifications of non-compliance reported by EPPO for the European countries for the period 1995-2005. Among the 1315 alien insect species today considered as established in Europe, a total of 183 species of which 114 phytophagous have been recorded to have established during the same period. Only 7 of these species (i.e., 6.1%) have been intercepted before their establishment.

Although the insects arriving from Asia were dominant in both interceptions and establishments, representing 38.2 and 35% of the species respectively, the taxonomic composition of the two groups significantly differed. Coleoptera (30.6% without any dominant family), followed by Hemiptera (25.7% with a dominance of aphids and scales) largely dominate the alien species established during 1995-2005. In contrast, the quarantine interceptions during the same period mostly consisted of Diptera (30.7% of which 66.7% of agromyzids) and Hemiptera (30.0% among which 82.7% of aleyrodids), far above Coleoptera (17.8%). Overall, the taxa including small-sized sucking insects appeared to be significantly more represented in the entomofauna which established than in the interceptions. A comparison of the biological traits shown by the species which were intercepted but never established with these of the species which were never intercepted but established is under progress and will be presented at the meeting.