



HAL
open science

Ectomycorrhizal fungi and mobilisation of organic phosphorus from forest soil: Novel data and actual role

Claude C. Plassard, Julien Louche, Muhammad Arif Ali, Francois Xavier Sauvage, Elvira Legname, M. Duchemin

► To cite this version:

Claude C. Plassard, Julien Louche, Muhammad Arif Ali, Francois Xavier Sauvage, Elvira Legname, et al.. Ectomycorrhizal fungi and mobilisation of organic phosphorus from forest soil: Novel data and actual role. 19. Goldschmidt Conference, Jun 2009, Davos, Switzerland. pp.A1033. hal-02756361

HAL Id: hal-02756361

<https://hal.inrae.fr/hal-02756361v1>

Submitted on 9 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC0 - Public Domain Dedication 4.0 International License

Ectomycorrhizal fungi and mobilisation of organic phosphorus from forest soil: Novel data and actual role

C. PLASSARD^{1*}, J. LOUCHE¹, M. ARIF ALI¹,
F.-X. SAUVAGE², E. LEGNAME¹ AND M. DUCHEMIN^{1,2,3}

¹INRA/IRD/SupAgro, UMR Eco&Sols, 34060 Montpellier
Cedex 01 – France

(* correspondence: plassard@supagro.inra.fr)

²NRA/Supagro/UM1, UMR SPO, 34060 Montpellier Cedex
01 – France

It has been demonstrated that forest soils contain a high proportion of phosphorus (P) in organic form (Po) that is represented by phosphomonoesters (such as glucose1-P, ATP, phytic acid, ...) or phosphodiester (such as nucleic acids or phospholipids). To be used by trees, the phosphate group (Pi) must be released by the hydrolysis of the strong ester bond by enzymes that are phosphatases (Pases). We are currently working on the hypothesis that phosphatases released by ectomycorrhizal (ECM) fungi play an important role in the recycling of P in forest soil, thus contributing to plant P nutrition. We addressed this question in the context of spodosols of “Landes de Gascogne” as these soils, ensuring around 40% of conifer wood production in France, are very poor in Pi and high in Po contents relative to total P contents, making P limiting for maritime pine (*Pinus pinaster*) growth.

We worked on (i) *Hebeloma cylindrosporum*, a model fungal ECM species able to release huge amounts of Pase activity when grown *in vitro* in P-starved conditions, and (ii) on native ECM species. We found that *H. cylindrosporum* released 4 acid Pase isoforms able to hydrolyse a broad range of phosphate monoesters and to a low extent, the phosphodiester bis-pNPP. However, the efficiency of each isoform to release Pi from Po extracted from spodosols was low. The role of native species was studied by cultivating *P. pinaster* seedlings in intact soil samples from plots of different ages or fertilisation design. Despite a high pNPPase activity in ECM tips, P mobilisation from soil Po, measured by plant P accumulation, was very low in those soils with the lowest mineral P availability and the highest Po concentrations. Surprisingly, we found a significant decrease in Po fractions in soils that were annually fertilised and irrigated. Finally, based on our findings, we will propose a novel scheme of the possible role of ECM fungal Pase in Po mobilisation from forest soil.