

## Endocrine regulation of cell proliferation and apoptosis in the esophagus of euryhaline fishes

H. Takahashi, C. Takagi, Patrick Prunet, T. Kitahashi, S. Kajimura, T.  
Hirano, E.G. Grau, T. Sakamoto

► **To cite this version:**

H. Takahashi, C. Takagi, Patrick Prunet, T. Kitahashi, S. Kajimura, et al.. Endocrine regulation of cell proliferation and apoptosis in the esophagus of euryhaline fishes. 23. Conference of European Comparative Endocrinologists, Aug 2006, Manchester, United Kingdom. 2006. hal-02756582

**HAL Id: hal-02756582**

**<https://hal.inrae.fr/hal-02756582>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

23<sup>rd</sup> CONFERENCE OF EUROPEAN COMPARATIVE  
ENDOCRINOLOGISTS

29 August – 2 September 2006

Manchester (UK)

**Abstracts Book**

### **ESCE Council**

Eric Roubos, The Netherlands, President  
Geoff Coast, United Kingdom, Vice-President  
Horst-Werner Korf, Germany, Secretary-Treasurer  
Arnold de Loof, Belgium, Hon. Vice-President  
Hubert Vaudry, France, Hon. Vice-President  
Richard Balment, United Kingdom  
Manuel Carrillo, Spain  
Valér Csernus, Hungary  
Elisabeth Eppler, Switzerland  
Sylvia Fasano, Italy  
Dalibor Kodrik, Czech Republic  
Dan Larhammar, Sweden  
Daniel Soye, France  
Hervé Tostivint, France  
Jozef Vanden Broeck, Belgium

### **ORGANISING COMMITTEES**

#### ***International Scientific Committee***

M Vallarino  
Eric Roubos  
Geoff Coast  
Horst-Werner Korf  
Manuel Carrillo  
Valér Csernus  
Elisabeth Eppler  
Sylvia Fasano  
Dalibor Kodrik  
Dan Larhammar  
Daniel Soye  
Hervé Tostivint  
Jozef Vanden Broeck  
Arnold de Loof  
Hubert Vaudry  
Richard Balment  
Robert Does

#### ***National Advisory Board***

Ann Brown  
Melody Clark  
Graham Goldsworthy  
Ian Henderson  
Andrew Loudon  
Richard Balment (*Chairperson*)

#### ***Local Organising Committee***

Julian Davis  
Wei-qun Lu  
Simon Luckman  
Hugh Piggins  
Cathy McCrohan  
Anne White  
Richard Balment (*Chairperson*)

### **HONORARY CONFERENCE CHAIRMAN**

Ian W. Henderson

### **SPONSORS**

AstraZeneca  
British Society for Neuroendocrinology  
Datasci  
Eppendorf  
Eurogentec  
Marketing Manchester  
Society for Endocrinology  
Society for Experimental Biology  
Stratagene  
The Physiological Society  
The University of Manchester  
Wolf Laboratories Ltd

P086

**EFFECTS OF HYPEROSMOTIC STRESS AND ARGININE VASOTOCIN ON EXPRESSION OF AN UREA TRANSPORTER IN THE KIDNEY AND URINARY BLADDER OF THE MARINE TOAD**

Konno, N., Hyodo, S., Matsuda, K., Uchiyama, M.

Department of Life &amp; Environmental Science, Graduate school of Science &amp; Engineering, University of Toyama, Japan. Laboratory of Physiology, Ocean Research Institute, University of Tokyo, Japan.

Email: d0470301@ems.u-toyama.ac.jp

Anuran amphibians accumulate a large amount of urea in their body fluids and maintain hyper-osmolality to tolerate a severe dehydration under dry and hyper-saline environments. To clarify the mechanisms of urea retention and effects of arginine vasotocin (AVT) on urea reabsorption, we examined molecular structure, distribution and functional expression of the urea transporter (UT) following exposure to dry and hyper-saline conditions in the kidney and urinary bladder of the marine toad (*Bufo marinus*). Bufo UT cDNA cloned from the kidney encodes a 390 amino acid residue protein, and the mRNA and protein were abundantly expressed in the kidney and urinary bladder. Immunohistochemically, the UT is localized at the apical membrane of epithelial cells along the early distal tubule, known as the diluting segment, in the kidney. When toads were acclimated to dry and hyper-saline environments for 7 days, plasma concentrations of urea and AVT were significantly elevated, and there were significant correlations among the plasma concentrations of urea and AVT, and the level of Bufo UT mRNA expression in both the kidney and urinary bladder. These results suggest that the Bufo UT probably contributes to urea reabsorption in the kidney and urinary bladder in response to hyperosmotic stresses and an increase of circulating plasma AVT level.

P087

**ENDOCRINE REGULATION OF CELL PROLIFERATION AND APOPTOSIS IN THE ESOPHAGUS OF EURYHALINE FISHES**

Takahashi, H., Takagi, C., Prunet, P., Kitahashi, T., Kajimura, S., Hirano, T., Grau, E.G., Sakamoto, T.

Ushimado Marine Laboratory, Graduate School of Natural Science and Technology, Okayama University, Japan. Group on Fish Physiology of Stress and Adaptation, INRA SCRIBE, France. Department of Physiology, Nippon Medical School, Japan. Dana-Farber Cancer Institute, Harvard Medical School, USA. Hawaii Institute of Marine Biology, University of Hawaii, USA.

Email: dns18408@cc.okayama-u.ac.jp

In the seawater (SW)-acclimated euryhaline fishes, the ion/water permeability of the gastrointestinal tract is generally greater than that of freshwater (FW)-acclimated fish. The esophageal epithelium of SW fishes is simple columnar in form, whereas that of FW fishes is stratified. To understand how environmental and hormonal stimuli affect the gastrointestinal-tract differentiation, we examined the esophageal cell turnover of Mozambique tilapia and amphibious mudskipper transferred to various environments. In both species, increased apoptosis was found throughout the esophageal epithelium during SW acclimation, whereas cell proliferation occurred randomly over the epithelium in FW. There was no significant change in the mudskipper kept out of water. In vivo treatment of the mudskipper with prolactin (PRL) induced the epithelial cell proliferation. Triiodothyronine or a teleostean mineralocorticoid candidate, 11-deoxycorticosterone, showed no significant effect on cell turnover. Cortisol stimulated both epithelial cell proliferation and apoptosis. In the tilapia, glucocorticoid receptor (GR) was observed in the epithelia including the apoptotic and proliferating cells, whereas the PRL receptor expression was high in FW and seemed to be localized in the proliferating cells. Thus, it is likely that in the esophageal epithelium of these fishes, PRL and cortisol induce cell proliferation in FW, whereas cortisol-GR stimulates apoptosis during SW acclimation.