



**HAL**  
open science

## **Analysis of the resistance to PPV in a F1 progeny derived from the apricot cultivars 'Goldrich' and 'Moniqui'. Identification of the QTLs involved**

Patrick Lambert, Guillaume Roch, Aurore Graillat, Jean Leonetti, Eric Martin, Anne Marie Ferréol, Alain Blanc, Jean Marc Audergon

### ► To cite this version:

Patrick Lambert, Guillaume Roch, Aurore Graillat, Jean Leonetti, Eric Martin, et al.. Analysis of the resistance to PPV in a F1 progeny derived from the apricot cultivars 'Goldrich' and 'Moniqui'. Identification of the QTLs involved. International symposium on Plum pox virus; SharCo research workshop, Sep 2010, Sofia, Bulgaria. hal-02756753

**HAL Id: hal-02756753**

**<https://hal.inrae.fr/hal-02756753>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Analysis of the resistance to PPV in a F1 progeny derived from the apricot cultivars ‘Goldrich’ and ‘Moniqui’ - Identification of the QTLs involved**

**P. Lambert, G. Roch, A. Graillat, J. Leonetti, E. Martin, A.M. Ferreol, A. Blanc and J. M. Audergon**

INRA– UR1052 UGAFL – Domaine St Maurice – BP94 – F84143 Montfavet cedex (France)  
Email: [patrick.lambert@avignon.inra.fr](mailto:patrick.lambert@avignon.inra.fr)

**Keywords:** *Prunus armeniaca*, sharka, *Plum pox virus*, genetic resistance, QTL, genetic map

Within the framework of the SharCo project, the genetic factors controlling resistance to PPV in the apricot ‘Goldrich’ cultivar were investigated using a F1 progeny. The aim was to complete our knowledge on the genetic determinism of the resistance carried by ‘Goldrich’ and to compare it with other apricot resistance sources.

A progeny of 180 hybrids derived from a cross between ‘Goldrich’ (resistant) and ‘Moniqui’ (susceptible) has been maintained and two genetic maps composed of 120 and 94 SSR markers widely distributed throughout the genome were then constructed for ‘Goldrich’ and ‘Moniqui’, respectively.

Concurrently, the susceptibility vs resistance to sharka disease, after inoculation with PPV-M, was scored in an insect-proof greenhouse, on all the F1 seedlings, and the parents, using 5 replicates for each. The observations were performed during each vegetative cycle onto both the rootstock and the scion. They were completed by a systematic ELISA test. A minimum of 5 vegetative cycles were conducted for each genotype.

The data obtained are consistent with those reported in previous studies on ‘Goldrich’ or other resistant cultivars. However, while ‘Goldrich’ carries a major resistant factor located on LG1 which is associated with symptom intensity and localization, an additional QTL was detected on LG7 and found associated with the so-called latency period, defined as the period (expressed in number of vegetative cycles) before the inoculation and the appearance of the first symptoms. ‘Moniqui’ carries two susceptible factors located on LG2 associated with the latency period before symptom appearance and the persistence of the observed symptoms.