



HAL
open science

Effect of phytobiotic blends on digestive microbiota of broiler chickens in two rearing densities

Sarah S. Guardia, F. Recoquillay, Herve H. Juin, Michel Lessire, Maryse Leconte, Patricia Rideaud, Carole C. Moreau-Vauzelle, Christele Dupont, J.F. Guillot, Irène Gabriel

► To cite this version:

Sarah S. Guardia, F. Recoquillay, Herve H. Juin, Michel Lessire, Maryse Leconte, et al.. Effect of phytobiotic blends on digestive microbiota of broiler chickens in two rearing densities. 7. Joint Symposium Gut Microbiology: new insights into gut microbial ecosystems, Jun 2010, Aberdeen, United Kingdom. Rowett-INRA, 2010, 7th Joint Symposium Gut Microbiology: new insights into gut microbial ecosystems. hal-02757050

HAL Id: hal-02757050

<https://hal.inrae.fr/hal-02757050>

Submitted on 5 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title :

Effect of phytobiotic blends on digestive microbiota of broiler chickens in two rearing densities

Abstract:

Sarah Guardia¹, François Recoquillay², Hervé Juin³, Michel Lessire¹, Maryse Leconte¹, Patricia Rideaud³, Carole Moreau-Vauzelle³, Christel Dupont³, Jean François Guillot⁴, Irène Gabriel¹

1-INRA, UR 83, URA, 37380 Nouzilly, FRANCE; 2-PHYTOSYNTHESE, 57, avenue Jean Jaures, Z.I. de Mozac Volvic, BP 50100, 63203 Riom, FRANCE; 3-INRA, UE 1206, UEASM, 17700 Surgeres, FRANCE; 4-I.U.T DE TOURS, Microbiology laboratory, 29 rue du Pont-Volant, 37082 Tours, France.

In poultry, phytobiotics are used in order to improve host growth performances. Several modes of action could be involved, the most probable being by effect on microbiota, immunity, antioxidant capacity, and digestive area. The effect on growth performances of these additives can show variations partly due to variation in breeding conditions .

In this study, the digestive microbiota and the performances of chicken fed with or without phytobiotics were studied according to rearing densities. Three experimental dietary treatments were performed: a control diet, a Exp1 diet incorporating a phytobiotic blend from d22 with an antimicrobial effect, and a Exp2 diet incorporating the previous phytobiotic blend from d10 preceded by another phytobiotic blend with antioxidant and immunomodulator effects. Body weights were recorded until d39. Digestive microbiota was analysed by TTGE in digestive content of crop, ileum and caeca at 3 and 6 weeks. The effect of the phytobiotics on the microbiota was assessed by ANOSIM performed on the Pearson distance matrix between the TTGE profiles.

Phytobiotics used during this study were efficient in promoting chickens growth from 3 to 6 weeks, specially at normal density. Moreover, changes were observed in digestive microbiota of chickens fed with phytobiotics, specially at 6 weeks. These changes were more important at high rearing density. With Exp1, main changes were observed in the caeca, whereas with Exp2, changes were observed in ileum and caeca. Thus, digestive microbiota may be one factor involved in the growth promoting effect of these phytobiotics, although probably not the only factor.