

**Is water use efficiency a useful and relevant trait for adaptation to local conditions and for breeding? Present knowledge and research needs**

Erwin Dreyer, Oliver Brendel, Didier Le Thiec, Nicolas Marron, Franck Brignolas

► **To cite this version:**

Erwin Dreyer, Oliver Brendel, Didier Le Thiec, Nicolas Marron, Franck Brignolas. Is water use efficiency a useful and relevant trait for adaptation to local conditions and for breeding? Present knowledge and research needs. Forest ecosystem genomics and adaptation, Jun 2010, San Lorenzo de El Escorial, Spain. n.p. hal-02757075

**HAL Id: hal-02757075**

**<https://hal.inrae.fr/hal-02757075>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## IS WATER USE EFFICIENCY A USEFUL AND RELEVANT TRAIT FOR ADAPTATION TO LOCAL CONDITIONS AND FOR BREEDING? PRESENT KNOWLEDGE AND RESEARCH NEEDS

*Erwin Dreyer<sup>1</sup>, Oliver Brendel<sup>1</sup>, Didier Le Thiec<sup>1</sup>, Nicolas Marron<sup>1</sup> and Franck Brignolas<sup>2</sup>*

<sup>1</sup> INRA-Nancy, Champenoux, France. Corresponding author email: dreyer@nancy.inra.fr

<sup>2</sup> Université d'Orléans, Orléans, France

---

Differences in water use efficiency (WUE) have been extensively investigated in trees using carbon-13 in leaf biomass as a surrogate. We now have evidence of considerable diversity for this trait in several tree species, and valuable insight into the genetic control of this diversity is available. Nevertheless, two main questions remain open: 1) Is the observed diversity of WUE in natural populations neutral, or is WUE subject to selective pressure? 2) Is enhanced WUE a valuable target for breeding in plantation species such as poplar or pines? Based on recent evidence obtained in natural populations of oaks and beech, as well as in clonal plantations of poplar, we discuss these two questions and provide some answers, as well as outline further research needs. Briefly, there is currently little evidence from natural populations that the observed genetic diversity in <sup>13</sup>C discrimination and WUE is organized along clear patterns related for instance to soil water availability. We therefore urgently need more data on the genetic variability in <sup>13</sup>C discrimination at local scales, i.e., common garden experiments. Breeding for enhanced WUE will not produce genotypes with enhanced drought resistance, but could possibly increase the sustainability of plantation forestry in regions where water availability is a major issue.

**Keywords:** <sup>13</sup>C, functional trait, isotopic discrimination, transpiration efficiency, water availability