



HAL
open science

Exploration of late blight quantitative resistance in potato diversity: QTL meta-analysis including the novel resistance sources *Solanum sparsiplum* and *Solanum spegazzinii*

Sarah Danan, Jean-Eric Chauvin, Bernard Caromel, Jean-Baptiste Veyrieras, Jean-Paul Dantec, Roland Pellé, Nadia Lama, Véronique Lefebvre

► To cite this version:

Sarah Danan, Jean-Eric Chauvin, Bernard Caromel, Jean-Baptiste Veyrieras, Jean-Paul Dantec, et al.. Exploration of late blight quantitative resistance in potato diversity: QTL meta-analysis including the novel resistance sources *Solanum sparsiplum* and *Solanum spegazzinii*. 5. Solanaceae Genome Workshop, Oct 2008, Cologne, Germany. hal-02757139

HAL Id: hal-02757139

<https://hal.inrae.fr/hal-02757139>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#154: Exploration of late blight quantitative resistance in potato diversity: QTL meta-analysis including the novel resistance sources *Solanum sparsipilum* and *Solanum spegazzinii*

S. Danan¹, J. Chauvin², B. Caromel³, J. Veyrieras⁴, J. Dantec⁵, R. Pellé⁵, N. Lama³, V. Lefebvre³

¹INRA, MONTFAVET, France

²INRA UMR 118 APBV, Ploudaniel, France

³INRA - UR 1052 - GAFL, Montfavet, France

⁴INRA UMR 320, Gif-sur-Yvette, France

⁵UMR 118 APBV, Ploudaniel, France

The resurgence of late blight epidemics in potato (*Solanum tuberosum*) is due to the apparition of new *Phytophthora infestans* strains, resistant to chemicals and overcoming deployed R-genes. The alternative exploitation of durable partial resistance controlled by QTLs requires to inventory resistance sources and to get a better insight of the genetic architecture and diversity of quantitative resistance to late blight in potato. Here, we report the identification of QTLs in two novel wild species and integrate the mapping results in a meta-analysis for late blight resistance in potato together with the data of 19 other published studies.

Two mapping diploid progenies were assessed with a stem assay in controlled conditions and a foliage assay in outdoor conditions. QTL analysis was performed with 16 traits describing late blight resistance on the 4 parental maps sharing common genetic markers. Considering both progenies, a total of 75 detected QTLs were mapped on 11 of the 12 potato chromosomes. High-effect QTLs were detected on chromosomes I, V, VIII and X with QTL colinearities across parental maps. Anchor literature markers highlighted colinearities with reported R-genes and QTLs.

Thanks to the QTL meta-analysis, a consensus map of potato with the projection of late blight resistance meta-QTLs was constructed. Also, markers with a high frequency occurrence among experiments were selected to constitute an advisable potato set for new genetic map construction. Results showed (i) the resistance potential of *S. sparsipilum* and *S. spegazzinii* for breeding programs, (ii) the efficiency of the stem test as a complement to the foliage test to assess quantitative resistance, (iii) close relationships between quantitative and qualitative resistance factors, and (iv) highlight the allelic diversity of late blight resistance QTLs.