



**HAL**  
open science

## Natural variation and functional analysis provide evidence for co-evolution between pepper eIF4E and potyviral Vpg

Carole Caranta, Carine Charron, Maryse Nicolaï, Jean-Luc J.-L. Gallois, Alain Palloix, Christophe Robaglia, Stéphane de Mita, Jacques David

### ► To cite this version:

Carole Caranta, Carine Charron, Maryse Nicolaï, Jean-Luc J.-L. Gallois, Alain Palloix, et al.. Natural variation and functional analysis provide evidence for co-evolution between pepper eIF4E and potyviral Vpg. 5. Solanaceae Genome Workshop, Oct 2008, Cologne, Germany. hal-02757223

**HAL Id: hal-02757223**

**<https://hal.inrae.fr/hal-02757223v1>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## #107: Natural variation and functional analysis provide evidence for co-evolution between pepper eIF4E and potyviral VPg

C. Caranta<sup>1</sup>, C. Charron<sup>1</sup>, M. Nicolai<sup>1</sup>, J. Gallois<sup>1</sup>, A. Palloix<sup>1</sup>, C. Robaglia<sup>2</sup>, S. de Mita<sup>3</sup>, J. David<sup>3</sup>

<sup>1</sup>INRA, UR1052, Unité de Génétique et Amélioration des Fruits et Légumes, Dom. Saint-Maurice, Montfavet, France

<sup>2</sup>Laboratoire de Génétique et Biophysique des Plantes, CEA-CNRS-Univ. Aix Marseille II, Faculté des Sciences de Luminy, Marseille, France

<sup>3</sup>INRA-UMR 1097, Diversité et Adaptation des Plantes Cultivées, Domaine de Melgueil, Mauguio, France

Amino acid (AA) substitutions in the eukaryotic translation initiation factor eIF4E result in recessive resistance to potyviruses in a range of plant species, including pepper and tomato [Robaglia and Caranta, 2006, Trends in Plant Science]. Correspondingly, AA changes in the viral protein VPg are responsible for the potyvirus's ability to overcome eIF4E-mediated resistance. With the aim to characterize selection pressures exerted on eIF4E genes in relation to resistance and mRNA translation functions, comparative analyses of polymorphism and functional analyses were conducted in pepper and *Arabidopsis thaliana*.

Polymorphism analyses show that *A. thaliana* eIF4E genes are very well conserved, under negative selection, similarly to house-keeping genes. Conversely, polymorphism analysis at the pvr2-eIF4E coding sequence in a worldwide sample of 25 *C. annuum* accessions identified ten allelic variants with exclusively non-synonymous variations. Resistance and genetic complementation assays demonstrated that pvr2 variants, each with signature amino acid changes, corresponded to potyvirus resistance alleles. Systematic analysis of the interactions between eIF4E proteins encoded by the ten pvr2 alleles and VPgs of virulent and avirulent Potato virus Y (PVY) and Tobacco etch virus (TEV) strains demonstrated that resistance phenotypes arose from the disruption of the interaction between eIF4E and VPg, and that viral adaptation to eIF4E-mediated resistance resulted from the restored interaction with the resistance protein. Complementation of an eIF4E knock-out yeast strain by *C. annuum* eIF4E proteins further shows that amino acid changes did not impede essential eIF4E functions. Moreover, eIF4E gene from *C. annuum* presents positive selection traces and AA under positive selection correspond to residues involved in PVY resistance. Altogether, these results argue in favour of a coevolutionary arms race between *Capsicum* eIF4E and potyviral VPg [Charron et al., 2008. Plant J].