



Contemporary seed and pollen dispersal abilities of silver fir (*Abies alba* Mill.) at range margins

Sylvie Oddou-Muratorio, Annabelle Amm, Jaroslaw Burczyk, Igor J. Chybicki, Andrzej Lewandowski, Monika Litkowiec, Christian Pichot, Etienne K. Klein

► To cite this version:

Sylvie Oddou-Muratorio, Annabelle Amm, Jaroslaw Burczyk, Igor J. Chybicki, Andrzej Lewandowski, et al.. Contemporary seed and pollen dispersal abilities of silver fir (*Abies alba* Mill.) at range margins. Forest ecosystem genomics and adaptation, Jun 2010, San Lorenzo de El Escorial, France. hal-02757428

HAL Id: hal-02757428

<https://hal.inrae.fr/hal-02757428>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTEMPORARY SEED AND POLLEN DISPERSAL ABILITIES OF SILVER FIR (*ABIES ALBA* MILL.) AT RANGE MARGINS

Sylvie Oddou-Muratorio¹, Annabel Amm¹, Jaroslaw Burczyk², Igor J. Chybicki², Andrzej Lewandowski³, Monika Litkowiec³, Christian Pichot¹ and Etienne K. Klein⁴

¹ Ecologie des Forêts Méditerranéennes, INRA, Avignon, France.

Corresponding author email: sylvie.oddou@avignon.inra.fr

² Uniwersytet Kazimierza Wielkiego, Institut Biologii Eksperymentalnej, Bydgoszcz, Poland

³ Instytut Dendrologii PAN, Kórnik, Poland

⁴ Biostatistique et Processus Spatiaux, INRA, Avignon, France

Effective gene flow from parent to offspring is a major process shaping plant population genetic diversity and evolutionary potential. However, little is still known about the extent of within-species variation in contemporary patterns of pollen and seed dispersal, and even less on ecological drivers shaping this variation. Silver fir (*Abies alba* Mill.) is a keystone tree species in low to mid-altitude European mountainous forest ecosystems, and appears to be particularly sensitive to ongoing climatic change. In this study, we investigated contemporary patterns of gene flow at southern and northern margins of the European distribution range of silver fir, using two populations in a Mediterranean mountainous forest (EVOLTREE ISS Ventoux, France) and two populations in an old-growth forest (EVOLTREE ISS Blyzin, Poland). We used spatially explicit parentage models and a combination of nuclear and chloroplast DNA microsatellite markers to estimate seed and pollen dispersal kernels in naturally established seedlings. Patterns of contemporary gene flow appeared to vary across populations. Selfing rates were higher in southern populations (average $s = 0.18$) as than in northern populations (average $s = 0.07$). Pollen dispersal also differed strongly among sites, with fat-tailed dispersal kernels and average pollen dispersal distance (δ_p) >100 m in northern populations, while southern populations showed exponential pollen dispersal kernels and restricted pollen dispersal distance ($\delta_p < 11$ m). Patterns of seed dispersal were less variable among populations, with a general tendency for exponential dispersal kernel and average seed dispersal distance ranging between 18.2 and 24.5 m. Both pollen and seed immigration rates were highly sensitive to assumptions on background pollen pool frequencies, due to low resolution of the marker set. These results indicate limited pollen production and dispersal abilities in trailing-edge populations as compared with leading-edge populations, as well as overall limited seed dispersal abilities relative to rates of ongoing and predicted climate change.

Keywords: contemporary gene flow, microsatellite, parentage analysis, pollen dispersal, seed dispersal, selfing