



HAL
open science

Genetic analysis of the resistance to downy and powdery mildews derived from cultivar Bronner

Paule Blasi, Sylvain Schnee, Sabine Wiedemann-Merdinoglu, Emilce Prado, S. Godard, Pascale Coste-Heinrich, Christine Onimus, Katia Gindro, Christophe Joseph Schneider, O. Viret, et al.

► To cite this version:

Paule Blasi, Sylvain Schnee, Sabine Wiedemann-Merdinoglu, Emilce Prado, S. Godard, et al.. Genetic analysis of the resistance to downy and powdery mildews derived from cultivar Bronner. 6. International Workshop of Grapevine Downy and Powdery Mildew, Jul 2010, Bordeaux, France. hal-02757775

HAL Id: hal-02757775

<https://hal.inrae.fr/hal-02757775v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic analysis of the resistance to downy and powdery mildews derived from cultivar Bronner

P. Blasi^a, S. Schnee^b, S. Wiedemann-Merdinoglu^a, E. Prado^a, S. Godard^b, P. Coste^a, C. Onimus^a, K. Gindro^b, C. Schneider^a, O. Viret^b, D. Merdinoglu^a

^aINRA-UDS, UMR 1131 Santé de la Vigne et Qualité du Vin, 28 rue de Herrlisheim, BP 20507, 68021 Colmar cedex, France ; ^bDépartement Fédéral de l'Economie DFE, Station de recherche Agroscope Changins-Wädenswil ACW, CP 1012, 1260 Nyon, Suisse

A wide range of pathogens threatens viticulture. Among them, downy and powdery mildews are the most important in Europe. The current strategy to control these diseases relies totally on the use of fungicides. This practice is not only expensive but also causes a slow and progressive damage to the environment. A cost-effective and environment friendly alternative to the use of chemicals is the development of varieties resistant to pathogens. All traditional European grapevine varieties are susceptible to the main pathogens. However *Vitis* species closely related to cultivated grapevine were shown to be potential sources of resistance to a wide spectrum of grapevine diseases (Boubals 1959, Staudt and Kassemeyer 1995).

The absence of private grapevine breeders in France led the INRA to design a breeding program dedicated to create new resistant varieties. The main goal of this programme is to create varieties durably resistant to downy and powdery mildews with a berry quality suitable to produce high quality wines (Merdinoglu *et al.* 2009). In order to successfully reach the double objective of high resistance efficiency and durability, the use of multiple sources of resistance was planned as soon as the project was designed. The project was developed in close interconnection with upstream research programmes which aim at understanding the genetic bases of the resistance to downy mildew derived from grapevine-related wild species by addressing four key questions: (i) exploring the diversity available in genetic resources to chose original genitors (ii) identifying and characterizing the relevant genes/QTLs to genetically improve the targeted traits, (iii) using the data acquired on genes/QTLs (position, effects) to assist the selection with markers, and (iv) assessing the durability of the identified resistance genes/QTLs.

In the presented study, we analysed the genetic determinism of the resistance to grapevine downy and powdery mildews derived from cultivar Bronner. We used two BC6 mapping populations, respectively consisting of 96 and 143 individuals from a cross between the resistant parent Bronner and an other resistant parent derived from *Muscadinia rotundifolia* var. Dearing. The two populations were screened with two SSR markers flanking Rpv1, a downy mildew resistance QTL from *Muscadinia*, in order to discard the individuals with this resistance factor and only keep the plants segregating for the resistance derived from Bronner. Resistance to downy mildew was assessed after artificial inoculation on both populations. Plants were genotyped at 65 SSR loci which allowed us to build a genetic map covering 648 cM. The interval mapping analysis revealed the presence of a major downy mildew

resistance QTL that was located on linkage group 9. This resistance factor accounted for 52% of the total phenotypic variation. Thus, we considered this QTL as a major gene.

Resistance to powdery mildew was assessed after artificial inoculation as well. The interval mapping analysis revealed the presence of two minor QTLs involved in powdery mildew resistance. The two QTLs were located on linkage group 3 and linkage group 19 and explained 8% and 16% of the total phenotypic variation, respectively.

Literature cited

- Boubals D. 1959. Contribution à l'étude des causes de la résistance des Vitacées au mildiou de la vigne et de leur mode de transmission héréditaire. *Ann. Amélior. Plant.* 9 :1-236.
- Merdinoglu D, Merdinoglu-Wiedemann S, Mestre P, Prado E, Schneider C. 2009. Apport de l'innovation variétale dans la réduction des intrants phytosanitaires au vignoble : exemple de la résistance au mildiou et à l'oïdium. *Progrès Agricole et Viticole* 126:244-247.
- Staudt G, Kassemeyer HH. 1995. Evaluation of downy mildew resistance in various accessions of wild *Vitis* species. *Vitis* 34:225-228.