



HAL
open science

Molecular mapping in *Capsicum annuum* of the tomato Ve2 gene

Lorenzo Barchi, E. Portis, Véronique Lefebvre, Alain Palloix, S. Lanteri

► **To cite this version:**

Lorenzo Barchi, E. Portis, Véronique Lefebvre, Alain Palloix, S. Lanteri. Molecular mapping in *Capsicum annuum* of the tomato Ve2 gene. 5. Solanaceae Genome Workshop, Oct 2008, Cologne, Germany. hal-02757830

HAL Id: hal-02757830

<https://hal.inrae.fr/hal-02757830v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#253: Molecular mapping in capsicum annuum of the tomato ve2 gene

L. Barchi¹, E. Portis¹, V. Lefebvre², A. Palloix², S. Lanteri¹

¹University of Turin DI.VA.P.R.A. - Plant Genetics and Breeding, Grugliasco, Italy

²INRA - UR 1052 - GAFL, Montfavet, France

Verticillium wilt is a fungal disease caused by two soil-borne pathogens: *Verticillium dahliae* and *Verticillium albo-athrum*. They are responsible for important yield and quality losses in many crops. Few strategies are at present available for the control of these pathogens. To date, tolerance to *Verticillium* wilt has been found in *Capsicum chinense*, *C. frutescens* and *C. baccatum* genotypes, while only reduced susceptibility have been identified in *C. annuum*.

In tomato, the genetic resistance is conferred by two independent genes: *Ve1* and *Ve2*, which have been mapped on the short arm of chromosome T9. We isolated sequences homologue to *Ve1* and *Ve2* in pepper, and identified their location in three previously developed intra-specific *C. annuum* maps: (i) F5 RILs 'Yolo Wonder' x 'Criollo de Morelos 334' (YC); (ii) doubled haploids 'Perennial' x 'Yolo Wonder' (PY); (iii) doubled haploids 'H3' x 'Vania' (HV).

Primers were designed on the tomato *Ve1* (AF272367) and *Ve2* (AF365929) gene sequences and applied for amplifications of pepper homologous cDNAs obtained from pepper leaves. PCR products were obtained only with *Ve2* primers. The amplified pepper 860-bp-sequence revealed 87% identity with the tomato *Ve2* gene. The full length of the pepper homologous *Ve2* gene (*CaVe2*) was then successfully isolated.

Primer pairs were designed for analyzing the allelic forms of *CaVe2* in the five parents of the 3 mapping populations. A SNP between 'Yolo Wonder' and 'Criollo de Morelos 334' was identified, and a specific co-dominant dCAPS marker developed. Another dCAPS marker was generated for PY population while a tetra-primer ARMS marker was obtained for HV population. Segregation analyses in the three mapping populations made possible to localize *CaVe2* locus on chromosome P9, in a position colinear to the tomato *Ve* loci.

Further studies are in progress to evaluate the role played by this gene in the pepper tolerance to *Verticillium*.