



HAL
open science

Caractérisation moléculaire, phénotypique et fonctionnelle d'isolats de *Trichoderma* sp. antagonistes de *Rhizoctonia solani*

Christian Steinberg, Muhammad Anees, Arne Tronsmo, Linda Gordon Hjeljord, Cécile Héraud, Véronique Edel-Hermann

► To cite this version:

Christian Steinberg, Muhammad Anees, Arne Tronsmo, Linda Gordon Hjeljord, Cécile Héraud, et al.. Caractérisation moléculaire, phénotypique et fonctionnelle d'isolats de *Trichoderma* sp. antagonistes de *Rhizoctonia solani*. Journées Jean Chevaugeon : 8. Rencontres de Phytopathologie - Mycologie, Jan 2010, Aussois, France. n.p. hal-02757858

HAL Id: hal-02757858

<https://hal.inrae.fr/hal-02757858>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Caractérisation moléculaire, phénotypique et fonctionnelle d'isolats de *Trichoderma* sp. antagonistes de *Rhizoctonia solani*.

Christian Steinberg^a, Muhammad Anees^a, Arne Tronsmo^b, Linda Gordon Hjeljord^b, Cécile Héraud^a, Véronique Edel-Hermann^a

^aINRA, Université de Bourgogne, UMR 1229 Microbiologie du sol et de l'environnement, CMSE, 17 rue Sully, BP 86510, 21065 Dijon, France; ^bDepartment of chemistry, biotechnology and food science, PO box 5003, NO-1432 Aas, Norway.

The aim of the present study was to characterize sixteen isolates of *Trichoderma* originating from a field of sugar beet where disease patches caused by *Rhizoctonia solani* were observed. The isolates were identified using both molecular and morphological characters and both approaches were found complementary. Production of water-soluble and volatile inhibitors, mycoparasitism and induced systemic resistance in plant host were investigated using *in vitro* and *in vivo* tests in disinfested and non-disinfested soils. This functional approach revealed the intraspecific diversity as well as biocontrol potential of the different isolates. Different antagonistic mechanisms were evident for different strains. The most antagonistic strain, T30 was identified as *T. gamsii*. This is the first report of an efficient antagonistic strain of *T. gamsii* being able to reduce the disease in different conditions. The ability to produce water-soluble inhibitors or coil around the hyphae of the pathogen *in vitro* was not related to the disease reduction *in vivo*. Additionally, the strains collected from the high disease areas in the field were better antagonists. The antagonistic activity was not characteristic of a species but that of a population of strains.

Key words: *Trichoderma gamsii*, *T. velutinum*, water-soluble inhibitors, volatile inhibitors, biocontrol