



HAL
open science

Nuclear microsatellites and SNPs reveal low diversity in stone pine (*Pinus pinea* L.), a genetically depauperated species

Giovanni G. Vendramin, Bruno Fady, Hadrien Lalagüe, Milvia Racchi, Federico Sebastiani, Remy R. Petit, Santiago C. Gonzalez-Martinez

► **To cite this version:**

Giovanni G. Vendramin, Bruno Fady, Hadrien Lalagüe, Milvia Racchi, Federico Sebastiani, et al.. Nuclear microsatellites and SNPs reveal low diversity in stone pine (*Pinus pinea* L.), a genetically depauperated species. Forest ecosystem genomics and adaptation, Jun 2010, San Lorenzo de El Escorial, Spain. <hal-02758127>

HAL Id: hal-02758127

<https://hal.inrae.fr/hal-02758127v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

NUCLEAR MICROSATELLITES AND SNPs REVEAL LOW DIVERSITY IN STONE PINE (*PINUS PINEA* L.), A GENETICALLY DEPAUPERATED SPECIES

Giovanni G. Vendramin¹, Bruno Fady², Hadrien Lalague^{1,2}, Milvia Racchi³, Federico Sebastiani¹, Remy J. Petit⁴ and Santiago C. González-Martínez⁵

¹ Istituto di Genetica Vegetale, Consiglio Nazionale delle Ricerche, Sesto Fiorentino, Italy.
Corresponding author email: giovanni.vendramin@igv.cnr.it

² INRA, Unité de Recherches Forestières Méditerranéennes, Avignon, France

³ Dipartimento di Biotecnologie Agrarie, Università di Firenze, Firenze, Italy

⁴ INRA, Biodiversité Gènes & Communautés, Cestas, France

⁵ Unidad Mixta de Genómica y Ecofisiología Forestal, INIA-UPM, Madrid, Spain

Genetic variation is generally considered a prerequisite for adaptation to changing environments. Thus, the discovery of genetically depauperate but geographically widespread species is unexpected. In a first work, 12 paternally inherited chloroplast microsatellites were used to estimate population genetic variation across the full range of an emblematic circum-Mediterranean conifer, stone pine (*Pinus pinea* L.). We found that the same chloroplast DNA haplotype was fixed in nearly all of the 34 populations investigated. New data at the nuclear level (14 nuclear microsatellite loci and SNPs scored among about 280 genes) also show a near absence of genetic variation. We screened the literature to identify other cases of genetically depauperate plants. This search indicates that *P. pinea* is truly exceptional among widespread, sexually reproducing plant species for its low level of genetic diversity in both chloroplast and nuclear molecular markers. Stone pine appears to have passed through at least one severe and prolonged demographic bottleneck during the Quaternary, followed by subsequent natural- and human-mediated dispersal across the Mediterranean basin during the Holocene. Measurements from provenance tests indicate that the species harbours a non-negligible amount of variation for adaptive traits. This illustrates that although there is little doubt that genetic variation is the raw material for adaptation, the relationships between neutral diversity, quantitative trait variation, and adaptability are not straightforward.

Keywords: depauperate species, diversity depletion molecular markers, *Pinus pinea*