



HAL
open science

Effect of Defoliation on Carbon Partitioning into Plant Soil Micro-Organisms System of Rye Grass (*Lolium perenne*) Sward

S. Bazot, J. Tavernier, Sylvain Plantureux, Christophe Robin

► **To cite this version:**

S. Bazot, J. Tavernier, Sylvain Plantureux, Christophe Robin. Effect of Defoliation on Carbon Partitioning into Plant Soil Micro-Organisms System of Rye Grass (*Lolium perenne*) Sward. 20. General meeting of the European Grassland Federation, Jun 2004, Luzern, Switzerland. hal-02760109

HAL Id: hal-02760109

<https://hal.inrae.fr/hal-02760109v1>

Submitted on 29 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of Defoliation on Carbon Partitioning into Plant Soil Micro-Organisms System of Rye Grass (*Lolium perenne*) Sward

Bazot S, Tavernier J, Plantureux S., Robin C.

Equipe Rhizosphère, UMR INPL-INRA « Agronomie et Environnement » ENSAIA, 54505 Vandoeuvre-les-Nancy, France.

Stephane.Bazot@ensaia.inpl-nancy.fr

Keywords : Carbon allocation, Defoliation, Rhizosphere, Microbial activity.

Consequences of defoliation on C partitioning and C release by roots are contradictory. (Mikola and Kytöviita 2002, Miller and Roses 1992, Paterson and Sim 1999, 2000, Todorovic *et al* 1999). In this context, this work aimed to quantify the effects of defoliation of ryegrass shoots on C partitioning in the plant-soil-micro-organisms system and on the short-term below ground carbon allocation under two N treatments (0 and 100U). After two days, rhizodeposition is not affected by defoliation. Four days after defoliation, a decrease of shoots C content could be explained by a decrease of nutrient assimilation by roots, resulting from root senescence, and reduced photosynthesis. Senescent tissues involve an increase of C availability for micro-organisms, especially under high N. This explain the enhance of the number of culturable cells (as colony forming units CFU), and the non significant increase of micro-organisms activity (test based on the short term utilization of ¹⁴C glucose by rhizosphere micro-organisms). Thus under high N treatment, defoliation involve, after 4 days, an increase of rhizodeposition mainly in the form of sloughed-off cells. Finally, the allocation of current assimilates to plant roots is not affected by defoliation, but roots senescence increase the root released organic C.

Session 2