



Dynamics of soil surface crusting: coupled study with image analysis and high resolution Digital Elevation Models

Sophie Leguédais, Isabelle I. Cousin, Frédéric Darboux, Olivier Fischer, Yves Le Bissonnais

► To cite this version:

Sophie Leguédais, Isabelle I. Cousin, Frédéric Darboux, Olivier Fischer, Yves Le Bissonnais. Dynamics of soil surface crusting: coupled study with image analysis and high resolution Digital Elevation Models. Soil erosion patterns: Evolution, spatio-temporal dynamics and connectivity, Oct 2002, Müncheberg, Germany. 1 p. hal-02761261

HAL Id: hal-02761261

<https://hal.inrae.fr/hal-02761261>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



COST 623 'Soil Erosion and Global Change' WG 1: Scale Issues, WG 2: Thresholds
ESSC (European Society for Soil Conservation)
DBG (German Society for Soil Science)
ZALF (Centre for Agricultural Landscape and Land Use Research, Müncheberg, Germany)

International Symposium

Soil Erosion Patterns: Evolution, Spatio-Temporal Dynamics and Connectivity

10th – 12th October 2002

at the

**Centre for Agricultural Landscape and Land Use Research (ZALF),
Müncheberg, Germany**

Book of Abstracts

www.zalf.de/essc/symp-pattern.html

Dynamics of soil surface crusting: coupled study with image analysis and high resolution Digital Elevation Models

Sophie Leguédais, Isabelle Cousin, Frédéric Darboux, Olivier Fischer, Yves Le Bissonnais

sophie.leguedois@orleans.inra.fr
INRA Orléans Soil Science, France

Key words: image analysis, roughness, soil surface crusting

Developments of soil erosion modelling need to take spatio-temporal dynamics of soil surface characteristics into account especially soil surface structure. Soil surface structure is a key factor influencing interrill soil erosion behaviour through soil hydrodynamic properties, runoff rate and sediment detachment. For cultivated soils and between field works, surface structure is driven essentially by evolution of soil crusting. As a consequence, soil surface structure is modified during rainfall event and varies with the topographic position. Soil surface structure is also soil dependent.

The aims of this study were:

- to develop a tool relevant to describe and to quantify spatio-temporal evolution of soil crusting, and that could be used for soil erosion modelling needs,
- to confront information about spatio-temporal evolution of soil crusting and dynamics of interrill soil erosion given by runoff rate, splash rate and sediment delivery, for a range of soil types.

Soil surface structure evolution was made by laboratory rainfall simulations. Rainfall simulations were carried out on air-dried soils samples packed in small trays of $0.5 \times 0.5 \text{ m}^2$ with a 5 % slope. Soil surface was initially moulded with an oriented roughness of about 2-cm high, parallel to the slope. Three cultivated soils with various susceptibilities to crusting were studied. Rainfall was applied with an intensity of about 30 mm.h^{-1} during 2 hours after the runoff initiation. Three replicated rainfall simulations were done for each soil type.

Soil erosion characteristics were measured by sampling runoff every 10 min and splashed particles every 20 min. Splash was collected with a box set at the upstream side of the tray.

Digital colour pictures of the soil surface were taken every 10 minutes during the whole rainfall event. Photographs were studied by image analysis in order to obtain spatial information about soil surface sealing and crust type.

Digital Elevation Models (DEMs) of the soil surface were obtained with a high-resolution profile laser scanner. Three acquisitions were done at different times:

- before the beginning of the rainfall simulation,
- when runoff coefficient reaches 10 %,
- after the rainfall event.

For each micro-topographic acquisition, a photograph of the soil surface was taken. DEMs were processed in order to quantify spatial evolution of soil surface roughness.

Finally, results obtained from image analysis and study of DEMs were coupled to quantitatively describe spatio-temporal evolution of crusting and surface roughness in order to follow dynamics of soil surface structure.