



HAL
open science

Effect of live yeast culture supplementation on rumen fermentation in lactating dairy goats

Sylvie Giger-Reverdin, Daniel Sauvant, Joseph-Alexandre J.-A. Tessier, Gaëlle Bertin, Pierre Morand-Fehr

► To cite this version:

Sylvie Giger-Reverdin, Daniel Sauvant, Joseph-Alexandre J.-A. Tessier, Gaëlle Bertin, Pierre Morand-Fehr. Effect of live yeast culture supplementation on rumen fermentation in lactating dairy goats. 8. International Conference on Goats, Jul 2004, Pretoria, South Africa. hal-02761790

HAL Id: hal-02761790

<https://hal.inrae.fr/hal-02761790>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of live yeast culture supplementation on rumen fermentation in lactating dairy goats

S. Giger-Reverdin^{1#}, D. Sauvant¹, J. Tessier¹, G. Bertin² and P. Morand-Fehr¹

¹UMR INRA INA P-G, Physiologie de la Nutrition et Alimentation, 16 rue Claude Bernard, 75005 Paris, France

²ALLTECH France, 2-4 avenue du 6 juin 1944, 95190 Goussainville, France

Abstract

Addition of yeast to dairy cow diets might be beneficial for milk production. However, data concerning goats are scarce, especially on the role of yeast on rumen metabolism. Thus, four goats, according to a cross-over design, received successively two diets with or without living yeasts. Animals were in mid-lactation and received a total mixed diet *ad libitum*. For the yeast diet (Y), each goat received 5 g of *Saccharomyces cerevisiae* CBS 493.94 twice a day. Samples of ruminal content were taken every two hours for eight hours after the morning feeding. Yeast addition did not have any statistical effect either on feeding pattern or on concentrations of volatile fatty acids, ammonia, lactate or soluble carbohydrate in ruminal fluid. The pH was numerically higher for the yeast diet compared to the control. Ruminal buffering (BC) capacity of the Y diet was significantly higher than that of the control. The BC increased as pH decreased. Dietary effects and ruminal soluble carbohydrate concentrations explained part of the residual of the equation linking buffering capacity and pH. Yeast addition also avoided some lactate peaks in the first hours of fermentation. This experiment clearly pointed out that the *Saccharomyces cerevisiae* CBS 493.94 yeast has an effect on ruminal metabolism when considering its BC and might have a smoothing effect on the appearance of lactate peaks or on a decreased pH. These effects are of particular interest for those diets which might induce acidosis such as some diets rich in rapidly fermentable energy given to high producing dairy goats in early lactation.

Keywords: *Saccharomyces Cerevisiae* yeast, rumen digestion, buffering capacity, dairy goats

[#]Corresponding author. E-mail: giger@inapg.inra.fr

Introduction

Several experiments using dairy cows or goats have shown that yeast addition to the diet might increase milk production (Gunther, 1990; Piva *et al.*, 1993; Adams *et al.*, 1995; Putnam *et al.*, 1997; Nocek *et al.*, 2003) while other studies did not find any difference (Arambel & Kent, 1990; Chiquette, 1995; Kamalamma *et al.*, 1996; Salama *et al.*, 2002). Studies where yeast has been supplemented to dairy goat diets are limited, in particular, studies where the ruminal mode of action of yeast has been studied (Flachowsky *et al.*, 1993; Hadjipanayiotou *et al.*, 1997; Salama *et al.*, 2002). Furthermore, parameters such as buffering capacity (BC) of ruminal content and lactate concentrations have been either seldom or never measured. These parameters are of particular interest when animals receive diets that might induce acidosis such as diets rich in rapidly fermentable carbohydrates given to high producing dairy goats in early lactation.

Materials and Methods

Four rumen fistulated goats received successively either the control diet (C) or the Yeast diet (Y) in a cross-over experimental design. For the Yeast diet, each goat received 5 g of *Saccharomyces cerevisiae* CBS 493.94 [Alltech Company, Nicholasville, Kentucky, USA] twice a day with a dosing gun. Animals were in mid-lactation and milked twice a day. They received *ad libitum* a total mixed diet containing on a dry matter (DM) basis: 40% maize silage, 10% dehydrated lucerne and 50% of a concentrate (10% wheat bran, 18% maize gluten meal, 10% lupine seeds, 28% sugar beet pulp, 29% citrus pulp, 2% molasses and 3% mineral and vitamins mixture). Water was available *ad libitum*.

After an adaptation of three weeks to the diet, samples of ruminal content were taken from the dorsal part of the rumen of each goat every two hours for eight hours after the morning feeding. About 100 mL were immediately filtered through cheesecloth, pH was recorded and thereafter the BC by means of titration

of 20 mL of rumen juice with 1N acetic acid (Giger-Reverdin *et al.*, 2000) which was performed until the pH decreased from its initial value to 4. The evolution in pH was expressed as an exponential function of the quantity of acetic acid added (in meq H⁺): $pH = a - b \text{ eqH} + c \exp^{-d \text{ eqH}}$

Buffering capacity, or the inverse of the derivative function at the initial pH, is equal to $1/(b + (c * d))$. Samples were kept at -20 °C until analysed for volatile fatty acids, soluble carbohydrates, ammonia and lactate concentrations. Statistical analyses were performed according to the “proc mixed” procedure of SAS[®] in order to take into account the repeated effect on a same day for a given goat. The diet and sampling time effects and their interactions were tested for each variate.

Results and Discussion

Yeast addition did not affect DM intake (DMI) (g DMI per kg body-weight (DMI BW⁻¹)) at any sampling time or any of the rumen parameters measured, except for buffering capacity.

Table 1 Effect of yeast culture supplementation on dry matter intake (DMI) and rumen fermentation parameters

	Control diet	Yeast diet	Diet effect	Time effect
DMI/body weight (g/kg)	32.8	31.3	NS	P < 0.0001
pH	6.09	6.15	NS	P < 0.03
Buffering capacity	0.0299 ^a	0.0448 ^b	P < 0.001	NS
Volatile fatty acids (mmole/L)	152	153	NS	P < 0.01
Ammonia (g/L)	0.0795	0.0867	NS	P < 0.001
Lactate (mmole/L)	0.455	0.207	NS	NS
Soluble carbohydrates(g/L)	4.27	4.67	NS	P < 0.01

^{a, b}Row means with common superscripts do not differ (P > 0.05)

Compared to the control diet, pH was numerically higher with the yeast addition. This tendency is in agreement with other researchers who did not find any impact of yeast on ruminal metabolism (Erasmus *et al.*, 1992; Doreau & Jouany, 1998), while others recorded that yeast increased pH (Kumar *et al.*, 1997; Roa *et al.*, 1997). There was no lactate peak with the yeast supplemented diet that explained the lower lactate value of goats receiving the Y diet compared to goats receiving the C diet. Yeast might have had a smoothing effect on lactate peaks appearance, which is in agreement with other results (Williams *et al.*, 1991). The small increase in soluble carbohydrate concentration suggests that yeast might act on carbohydrate metabolism in the rumen. Furthermore, BC was higher with the yeast diet compared to the control diet (Figure 1).




Figure 1 Dietary effect on buffering capacity and pH

Buffering capacity increased as pH decreased:

$$BC = 0.187 - 0.0244 \text{ pH} \quad (r^2 = 0.437, n = 32, \text{RSD} = 0.008484)$$

Dietary effect explained part of the residual of this equation (P < 0.001), such as the ruminal soluble carbohydrates concentration (P < 0.01).

Conclusion

The *Saccharomyces cerevisiae* CBS 493.94 yeast improved BC of the rumen significantly and tended to indicate level of significance to an increase in pH and decrease lactate concentrations. This effect might be more significant for diets higher in NSC than the one tested. These effects are of particular significance for diets which might induce acidosis such as diets rich in rapidly fermentable carbohydrates given to high producing dairy goats in early lactation.

References

- Adams, A.L., Harris, B., Jr., Horn, H.H.v. & Wilcox, C.J., 1995. Effects of varying forage types on milk production responses to whole cottonseed, tallow, and yeast. *J. Dairy Sci.* 78, 573-581.
- Arambel, M.J. & Kent, B.A., 1990. Effect of yeast culture on nutrient digestibility and milk yield response in early- to midlactation dairy cows. *J. Dairy Sci.* 73, 1560-1563.
- Chiquette, J., 1995. *Saccharomyces cerevisiae* and *Aspergillus oryzae*, used alone or in combination, as a feed supplement for beef and dairy cattle. *Can. J. Anim. Sci.* 75, 405-415.
- Doreau, M. & Jouany, J.P., 1998. Effect of a *Saccharomyces cerevisiae* culture on nutrient digestion in lactating dairy cows. *J. Dairy Sci.* 81, 3214-3221.
- Erasmus, L.J., Botha, P.M. & Kistner, A., 1992. Effect of yeast culture supplement on production, rumen fermentation, and duodenal nitrogen flow in dairy cows. *J. Dairy Sci.* 75, 3056-3065.
- Flachowsky, G., Tiroke, K. & Matthey, M., 1993. Influence of yeast (*Saccharomyces cerevisiae* as Yea-Sacc or Levaferm) on in sacco dry matter degradability and ruminal parameters of variously fed small ruminants. *Archives Anim. Nutr.* 42, 159-169.
- Giger-Reverdin, S., Maaroufi, C., Jarnot, L., Müller, R., Bontems, V. & Sauvant, D., 2000. The influence of pea grinding and dietary nitrogen content on buffering capacity of diets offered to lactating goats. (Influence de la granulométrie du pois et du niveau azoté du régime sur son pouvoir tampon pour des chèvres en lactation). In: 7th Int. Conf. on Goats, Tours (France). 2000/05/15-18, Eds. Institut de l'Elevage and INRA. pp. 86-88.
- Gunther, K.D., 1990. Lebende Hefekulturen als Zusatzstoff in der Milchviehfütterung. (Live yeast cultures as additives in dairy cattle feeds). *Kraftfutter*, 168-172.
- Hadjipanayiotou, M., Antoniou, I. & Photiou, A., 1997. Effects of the inclusion of yeast culture on the performance of dairy ewes and goats and the degradation of feedstuffs. *Livest. Prod. Sci.* 48, 129-134.
- Kamalamma, U., Krishnamoorthy, U. & Krishnappa, P., 1996. Effect of feeding yeast culture (Yea-sacc1026) on rumen fermentation *in vitro* and production performance in crossbred dairy cows. *Anim. Feed Sci. Technol.* 57, 247-256.
- Kumar, U., Sareen, V.K. & Singh, S., 1997. Effect of yeast culture supplement on ruminal microbial populations and metabolism in buffalo calves fed a high roughage diet. *J. Sci. Food Agric.* 73, 231-236.
- Nocek, J.E., Kautz, W.P., Leedle, J.A.Z. & Block, E., 2003. Direct-fed microbial supplementation on the performance of dairy cattle during the transition period. *J. Dairy Sci.* 86, 331-335.
- Piva, G., Belladonna, S., Fusconi, G. & Sicbaldi, F., 1993. Effects of yeast on dairy cow performance, ruminal fermentation, blood components, and milk manufacturing properties. *J. Dairy Sci.* 76, 2717.
- Putnam, D.E., Schwab, C.G., Socha, M.T., Whitehouse, N.L., Kierstead, N.A. & Garthwaite, B.D., 1997. Effect of yeast culture in the diets of early lactation dairy cows on ruminal fermentation and passage of nitrogen fractions and amino acids to the small intestine. *J. Dairy Sci.* 80, 374-384.
- Roa, V.M.L., Barcena-Gama, J.R., Gonzalez, M., S., Mendoza, M., G., Ortega, C.M.E. & Garcia, B.C., 1997. Effect of fiber source and a yeast culture (*Saccharomyces cerevisiae*1026) on digestion and the environment in the rumen of cattle. *Anim. Feed Sci. Technol.* 64, 327-336.
- Salama, A.A.K., Caja, G., Garin, D., Albanell, E., Such, X. & Casals, R., 2002. Effects of adding a mixture of malate and yeast culture (*Saccharomyces cerevisiae*) on milk production of Murciano-Granadina dairy goats. *Anim. Res.* 51, 295-303.
- Williams, P.E., Tait, C.A., Innes, G.M. & Newbold, C.J., 1991. Effects of the inclusion of yeast culture (*Saccharomyces cerevisiae* plus growth medium) in the diet of dairy cows on milk yield and forage degradation and fermentation patterns in the rumen of steers. *J. Anim. Sci.* 69, 3016-3026.