

## Potentialities of the multiobjective approach to assimilate remote sensing data in SVAT models

Jérôme Demarty, Albert Olioso, Catherine Ottle, Olivier Marloie, Jean-Pierre Wigneron, Mickaël Pardé, Isabelle Braud, Joël Noilhan, Jean-Christophe Calvet

## ▶ To cite this version:

Jérôme Demarty, Albert Olioso, Catherine Ottle, Olivier Marloie, Jean-Pierre Wigneron, et al.. Potentialities of the multiobjective approach to assimilate remote sensing data in SVAT models. EGS-AGU-EUG Joint Assembly, Apr 2003, Nice, France. hal-02762470

HAL Id: hal-02762470 https://hal.inrae.fr/hal-02762470

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## POTENTIALITIES OF THE MULTIOBJECTIVE APPROACH TO ASSIMILATE REMOTE SENSING OBSERVATIONS IN SVAT MODELS

- J. Demarty (1), A. Olioso (1), C. Ottlé (2), O. Marloie (1), J. P. Wigneron (3), M. Pardé (3), I. Braud (4), J. Noilhan (5) and J. C. Calvet (5)
- (1) INRA Avignon, France, (2) CETP Vélizy, France, (3) INRA Bordeaux France, (4) CEMAGREF Lyon, France, (5) CNRM, Toulouse, France (jdemarty@avignon.inra.fr/ Fax: +33 4 32 72 23 62)

The purpose of this paper concerns remote sensing data assimilation in physically based model by the way of a multiobjective calibration procedure. Generally, the calibration of Soil Vegetation Atmosphere Transfer (SVAT) models consists to determine the best parameter set in terms of model performances. Due to model structural uncertainty and to errors on the observations, a such determination of a unique parameter set is generally impossible. This problem of non-uniqueness of the solution can be investigated by a multiobjective approach, in which different objective functions are optimized simultaneously.

In this context we developed a multiobjective approach allowing to combine sensitivity analyses and an iterative optimisation procedure of the model parameters in terms of uncertainty ranges. This paper investigated the potentialities of a such calibration approach in a remote sensing context, in order to drive SVAT models only by the way of remote sensing observations. The consequences of the calibration procedure on the simulated surface processes were analysed for different studied cases. Main results showed that physically based SVAT models can be constrained by remote sensing observations without initial a priori information about soil and vegetation properties.

Potentialities of the multiobjective approach for remote Sensing data assimilation in SVAT models.