



HAL
open science

SSR-typing of *Botrytis cinerea* reveals rapid evolution of population structure following introduction of marked isolates in tomato greenhouses

Véronique Decognet, Philippe C. Nicot, Marc Bardin, Yannie Trottin-Caudal

► To cite this version:

Véronique Decognet, Philippe C. Nicot, Marc Bardin, Yannie Trottin-Caudal. SSR-typing of *Botrytis cinerea* reveals rapid evolution of population structure following introduction of marked isolates in tomato greenhouses. 1. Workshop, Oct 2005, Kaiserslautern, Germany. hal-02763173

HAL Id: hal-02763173

<https://hal.inrae.fr/hal-02763173v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

Decognet, V., Nicot, P. C., Bardin, M., Trottin-Caudal, Y. (2005). SSR-typing of *Botrytis cinerea* reveals rapid evolution of population structure following introduction of marked isolates in tomato greenhouses. Presented at 1. *Botrytis cinerea* genome Workshop, Kaiserslautern, Germany (2005/10/06-08).

Decognet¹, V.; **Nicot¹, P.C.**; Bardin¹, M.; Trottin-Caudal², Y.

¹INRA, Plant Pathology Unit, BP94, Domaine St Maurice, F-84140 Montfavet, France

²CTIFL, Centre de Balandran, BP32, F-30127 Bellegarde, France

Botrytis cinerea can rapidly produce massive amounts of inoculum on diseased plants and control methods have been developed to reduce spore production. In addition to spores produced within a crop (endogenous inoculum), the airborne spora of *Botrytis* over a canopy may also include exogenous inoculum carried from a variety of places and hosts. The relative proportion of both types of inoculum may influence the efficacy of control methods and the impact of selective pressures resulting from their implementation.

A trial was set up to quantify the impact of endogenous inoculum on disease development and on the population structure of *B. cinerea* in four tomato glasshouse compartments of CTIFL-Balandran. Plants were grown in quasi-commercial conditions. Isolates were collected from the air spora 4 days before and 14 days after inoculation of 6 plants per compartment with one of two reference strains (differing in their SSR profile and aggressiveness on tomato). Disease development was monitored and isolates were collected on stem lesions 60 days after inoculation. All isolates were characterised using 9 microsatellites described by E. Fournier, INRA-Versailles.

Among 80 isolates collected in the air spora prior to inoculation (while all plants were healthy), none had SSR profiles similar to either of the reference strains. Following inoculation with the reference strains, lesions developed and sporulation was observed on all inoculation points. Within 14 days, SSR profiles identical to those of either introduced strain represented 63% of the 323 samples characterised from the air spora of the four compartments.

The disease spread steadily to non-inoculated plants and incidence reached an average of 3-7 lesions per plant (depending on the compartment) by 90 days after inoculation. Among 224 isolates collected from stem lesions at 60 days after inoculation, 61% and 39% had SSR profiles similar to the aggressive and to the less aggressive reference strains, respectively.

This suggests that the displacement of the initially dominant population of *Botrytis* was concomitant with its negligible contribution to the epidemic on tomatoes. These results are compatible with the hypothesis of a polycyclic development of *Botrytis* epidemics in tomato greenhouses and illustrate the importance of endogenous inoculum in this growing system. Furthermore, they raise the question of possible host specificity within *Botrytis cinerea* in naturally occurring epidemics.

Comment citer ce document :

Decognet, V., Nicot, P. C., Bardin, M., Trottin-Caudal, Y. (2005). SSR-typing of *Botrytis cinerea* reveals rapid evolution of population structure following introduction of marked isolates in tomato greenhouses. In: *Botrytis cinerea* genome (p. 1 p.). Presented at 1. Workshop, Kaiserslautern, DEU (2005-10-06 - 2005-10-08).