



HAL
open science

Satellite measurements and vegetation models for global carbon cycle studies

G rard Dedieu, Anne Ruimy, Philippe Maisongrande, Laurent Kergoat,
Sophie Moulin, Bernard Saugier

► **To cite this version:**

G rard Dedieu, Anne Ruimy, Philippe Maisongrande, Laurent Kergoat, Sophie Moulin, et al.. Satellite measurements and vegetation models for global carbon cycle studies. 7. International Symposium Physical Measurements and Signatures in Remote Sensing, Apr 1997, Courchevel, France. hal-02770389

HAL Id: hal-02770389

<https://hal.inrae.fr/hal-02770389>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.

SATELLITE MEASUREMENTS AND VEGETATION MODELS FOR GLOBAL CARBON CYCLE STUDIES

G. Dedieu¹, A. Ruimy², P. Maisongrande¹, L. Kergoat¹, S. Moulin¹ and B. Saugier²

1Centre d'Etudes Spatiales de la Biosphère (CESBIO)
18 avenue Edouard Belin, 31401 Toulouse Cedex 4, France
2Laboratoire d'Ecophysiologie Végétal (LEV), U.R.A. 1492,
Université de Paris Sud, Centre d'Orsay, Bât 362, 91405 Orsay Cedex, France

Global Terrestrial Biosphere Models (TBMs) are being developed by the scientific community to understand and forecast interactions (impacts and feedbacks) between terrestrial biosphere, soils, and climate. One specific objective is to predict the evolution of land carbon sources and sink and atmospheric CO₂ concentrations under various scenarios.

TBMs have proved very useful tools for understanding the role of the terrestrial biosphere in the current carbon cycle. However intercomparisons performed under the auspices of IGBP have shown that discrepancies exist between model results such as Net Primary Productivity (NPP) and above all intermediate prognostic variables (e.g. light use efficiency, autotrophic respiration). Further improvements and validations are needed, especially for predicting biospheric carbon fluxes and budget under future environmental conditions. TBM improvement has to cope with several difficulties such as accounting for main vegetation processes, specification of space and time dependent parameters, and validation of the models.

We have developed both diagnostic and prognostic models for estimating global NPP (Ruimy et al., 1994, Ruimy et al., 1996, Kergoat, 1996). The diagnostic model is driven by satellite data: NPP is computed as the product of photosynthetically active radiation (PAR) that plant canopies absorb by radiation use efficiency. Coefficients of PAR absorption are derived weekly from NOAA/AVHRR visible and near infrared measurements. This diagnostic model, strongly constrained by satellite observations, is a useful tool for understanding the current carbon cycle and its fluctuations. Prognostic models are also needed to simulate future feedbacks between climate and the terrestrial biosphere. A satisfactory representation of plant production processes is necessary to get a realistic simulation of the evolution of global plant productivity with increased CO₂ and associated climatic change.

The prognostic model we have developed incorporates main physiological processes such as photosynthesis, evapotranspiration, and assimilate allocation. In its current version, it predicts canopy development (LAI) by assuming that vegetation makes an optimal use of available resources (water, light, ...). This model can be tested by comparing predicted variables (LAI, ETR, soil moisture, ...) to independent estimates (ground measurements or satellite retrievals).

We will first present briefly the principles of these two models, and some of our achievements, such as global mapping of NPP, LAI and evapotranspiration, and interannual fluctuations of NPP, including the consequences of El-Niño events. Then, the focus will be on the use of satellite measurements (NOAA/AVHRR, SPOT-HRV) for providing model parameters or for validating the results. Satellite based retrieval of model parameters includes "forcing" strategies (e.g. fPAR) and assimilation techniques. Validation of TBMs is a crucial issue, and we will discuss the comparison of model results with in-situ flux measurements, atmospheric CO₂ concentration and satellite data. We will show that satellite measurements and observations of interannual variabilities could play an important role to make models more reliable, particularly if better quality data sets become available.