



**HAL**  
open science

## Ontogeny of the GH/IGF system mRNA during embryonic development in rainbow trout (*Oncorhynchus mykiss*)

Jean-Charles Gabillard, Pierre-Yves Rescan, Daniel Boujard, Chantal Cauty, Claudine Weil, Pierre-Yves Le Bail

### ► To cite this version:

Jean-Charles Gabillard, Pierre-Yves Rescan, Daniel Boujard, Chantal Cauty, Claudine Weil, et al.. Ontogeny of the GH/IGF system mRNA during embryonic development in rainbow trout (*Oncorhynchus mykiss*). 20. Conference of European comparative endocrinologists, Universidade do Algarve. PRT., Sep 2000, Faro, Portugal. hal-02770523

**HAL Id: hal-02770523**

**<https://hal.inrae.fr/hal-02770523>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**EUROPEAN SOCIETY FOR COMPARATIVE  
ENDOCRINOLOGY**

**20<sup>TH</sup> CONFERENCE OF EUROPEAN  
COMPARATIVE ENDOCRINOLOGISTS**

5-9<sup>th</sup> September 2000  
Faro, Portugal

**Programme and Abstracts**

Centre of Marine Sciences  
University of Algarve

5-9th September 2000  
Faro, Portugal  
Centre of Marine Sciences  
University of Algarve

**ONTOGENY OF THE GH/IGF SYSTEM mRNA DURING  
EMBRYONIC DEVELOPMENT IN RAINBOW TROUT**

*(Oncorhynchus mykiss)*

Gabillard, J.-C.<sup>1</sup>, Rescan, P.-Y.<sup>1</sup>, Boujard, D.<sup>2</sup>, Cauty, C.<sup>1</sup>, Weil, C.<sup>1</sup>  
and Le Bail, P.-Y.<sup>1</sup>

<sup>1</sup> INRA/SCRIBE, <sup>2</sup> UPRESA/Université de Rennes I, Campus de  
Beaulieu 35042 Rennes cedex, France

In fish, the GH/IGF system has been shown to be important for growth. In rainbow trout, we have studied its development during the first embryonic stages, as defined by Ballard, 1973 (J. Exp. Zool., 184,7-26). The presence of the different molecules (IGF-1, IGF-2, IGF-R1a, IGF-R1b and GH) of this system was examined by using whole *in situ* hybridization. Concerning GH, a strong signal was observed from stage 17 (around 20 pair of postanal somites). At this stage around 5 cells were visualized and the number increased until hatching, stage 23 (around 50 cells). On the other hand, at stage 17 the GH protein was not detected by using either an homologous radioimmunoassay or immunocytochemistry, while it was possible at stage 23. This result seems to show that there is probably a delay between the transcription and the translation of the GH genes during rainbow trout ontogenesis. Concerning the other molecules (IGF-1, IGF-2, IGF-R1a, IGF-R1b), the whole *in situ* hybridization technique did not allow to evidence them without ambiguity while their presence was detected during rainbow trout embryogenesis by other authors, using RT-PCR and RPA techniques. These observations suggest that the expression of these different molecules is low and probably ubiquitous at this period of development.