



HAL
open science

Below-ground nitrogen transfer from oak seedlings facilitates *Molinia* growth: 15 N pulse-chase labelling

Marine Fernandez, Philippe Malagoli, Antoine Vernay, Thierry Ameglio,
Philippe Balandier

► **To cite this version:**

Marine Fernandez, Philippe Malagoli, Antoine Vernay, Thierry Ameglio, Philippe Balandier. Below-ground nitrogen transfer from oak seedlings facilitates *Molinia* growth: 15 N pulse-chase labelling. *Plant and Soil*, 2020, 449 (1-2), pp.343-356. 10.1007/s11104-020-04473-9 . hal-02775498

HAL Id: hal-02775498

<https://hal.inrae.fr/hal-02775498v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Below-ground nitrogen transfer from oak seedlings facilitates *Molinia* growth: ^{15}N pulse-chase labelling

Marine Fernandez · Philippe Malagoli · Antoine Vernay ·
Thierry Améglio · Philippe Balandier

Received: 20 June 2019 / Accepted: 24 February 2020 / Published online: 20 March 2020
© Springer Nature Switzerland AG 2020

Abstract

Aims Belowground carbon transfer from plant to plant has been extensively described, but such transfer for nitrogen has been less thoroughly investigated when the donor is a non- N_2 -fixing species. This study, applied to forest regeneration, aimed to determine whether tree seedlings facilitated neighbouring grass growth through

nitrogen transfer at an early stage of development, thus facilitating nitrogen acquisition by understory species. **Methods** *Quercus petraea* seedlings were planted in pots either sole-grown or mixed-grown with *Molinia caerulea* tufts or another oak seedling. ^{15}N -urea pulse-chase labelling (cotton wick method) was performed in oak shoots and the fate of ^{15}N in each soil and plant compartment was tracked for one year. N transfer pathways were investigated using two degrees of physical separation between root systems.

Results *Molinia* dry weight was higher when mixed-grown with oak seedlings than when sole-grown. Increase in grass dry weight correlated with N transfer from donor oak to receiver *Molinia*. Interestingly, the presence of *Molinia* increased N rhizodeposition of oak. N allocation in donor oak towards root in winter and shoot in spring was enhanced.

Conclusions Oak seedlings facilitated *Molinia* growth through rapid N transfer, underlining the ability of non- N_2 -fixing species to supply N to neighbours. ^{15}N allocation within donor oak and its rhizodeposition depended on neighbour identity.

Responsible Editor: Xinhua He.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11104-020-04473-9>) contains supplementary material, which is available to authorized users.

M. Fernandez · P. Malagoli (✉) · T. Améglio
Université Clermont Auvergne, INRAE, PIAF,
F-63000 Clermont-Ferrand, France
e-mail: philippe.malagoli@uca.fr

M. Fernandez
e-mail: marine.fernandez@uca.fr

T. Améglio
e-mail: thierry.ameglio@inrae.fr

A. Vernay
Department of Forest Ecology and Management, Swedish
University of Agricultural Sciences (SLU), SE-901 83 Umeå,
Sweden
e-mail: antoine.vernay@slu.se

P. Balandier
INRAE, UR EFNO, Domaine des Barres,
F-45290 Nogent-sur-Vernisson, France
e-mail: philippe.balandier@inrae.fr

Keywords Facilitation · Nitrogen · Belowground transfer · Rhizodeposition · *Quercus petraea* · *Molinia caerulea*

Abbreviations

SG	Sole-grown
Mesh1	1 μm Mesh
Mesh30	30 μm Mesh
MG	Mixed-grown

Introduction

Clements et al. (1929) describing plant interactions defined resource competition as a “physical process” and “a combined need in excess of the supply”. This implies that plants compete for light, water and nutrients only when they occupy the same zone of air or soil. Competition then occurs when one individual successfully reaches and captures limiting resources to the detriment of another individual. In this way it lowers the performance of its neighbours in biomass production, growth or reproduction. For many years, work on interactions has mainly focused on strategies of competition for resources (Bleasdale 1960; Grime 1974; Tilman 1990; Craine 2005; Craine and Dybzinski 2013), seen as a passive mechanism by which plants draw resources from a common reservoir (*i.e.* soil) with ranging efficiency.

Schreiner and Reed (1907) in an early literature review on the capacity of living roots to excrete matter in the soil pointed out that toxic excretions from roots exhibited deleterious effects on the growth and development of other plants. Allelopathy, or more generally chemical interference, is the process by which plants release allelochemicals into the environment that have significant direct or indirect effects on other plant processes (Molisch 1937; Muller 1969; Rice 1985; Grove et al. 2013). Resource competition and chemical interference can thus occur simultaneously (Mallik 1998; San Emeterio et al. 2007) and are species-specific (Fernandez et al. 2016). For example, the presence of beech (*Fagus sylvatica*) reduced the ability of maple (*Acer pseudoplatanus*) to take up inorganic nitrogen, so slowing its growth (Li et al. 2015), although the precise mechanism was not described.

Chemical compounds can also exert a beneficial influence (Rice 1985; Callaway 1995; Kohli et al. 1998). The molecular entities exuded by living roots include metabolites that include carbon, phosphorus and nitrogen compounds (Virtanen and Laine 1939; Slankis et al. 1964; Rovira 1969). Recent studies have emphasized that nutrient and information exchanges among plants are more generalized than had been thought (Pierik et al. 2013), and play a key role in plant interactions. The process by which living and dead roots exude substances in soil, shed fine roots and sloughed-off cells and tissue, along with cell lysates and decomposed root materials is termed rhizodeposition (Philipps et al. 2006; Scandellari et al. 2010; Mommer

et al. 2016). Legumes offer a well-known case of positive substance release in the soil. Indeed, they can release about 50% of their total N into the soil by rhizodeposition (Boulter et al. 1966; Shamoot et al. 1968; Friedrich and Dawson 1984; Khan et al. 2002; Mahieu et al. 2009; Fustec et al. 2010). Some authors have described the ability of neighbouring grass to take up nitrogen released by legumes (Henzell et al. 1964; Vallis et al. 1967; Whitney et al. 1967). N transfer from legumes to non-leguminous species has been reported in intercrops (Stagnari et al. 2017) such as soybean-weed (Moyer-Henry et al. 2006), pea-corn (van Kessel et al. 1985; Bethlenfalvay et al. 1991), pea-barley (Jensen 1996; Johansen and Jensen 1996) and clover-ryegrass (Haystead et al. 1988). Transfer of nutrients has mainly been studied for N₂-fixing plants, and only very few examples (“less than a handful” – Teste et al. 2015) have shown exchange of nitrogen between non-fixing plants. These N transfers can be as high as 4% from donor plant N, which is not marginal for a non-N₂-fixing species (Teste et al. 2015) especially in infertile soils (Eissenstat 1990). Haystead and Marriott (1978, 1979) concluded that transfer was a complex multicomponent process that could involve key actors such as soil microorganisms. N transfer pathways can include root intermingling, potential common mycorrhizal networks or rhizodeposition of diverse molecular entities (Ek et al. 1996; Teste et al. 2015; He et al. 2019). All these interactions may result in facilitation of the targeted species. Studies on forest trees found that birch (*Betula papyrifera*) and Douglas fir (*Pseudotsuga menziesii*) were colonized by the same ectomycorrhizal fungi (Simard and Perry 1997), allowing bidirectional carbon transfer (Simard et al. 1997a, b).

Facilitation is defined as a positive effect of plants on the establishment or growth of other plant species, and that causes harm to neither (Hunter and Aarssen 1988; Bertness and Callaway 1994; Callaway 1995; Bruno et al. 2003). As an example for nitrogen, in a silvopastoral systems in Patagonia, Gargaglione et al. (2014) showed that *Nothofagus* facilitated grass N uptake, probably owing to improved microclimatic conditions or reduced competition for N between soil microorganisms and grasses. However, in some cases, both competition and facilitation can occur between plants. Antagonistic facilitation occurs when a species A has a positive effect on a species B, while B has a negative effect on the species A (Bronstein 2009; Schöb et al. 2014). In the context of forest regeneration, Vernay et al.

(2018) described such a negative effect of the grass *Deschampsia cespitosa* on *Quercus petraea* seedling growth, while *Q. petraea* improved *D. cespitosa* biomass production. The strong ability of grasses to take up nutrients, and particularly nitrogen, has been demonstrated in many cases (e.g. Coll et al. 2004; Balandier et al. 2006), but the mechanisms underlying the facilitative effect of young oak seedlings on the grass are not known. From the literature briefly reviewed above, we hypothesized that nitrogen transfer between interacting plants occurs, even in the absence of N₂ fixation, through facilitation of *Molinia* growth by oak seedlings. We specifically tested this hypothesis on *Q. petraea* seedlings in competition with a common perennial forest grass in temperate forests, *Molinia caerulea*. Plants were grown in pots under controlled conditions. ¹⁵N pulse-chase labelling (He et al. 2009) was applied to the oak, and ¹⁵N retrieval in soil and *Molinia* was quantified after 3, 5, 8 and 11 months to characterize N fate seasonally. N transfer pathways were also indirectly investigated by inserting a mesh between the two species. Two mesh sizes were used: 1 µm to allow only chemical transfer (Mesh1), and 30 µm to prevent root contact but allow both hyphae contact and chemical transfer (Mesh30).

Materials and methods

Experimental design

The experiment was conducted in pots under outdoor conditions in Clermont-Ferrand (Auvergne, France, 45°45'N 3°07'E, altitude 394 m a.s.l.) from April 2017 to May 2018 ($T_{\text{mean}} = 12.9$ °C, Rainfall = 546 mm). 168 one-year-old bare-root oak seedlings (*Quercus petraea* (Matt.) Liebl.) and 56 grass tufts (*Molinia caerulea*, three tufts on average) were planted in plastic pots, separately or together. Oak seedlings were sourced from a local nursery. They were 21 ± 6.6 g (mean \pm SE) fresh weight, 41.47 ± 5.27 cm in height, and 4.94 ± 0.98 cm in diameter on average. *Molinia caerulea* (three tufts on average) was collected in a local oak forest at Paray-le-Frésil (Auvergne, France; 46°39'N 3°36'E) with 2.04 ± 0.97 g fresh weight per lot. A total of 80 10 L pots and 64 5 L pots were filled with soil (typical luvisol-redoxisol pseudogley, sandy loam) collected in the same forest as the *Molinia*. Forest soil was used to preserve natural colonization by mycorrhizae. Four treatments

based on root system separation or interaction were set up (Fig. 1): (i) two 5 L pots containing either one oak or one *Molinia* tuft were placed side-by-side such that no root interactions were possible (root contact, hyphae contact or chemical transfer) (sole-grown), (ii) oak and *Molinia* tufts were placed in the same 10 L pots but their rooting zones were separated by a nylon (Nitex®) cloth with one of two mesh sizes: either 1 µm to allow only chemical transfer (Mesh1) or 30 µm to prevent root contact but allow both hyphae contact and chemical transfer (Mesh30), and (iii) oak and *Molinia* tufts were placed in the same 10 L pot to allow full belowground interactions through both root and hyphae contacts and chemical transfer (mixed-grown). Effect of species identity in the interaction, i.e. oak or *Molinia* as a neighbour, was tested, replicating every treatment with receiver oak instead of receiver *Molinia* (Fig. 1). Each treatment was replicated four times with a random spatial pot arrangement. To prevent interaction with water availability, the pots were fitted with probes and irrigated to the field capacity throughout the experiment. No fertilizer of any kind was added to the pots during the experiment.

Oak shoot ¹⁵N pulse-chase labelling by the cotton wick technique

To track possible nitrogen transfer from donor oak to receiver *Molinia* or oak, ¹⁵N was supplied to donor oaks using the “cotton wick” method. This method supplies ¹⁵N through stem injection. Oak stems were pierced using a drill (1 mm hole 2 cm above the ground) to push through a cotton wick, both ends of which were dipped into a ¹⁵N solution in a 5 ml Eppendorf tube (¹⁵N-urea, At. ¹⁵N 98%) through two holes in a cap on the top of the tube (Mahieu et al. 2007; Fustec et al. 2010). Drying and loss of solution from the reservoir and the cotton wick were prevented by sheathing the wick with two silicone tubes sealed to the stem and cap (Terostat®, Henkel Surface Technologies, Gulph Mills, USA).

¹⁵N-urea was applied on June 13 and 14, 2017. After oak seedlings had absorbed more than 3 mL of solution (at 0.5%, weight/volume), the reservoir was replenished with a further 3 mL of ¹⁵N urea solution at 0.375% (w/v). When all the solution of ¹⁵N was fully absorbed, the reservoir was washed with 1 mL of deionized water, which was in turn absorbed by the plant. This procedure supplied around 18 mg of ¹⁵N to oak shoots. 5 L pots without cotton wicks were used as controls to determine




Fig. 1 Experimental set-up. Oak was mixed-grown with either another oak plant or a *Molinia* tuft in one pot. When mixed-grown, belowground compartments were either fully intermingled (root contact; mixed-grown) or separated with a mesh of size 30 μM (Mesh30) or 1 μM (Mesh1). Mesh limits (cross in the table) the number of pathways of nutrient exchange between two root

compartments: hyphae and molecules could go through Mesh30 (check mark in the table), but only molecules through Mesh1. Oak was sole-grown beside another oak plant or a *Molinia* tuft with no root contact (sole-grown). Superscript ^{15}N indicates oak was labelled with ^{15}N urea through cotton wick application (see Materials and methods for details)

the natural ^{15}N abundance in each compartment (oak shoots and roots, soil, *Molinia* shoots and roots).

Plant harvesting

Plants were harvested sequentially after total ^{15}N urea uptake was completed (August 2017), during leaf senescence (October 2017), during dormancy (February 2018) and after bud break (May 2018). In winter, only sole-grown and mixed-grown conditions were collected, to estimate ^{15}N allocation, as there is no transfer during dormancy. For this harvest, oak shoots were stem and marcescent leaves, and *Molinia* shoots were senescent leaves.

Plant shoots and roots were collected for both species. For Mesh1 and Mesh30 treatments, soils separated by the mesh were collected individually, and soil in sole-grown and mixed-grown pots was collected as a whole. Soil was then sieved (2 mm) and visible roots were

collected. Roots were washed with 200 mL of a 0.5 mM CaCl_2 solution to remove ^{15}N from their surface and to maintain cell membrane integrity (Epstein 1961). This solution was then sieved and mixed with the soil. About 200 mL of this mixture was used to determine N content and isotope abundance. At each harvest and for each treatment, four replicates of each compartment were collected.

N content and isotopic analysis

Root, shoot and soil samples were dried at 60 $^\circ\text{C}$ for at least 48 h, weighed and ground to a fine powder. Total N content and ^{15}N abundance were then determined with an elemental analyser (vario ISO-TOPE cube, Elementar, Hanau, Germany) in line with a gas isotope ratio mass spectrometer (IsoPrime 100, Isoprime Ltd, Cheshire, UK) at the Silvatech platform, INRA Nancy-Lorraine.

Calculations

Total N amount (mg) was calculated as follows:

$$N_{\text{tot}} = \frac{\%N_{\text{total}} \times \text{DW}}{100}, \quad (1)$$

where $\%N_{\text{total}}$ is N content (% DW) and DW is dry weight (mg).

^{15}N amount (mg) in each tissue sample was calculated as follows:

$$^{15}\text{N}_{\text{amount}} = \frac{(\text{At.}\%^{15}\text{N} - \text{At.}\%^{15}\text{N}_{\text{unlabelled sample}}) \times N_{\text{tot}}}{100}, \quad (2)$$

where At. $\%^{15}\text{N}$ is isotopic abundance in samples of donor oak and receiver neighbour, defined as follows:

$$\text{At.}\%^{15}\text{N} = \frac{^{15}\text{N}}{(^{14}\text{N} + ^{15}\text{N})} \times 100. \quad (3)$$

Relative ^{15}N allocation (supp. data, Figure S1 and S2) was calculated as follows:

$$\%^{15}\text{N} = \frac{^{15}\text{N}_{\text{amount in compartment } i} \times 100}{\sum ^{15}\text{N}_{\text{amount}}}, \quad (4)$$

where $\sum ^{15}\text{N}$ amount is the sum of ^{15}N amounts in each compartment (shoots and roots in donor, receiver *Molinia*, receiver oak and in the soil).

Statistics

Statistical analysis was performed using R software (R studio, Version 1.0.153). Data are means of $n = 4$ biological replicates (\pm SE). All variables were tested for normality and homoscedasticity using the Shapiro-Wilk and Levene tests. Statistical analyses were conducted using one-way ANOVA or t test. For factors with more

than two levels (seasons, root separation treatment and compartment), ANOVA was followed by Tukey's honestly significant difference *post hoc* test for mean comparison at a 90% level.

Results

Molinia dry weight increased when mixed-grown with oak (Fig. 2 and Tables S1–S5)


Donor oak dry weight was constant across seasons either sole-grown or mixed-grown with *Molinia* (Fig. 2).

By contrast, growth in *Molinia* tufts increased appreciably from summer (7.95 ± 2.31 g for shoots and 6.62 ± 1.65 g for roots) to autumn (14.99 ± 1.11 g for shoot and 26.18 ± 4.3 g for root) in sole-grown ($p = 0.01$ for shoot and $p = 0.0007$ for roots) and then remained constant until spring (Tables S1, S2, S3 and S>4). In mixed-grown conditions, *Molinia* shoot growth increased from summer (8.11 ± 1.81 g) to autumn (24.61 ± 4.29 g, $p = 0.005$) (Table S1), but larger root biomass was only observed in winter. Remarkably, *Molinia* shoot biomass tended to be higher when grown with oak rather than sole-grown in autumn ($p = 0.07$), winter (11.06 ± 0.12 g in sole-grown and 16.7 ± 1.55 g in mixed-grown, $p = 0.04$) and spring (9.83 ± 0.8 g and 18.74 ± 2.37 g in mixed-grown, $p = 0.027$) (Fig. 2 and Tables S1–S5).

Fate of rhizodeposited N from donor oak seedlings (Figs. 3, 4 and Tables S6–S14)

Overall, about 73% of the measured ^{15}N amount in all compartments was found in donor oak, either sole-grown or mixed-grown with *Molinia* in summer (Fig.

Fig. 2 Above- and belowground dry weight for donor oak and receiver *Molinia*, for each season (summer, autumn, winter and spring). Plants were grown with either no root contact (sole-grown, white bars) or root contact (mixed-grown, grey bars). Values are reported as means \pm SE ($n = 4$). *, ** and *** correspond to $p < 0.1$, 0.05 and 0.01, respectively






Fig. 3 ^{15}N allocation (expressed as % of total ^{15}N measured in all compartments) among shoots, roots and soil in donor oak and receiver *Molinia*. Species were sole-grown (a, b, c and d) or mixed-grown (e, f, g and h). Values are reported as means \pm SE ($n = 4$)




Fig. 4 ¹⁵N allocation (expressed as % of total ¹⁵N measured in all compartments) among shoots, roots and soil in donor oak and receiver oak. Species were sole-grown (a, b, c and d) or mixed-grown (e, f, g and h). Values are reported as means ± SE (n = 4)

3a and e). ^{15}N percentage in soil was statistically similar among seasons, treatments and neighbours.

Sole-grown plants

Surprisingly, ^{15}N was detected in receiver neighbour (significantly different from zero) but at a very low level (Figs. 3 and 4 – sole-grown and Table S15). In receiver oak, ^{15}N amount was significantly different from zero in summer, autumn and winter in shoots, roots and soil respectively. In spring, both donor oak shoots and soil contained significant amounts of ^{15}N . In *Molinia*, ^{15}N amount was significantly different from zero in all cases except in roots in summer ($p = 0.17$). In donor oak, ^{15}N proportion in oak donor shoots decreased from summer to winter (from about 75% to 50%, $p = 0.004$) while that in the roots and soil rose (Figs. 3c, 4c, S1 – sole-grown and Table S15 – Receiver *Molinia*) near either receiver oak or *Molinia*.

Mixed-grown plants

When mixed-grown, ^{15}N transfer from donor oak to receiver neighbour, either *Molinia* or oak, was statistically significant. ^{15}N allocation patterns in shoots and roots were similar to that of sole-grown, but ^{15}N amount in receiver was higher (Fig. 3, 4 and S2). However, decrease in ^{15}N allocation to shoots and a concomitant increase in roots in winter were much larger when mixed-grown with receiver *Molinia* (Fig. 3e and 3g; from about 70% to 20%, $p < 0.0001$) than with receiver oak (Fig. 4e and g; from about 75% to 40%, $p < 0.001$) (Figure S1j and k). In spring, ^{15}N in donor oak was reallocated to shoots more when mixed-grown with *Molinia* ($p < 0.001$) than with another oak ($p \approx 1$). Allocation was much greater in receiver *Molinia* roots from autumn onwards than in receiver oak (Figure S2k). In autumn, ^{15}N relative amount in shoots was higher in *Molinia* than in receiver oak (Figure S2j).

Mesh (supp. data)

Overall, mesh had a very small effect on ^{15}N allocation in either donor oak, receiver oak or receiver *Molinia* (Figure S1 and S2). However, in Mesh30, ^{15}N allocation was greater in shoots in autumn in donor oak mixed-grown with another oak than in donor oak grown with *Molinia* (Figure S1g). ^{15}N allocation to shoots was much higher in receiver *Molinia* than in receiver oak

in summer and autumn (Figure S2g). ^{15}N allocation patterns in root and soil were not statistically different according to receiver oak or *Molinia* (Figure S2h, i).

Discussion

Our results show that *Molinia* biomass was rapidly favoured (within a few months) when mixed-grown with oak. Tracking ^{15}N fate validated our hypothesis that the presence of *Molinia* in the same pot with oak drove higher oak root N release for the benefit of *Molinia*.

Oak facilitated *Molinia* growth

The positive effect of oak seedlings on *Molinia* appeared in the early stage of cohabitation. When mixed-grown with oak, *Molinia* above- and belowground biomass values were greater from autumn and winter onwards, respectively, than those measured in *Molinia* sole-grown. These results are in line with previous findings that highlighted a facilitative effect of oak saplings on grass (Vernay et al. 2018). Unexpectedly, we did not find a detrimental effect of *M. caerulea* on oak seedling biomass. These results are thus not consistent with antagonistic facilitation, which had been shown in a previous experiment when *Q. petraea* and *D. cespitosa* were grown together, possibly owing to higher density of tufts in pots (Vernay et al. 2018). No significant biomass difference was observed in oak among seasons, not even when mixed-grown. Low growth may have failed to reveal an effect of *Molinia* on oak seedling growth.

N allocation pattern in oak was affected by the presence of *Molinia*

After oak shoot labelling by the cotton wick method, ^{15}N was rapidly found in oak stems and roots in summer. As expected, a higher ^{15}N allocation from shoots towards roots was observed in autumn and winter irrespective of the treatment (*i.e.* mixed-grown or sole-grown), in accordance with the conservative strategy of oaks. ^{15}N in roots was then re-allocated towards shoots to sustain bud break and early growth of emerging leaflets (Millard 1994). Interestingly, *Molinia* presence accentuated both allocations of ^{15}N to roots during winter and remobilization of ^{15}N towards shoots in oak during spring. Larger N mobilization to leaves might improve C capture, and ultimately shoot and root growth in oak mixed-grown with




Fig. 5 Correlation between dry weight and ^{15}N relative amount in receiver *Molinia* in summer (a), autumn (b), winter (c) and spring (d) and in receiver oak in summer (e), autumn (f), winter (g) and

spring (h). Dark point and regression line represent shoots and grey triangle and regression line represent roots. Equation line (y), r^2 and p -value are reported above each graph

Molinia (Bhatt et al. 2011). However, N distribution to roots was uncorrelated with concomitant biomass increment, suggesting greater N storage. Such a conservative strategy has already been reported, namely higher N allocation in oak roots than in oak shoots mixed-grown with *D. cespitosa* (Vernay et al. 2018). In spring, harvest coincided with bud bursting and early foliar development, a period generally associated with N remobilization (see Millard and Gwen.-Aelle 2010 for a review), which was particularly accentuated from roots to shoots in oak grown with *Molinia*. One explanation for this observation might be a greater soil N depletion by *Molinia*, leading to a higher demand for internal N remobilization in oak to sustain budburst and leaf development (Nielsen et al. 2000; Bausenwein et al. 2001). It has been shown that the level of N availability during bud break in oak is a critical factor affecting leaf C capture (Bazot et al. 2016; Vernay et al. 2016, 2018). Thus the magnitude of internal N recycling from roots might not depend only on limited soil inorganic N amounts but also on neighbour presence and neighbour identity (Welker et al. 1991; Kaitaniemi et al. 2018). Moreover, some internal N can be released in soil according to neighbour presence and identity (Dhamala et al. 2016).

A non-negligible part of ^{15}N is found in the soil

Nitrogen rhizodeposition processes have been described in legume and grass species and are known to occur a few days after labelling (Ribeiro Paula et al. 2015). Conversely, in fruit trees, N rhizodeposition is mainly attributed to root mortality, so ^{15}N is found in soil to a large extent in winter (Scandellari et al. 2010). However, in our study, oak seedlings exuded ^{15}N during the first weeks after labelling, and the presence of a neighbour modified N flux from roots to soil. We hypothesized that neighbours (both oak and *Molinia*) might exude specific molecular entities to increase active exudation by oak roots. Abiotic factors influencing root rhizodeposition such as soil nitrogen availability or CO_2 level (Hale et al. 1971; Philipps et al. 2006; de Graaff et al. 2007; Tückmantel et al. 2017; Bowsher et al. 2018) and interaction through root exudates (Suriyagoda et al. 2012; Semchenko et al. 2014; Mommer et al. 2016) are well-described. Recent studies on belowground signalling interactions have demonstrated that neighbour detection, response strategies and communication are mediated by root-secreted signalling chemicals (Chen et al. 2012; Rasmann and Turlings 2016; Kong et al. 2018;

Canarini et al. 2019; Huang et al. 2019). Intraspecific neighbours can improve or impair nutrient uptake of nitrogen, involving both a mutually beneficial cooperative relationship and a competitive relationship among neighbours (Hong et al. 2017). However, to our knowledge, the effect of neighbour presence and identity on nutrient rhizodeposition has not been described. This is therefore probably the first time that such a flux from roots to soil in oak-*Molinia* interference has been characterized and given an ecologically sound basis.

In winter, the increase of ^{15}N in soil was probably due to the shedding of fine roots and sloughing of living cells (Scandellari et al. 2010).

N transfer between plants may contribute to the growth facilitation process

In *Molinia*, biomass increase was positively correlated with ^{15}N relative amounts in shoots and roots (Fig. 5 and Table S16) especially in summer, autumn and winter (Fig. 5a, b and c, respectively). Conversely, the relationship was never significant for receiver oak, whatever the season. Overall, these results suggest that *Molinia* was able to obtain ^{15}N from oak for its own growth (Fig. 5e, f, g and h). These results support our hypothesis of *Molinia* growth facilitation by oak seedlings through N transfer. It emphasizes the importance of the strategy and identity of neighbours in influencing plant-plant N transfer.

Our experiment clearly shows that a significant albeit relatively modest amount of ^{15}N was transferred from oak to *Molinia* (maximum 6.5% of the total ^{15}N found in all compartments). However, we used a pulse labelling technique (at a given limited time) so that we do not know what the contribution of oak N was to the nutrition of *Molinia* over the whole season with a permanent N flux. Previous reports on the contribution of N released by a donor to total N amount in a receiver species showed that N amount varied widely according to combination of species, species age and the method used to estimate N transfer (Teste et al. 2015; Montesinos-Navarro et al. 2017). This contribution ranged from 5% (Frey and Schüepp 1993) to 80% in soybean-weed (Moyer-Henry et al. 2006). The cotton wick method was most often used on non-woody plants, but the technique can succeed on tree seedlings as demonstrated. Such transfer from plant to plant has been extensively described for carbon (Simard et al. 1997b; Simard and Perry 1997; Klein 2016) but this is the first time to our

knowledge that N transfer from a non-fixing nitrogen species to a graminoid has been evidenced at the very early stage of development in oak. This underlines the importance of nutrient exchanges, such as that of nitrogen, occurring below ground.

What are the different possible pathways of N transfer between plants?

When plants were grown with a 1 μm mesh as a barrier to roots and mycorrhizal hyphae, ^{15}N was found in receiver neighbour and soil, suggesting that some N transfer occurred as free molecules moving through the mesh (Figure S2 and Table S6). Mesh30 was designed to let through mycorrhizal hyphae, and a significant ^{15}N amount was found on the other side, at a level higher than in Mesh1, at least in autumn. This suggests that mycorrhizae might contribute to N transfers, in line with other studies (He et al. 2003, 2006, 2009; Govindarajulu et al. 2005; Jalonen et al. 2009), but our results do not enable us to conclude on the role of mycorrhizae in N transfer between oak and *Molinia*. Further experiments are now required to identify and quantify mycorrhizal networks in both oak and *Molinia* roots. We hypothesized that only vesicular arbuscular mycorrhizae (VAM) were involved in N transfer between young oak and *Molinia*, as *Molinia* is only colonized by VAM (Taylor et al. 2001). However, to our knowledge, it is not known whether *Q. petraea* is colonized by VAM, although it has been observed in other oak species (Dickie 2001).

Unexpectedly, ^{15}N was also found in the neighbouring pot in the separate-pot treatment (sole-grown). ^{15}N amounts were significantly different from natural abundance in many cases (Table S15), although the difference was small (1% at most). In receiver oak, ^{15}N was first different from zero in shoots (summer), in roots (autumn) and then in soil (winter). In spring, ^{15}N was significantly different from zero in all compartments. This N flux suggests emission of volatile nitrogen compounds from donor oak and uptake by its neighbour, first in the aboveground part. The ability of plants to take up nitrogen by leaves has been demonstrated in several studies (Wittwer and Teubner 1959; Eilers et al. 1992; Feng et al. 2015; Guo et al. 2017). After above-ground uptake, N may have been transferred to roots and translocated from roots to soil during dormancy. This spatial and temporal pathway from shoot to root, and then to soil, suggests that volatile

^{15}N was first absorbed by shoots, then transferred to roots in autumn, and rhizodeposited in winter.

In *Molinia* sole-grown, ^{15}N amount was different from zero in each condition except in roots in summer ($p < 0.15$) and in soil in winter ($p < 0.095$), supporting the hypothesis of volatile nitrogen compounds. Emissions of volatile organic compounds by plants is known (Fehsenfeld et al. 1992; Guenther et al. 1995; Holzinger et al. 2000), but only a few studies have described emissions of volatile nitrogen compounds by above-ground organs (Tukey 1966). To our knowledge it is not known whether volatile nitrogen transfer between plants occurs, or to what extent ^{15}N in receiver neighbours could derive from soil N volatilization. The belowground pathway is still the main contributor to N transfer from oak to either oak or *Molinia* (Table S17 – Ratio MG/SG). Additionally, total ^{15}N amount in all pots decreased across seasons especially between August and October (June–August: –9%; August–October: –31%; October–February: –0.4%; February–May: –18%) supporting the hypothesis of soil ^{15}N loss. Involvement of microbial communities in nitrogen transformation (Robertson and Groffman 2007; Hayatsu et al. 2008) can explain ^{15}N diminution across seasons. Although the impact of rhizodeposits on soil microorganisms has been well-studied in the past 20 years (Paterson et al. 2006; Wichern et al. 2008; Schenck Zu Schweinsberg-Mickan et al. 2012), it would be of interest to characterize more precisely (quantitatively and qualitatively) their role in rhizodeposition processes, according to species, and how it can impact on nutrient transfer (Gorka et al. 2019).

Conclusions

Our experiment clearly demonstrates N transfer from oak to *Molinia* qualitatively. However, pulse-chase labelling only enabled us to track the fate of ^{15}N supplied in summer. A continuous supply of ^{15}N throughout the whole experiment would have allowed quantification of the actual N amount effectively transferred from oak to *Molinia*. Compartmental modelling based on testing different pathways of N transfer may help identify and rank these fluxes. Soil properties and microorganisms such as mycorrhizae need to be considered in further studies to gain a better understanding of transfer processes between plants. Some substances might be emitted by roots and/or shoots: such allelopathic substances have not been well-identified and their role in such interference is under-

researched. Applying exudates from *Molinia* roots around oak seedlings to monitor any changes in growth and functioning (such as mycorrhizal symbiosis) and then analysing these exudates chemically may prove helpful in identifying allelopathic substances.

Acknowledgements The authors thank André Marquier, Christophe Serre, Pascal Walser, Camille Couteau and Brigitte Saint-Joanis for their valuable help in processing plant material.

References

- Balandier P, Collet C, Miller JH et al (2006) Designing forest vegetation management strategies based on the mechanisms and dynamics of crop tree competition by neighbouring vegetation. *Forestry* 79:3–27. <https://doi.org/10.1093/forestry/cpi056>
- Bausenwein U, Millard P, Thornton B, Raven JA (2001) Seasonal nitrogen storage and remobilization in the forb *Rumex acetosa*. *Funct Ecol* 15:370–377. <https://doi.org/10.1046/j.1365-2435.2001.00524.x>
- Bazot S, Fresneau C, Damesin C, Barthes L (2016) Contribution of previous year's leaf N and soil N uptake to current year's leaf growth in sessile oak. *Biogeosciences* 13:3475–3484. <https://doi.org/10.5194/bg-13-3475-2016>
- Bertness MD, Callaway RM (1994) Positive interactions in communities. *Tree* 9:191–193. <https://doi.org/10.1201/9780203738559>
- Bethlenfalvay GJ, Reyes-Solis MG, Camel SB, Ferrera-Cerrato R (1991) Nutrient transfer between the root zones of soybean and maize plants connected by a common mycorrhizal mycelium. *Physiol Plant* 82:423–432. <https://doi.org/10.1034/j.1399-3054.1991.820315.x>
- Bhatt MV, Khandelwal A, Dudley SA (2011) Kin recognition, not competitive interactions, predicts root allocation in young *Cakile edentula* seedling pairs. *New Phytol* 189:1135–1142. <https://doi.org/10.1111/j.1469-8137.2010.03548.x>
- Bleasdale JKA (1960) Studies on plant competition. In: Harper JL (ed) *Biology of weeds*. Oxford, England, Blackwell Science Pub, pp 133–142
- Boulter D, Jeremy JJ, Wilding M (1966) Amino acids liberated into the culture medium by pea seedling roots. *Plant Soil* 24: 121–127. <https://doi.org/10.1007/BF01373078>
- Bowsher AW, Evans S, Tiemann LK, Friesen ML (2018) Effects of soil nitrogen availability on rhizodeposition in plants: a review. *Plant Soil* 423:59–85. <https://doi.org/10.1007/s11104-017-3497-1>
- Bronstein JL (2009) The evolution of facilitation and mutualism. *J Ecol* 97:1160–1170. <https://doi.org/10.1111/j.1365-2745.2009.01566.x>
- Bruno JF, Stachowicz JJ, Bertness MD (2003) Inclusion of facilitation into ecological theory. *Trends Ecol Evol* 18:119–125. [https://doi.org/10.1016/S0169-5347\(02\)00045-9](https://doi.org/10.1016/S0169-5347(02)00045-9)

- Callaway RM (1995) Positive interactions among plants. *Bot Rev* 61:306–349. <https://doi.org/10.1007/BF02912621>
- Canarini A, Kaiser C, Merchant A et al (2019) *Corrigendum*: root exudation of primary metabolites: mechanisms and their roles in plant responses to environmental stimuli. *Front Plant Sci* 10:1–19. <https://doi.org/10.3389/fpls.2019.00420>
- Chen J, Wu F, Liu T et al (2012) Emissions of nitric oxide from 79 plant species in response to simulated nitrogen deposition. *Environ Pollut* 160:192–200. <https://doi.org/10.1016/j.envpol.2011.09.007>
- Clements FE, Hanson HC, Weaver JE (1929) Plant competition; an analysis of community functions. 340 pp
- Coll L, Balandier P, Picon-Cochar C (2004) Morphological and physiological responses of beech (*Fagus sylvatica*) seedlings to grass-induced belowground competition. *Tree Physiol* 24: 45–54
- Craine JM (2005) Reconciling plant strategy theories of Grime and Tilman. *J Ecol* 93:1041–1052. <https://doi.org/10.1111/j.1365-2745.2005.01043.x>
- Craine JM, Dybzinski R (2013) Mechanisms of plant competition for nutrients, water and light. *Funct Ecol* 27:833–840. <https://doi.org/10.1111/1365-2435.12081>
- de Graaff M-A, Six J, van Kessel C (2007) Elevated CO₂ increases nitrogen rhizodeposition and microbial immobilization of root-derived nitrogen. *New Phytol* 173:778–786. <https://doi.org/10.1111/j.1469-8137.2006.01974.x>
- Dhamala N, Rasmussen J, Carlsson G et al (2016) N transfer in three-species grass-clover mixtures with chicory, ribwort plantain or caraway. *Plant Soil* 413:1–14. <https://doi.org/10.1007/s11104-016-3088-6>
- Dickie IA (2001) Vesicular-arbuscular mycorrhizal infection of *Quercus rubra* seedlings. *New Phytol* 151:257–264. <https://doi.org/10.1046/j.1469-8137.2001.00148.x>
- Eilers G, Brunnmc R, Matznep E (1992) Above-ground N-uptake from wet deposition by Norway spruce (*Picea abies* Karst.). *For Ecol Manage* 51:239–247. [https://doi.org/10.1016/0378-1127\(92\)90489-V](https://doi.org/10.1016/0378-1127(92)90489-V)
- Eissenstat DM (1990) A comparison of phosphorus and nitrogen transfer between plants of different phosphorus status. *Oecologia* 82:342–347. <https://doi.org/10.1007/BF00317481>
- Ek H, Anderson S, Söderström B (1996) Carbon and nitrogen flow in silver birch and Norway spruce connected by a common mycorrhizal mycelium. *Mycorrhiza* 6:465–467
- Epstein E (1961) The essential role of calcium in selective cation transport by plant cells. *Plant Physiol* 36:437–444. <https://doi.org/10.1104/pp.36.4.437>
- Fehsenfeld F, Calvert J, Fall R et al (1992) Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry. *Global Biogeochem Cycles* 6:389–430. <https://doi.org/10.1029/92GB02125>
- Feng W, Guo B, Zhang H et al (2015) Remote estimation of above ground nitrogen uptake during vegetative growth in winter wheat using hyperspectral red-edge ratio data. *F Crop Res* 180:197–206. <https://doi.org/10.1016/j.fcr.2015.05.020>
- Fernandez C, Monnier Y, Santonja M et al (2016) The impact of competition and allelopathy on the trade-off between plant defense and growth in two contrasting tree species. *Front Plant Sci* 7:1–14. <https://doi.org/10.3389/fpls.2016.00594>
- Frey B, Schüepp H (1993) A role of vesicular-arbuscular (VA) mycorrhizal fungi in facilitating interplant nitrogen transfer. *Soil Biol Biochem* 25:651–658. [https://doi.org/10.1016/0038-0717\(93\)90104-J](https://doi.org/10.1016/0038-0717(93)90104-J)
- Friedrich JM, Dawson J (1984) Soil nitrogen concentration and *Juglans nigra* growth in mixed plots with nitrogen-fixing *Alnus*, *Elaeagnus*, *Lespedeza* and *Robinia* species. *Can J For Res* 14:864–868
- Fustec J, Lesuffleur F, Mahieu S, Cliquet J (2010) Nitrogen rhizodeposition of legumes. A review. *Agron Sustain Dev* 30:57–66. <https://doi.org/10.1051/agro/2009003>
- Gargaglione V, Peri PL, Rubio G (2014) Tree-grass interactions for N in *Nothofagus antarctica* silvopastoral systems: Evidence of facilitation from trees to underneath grasses. *Agrofor Syst* 88:779–790. <https://doi.org/10.1007/s10457-014-9724-3>
- Gorka S, Dietrich M, Mayerhofer W et al (2019) Rapid transfer of plant photosynthates to soil bacteria via ectomycorrhizal hyphae and its interaction with nitrogen availability. *Front Microbiol* 10:1–20. <https://doi.org/10.3389/FMICB.2019.00168>
- Govindarajulu M, Pfeffer PE, Jin H et al (2005) Nitrogen transfer in the arbuscular mycorrhizal symbiosis. *Nature* 435:819–823. <https://doi.org/10.1038/nature03610>
- Grime JP (1974) Vegetation classification by reference to strategies. *Nature* 250:26–31. <https://doi.org/10.1038/252497a0>
- Grove S, Haubensak KA, Parker IM (2013) Direct and indirect effects of allelopathy in the soil legacy of an exotic plant invasion. *Plant Ecol* 213:1869–1882. <https://doi.org/10.1007/s11258-012-0079-4>
- Guenther A, Nicholas Hewitt C, David E et al (1995) A global model of natural volatile organic compound emissions. *J Geophys Res* 100:8873–8892. <https://doi.org/10.1029/94JD02950>
- Guo B, Qi S, Heng Y et al (2017) Remotely assessing leaf N uptake in winter wheat based on canopy hyperspectral red-edge absorption. *Eur J Agron* 82:113–124. <https://doi.org/10.1016/j.eja.2016.10.009>
- Hale MG, Foy C, Shay FJ (1971) Factors affecting root exudation. *Adv Agron* 23:89–109. [https://doi.org/10.1016/S0065-2113\(08\)60151-0](https://doi.org/10.1016/S0065-2113(08)60151-0)
- Hayatsu M, Tago K, Saito M (2008) Various players in the nitrogen cycle: diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Sci Plant Nutr* 54:33–45. <https://doi.org/10.1111/j.1747-0765.2007.00195.x>
- Haystead A, Marriott C (1978) Fixation and transfer of nitrogen in a white clover-grass sward under hill conditions. *Proc Assoc Appl Biol* 3:453–457. <https://doi.org/10.1111/j.1744-7348.1978.tb00740.x>
- Haystead A, Marriott C (1979) Transfer of legume nitrogen to associated grass. *Soil Biol Biochem* 11:99–104. [https://doi.org/10.1016/0038-0717\(79\)90083-X](https://doi.org/10.1016/0038-0717(79)90083-X)
- Haystead A, Malajczuk N, Grove TS (1988) Underground transfer of nitrogen between pasture plants infected with vesicular-arbuscular mycorrhizal fungi. *New Phytol* 108:417–423. <https://doi.org/10.1111/j.1469-8137.1988.tb04182.x>
- He X-H, Critchley C, Bledsoe C (2003) Nitrogen transfer within and between plants through common mycorrhizal networks (CMNs). *CRC Crit Rev Plant Sci* 22:531–567. <https://doi.org/10.1080/713608315>
- He X, Bledsoe CS, Zasoski RJ et al (2006) Rapid nitrogen transfer from ectomycorrhizal pines to adjacent ectomycorrhizal and

- arbuscular mycorrhizal plants in a California oak woodland. *New Phytol* 170:143–151. <https://doi.org/10.1111/j.1469-8137.2006.01648.x>
- He X, Xu M, Qiu GY, Zhou J (2009) Use of ^{15}N stable isotope to quantify nitrogen transfer between mycorrhizal plants. *J Plant Ecol* 2:107–118. <https://doi.org/10.1093/jpe/rtp015>
- He Y, Comelissen JHC, Wang P et al (2019) Nitrogen transfer from one plant to another depends on plant biomass production between conspecific and heterospecific species via a common arbuscular mycorrhizal network. *Environ Sci Pollut Res* 26:8828–8837. <https://doi.org/10.1007/s11356-019-04385-x>
- Henzell EF, Martin AE, Ross PJ, Haydock KP (1964) Isotopic studies on the uptake of sulphur by pasture plants. *Aust J Agric Res* 15:876–884. <https://doi.org/10.1080/00288233.1977.10427327>
- Holzinger R, Sandoval-Soto L, Rottenberger S et al (2000) Emissions of volatile organic compounds from *Quercus ilex* L. measured by proton transfer reaction mass spectrometry under different environmental conditions. *J Geophys Res Atmos* 105:573–579. <https://doi.org/10.1029/2000jd900296>
- Hong J, Ma X, Zhang X, Wang X (2017) Nitrogen uptake pattern of herbaceous plants: coping strategies in altered neighbor species. *Biol Fertil Soils* 53:729–735. <https://doi.org/10.1007/s00374-017-1230-0>
- Huang W, Gfeller V, Erb M (2019) Root volatiles in plant-plant interactions II: Root volatiles alter root chemistry and plant-herbivore interactions of neighboring plants. *Plant Cell Environ* 42:1946–1973. <https://doi.org/10.1101/441790>
- Hunter AF, Aarssen LW (1988) Plants helping plants: new evidence indicates that beneficence is important in vegetation. *Am Inst Biol Sci* 38:34–40. <https://doi.org/10.2307/1310644>
- Jalonen R, Nygren P, Sierra J (2009) Transfer of nitrogen from a tropical legume tree to an associated fodder grass via root exudation and common mycelial networks. *Plant, Cell Environ* 32:1366–1376. <https://doi.org/10.1111/j.1365-3040.2009.02004.x>
- Jensen ES (1996) Rhizodeposition of N by pea and barley and its effect on soil N dynamics. *Soil Biol Biochem* 28:65–71. [https://doi.org/10.1016/0038-0717\(95\)00116-6](https://doi.org/10.1016/0038-0717(95)00116-6)
- Johansen A, Jensen ES (1996) Transfer of N and P from intact or decomposing roots of pea to barley interconnected by an arbuscular mycorrhizal fungus. *Soil Biol Biochem* 28:73–81. [https://doi.org/10.1016/0038-0717\(95\)00117-4](https://doi.org/10.1016/0038-0717(95)00117-4)
- Kaitaniemi P, Lintunen A, Sievänen R, Perttunen J (2018) Computational analysis of the effects of light gradients and neighbouring species on foliar nitrogen. *Ecol Inform* 48:171–177. <https://doi.org/10.1016/j.ecoinf.2018.09.009>
- Khan WDF, Peoples MB, Herridge DF (2002) Quantifying below-ground nitrogen of legumes. *Plant Soil* 245:327–334. <https://doi.org/10.1023/A:1020407006212>
- Klein T (2016) Belowground carbon trade among talltrees in a temperate forest. *Science* (80-) 352:342–344. <https://doi.org/10.6084/m9.figshare.3081127>
- Kohli RK, Batish D, Singh HP (1998) Allelopathy and its implications in agroecosystems. *J Crop Prod* 1:169–202. https://doi.org/10.1300/j144v01n01_08
- Kong C-H, Zhang S-Z, Li Y-H et al (2018) Plant neighbor detection and allelochemical response are driven by root-secreted signaling chemicals. *Nat Commun* 9:1–9. <https://doi.org/10.1038/s41467-018-06429-1>
- Li X, Rennenberg H, Simon J (2015) Competition for nitrogen between *Fagus sylvatica* and *Acer pseudoplatanus* seedlings depends on soil nitrogen availability. *Front Plant Sci* 6:1–11. <https://doi.org/10.3389/fpls.2015.00302>
- Mahieu S, Fustec J, Faure ML et al (2007) Comparison of two ^{15}N labelling methods for assessing nitrogen rhizodeposition of pea. *Plant Soil* 295:193–205. <https://doi.org/10.1007/s11104-007-9275-8>
- Mahieu S, Fustec J, Jensen ES, Crozat Y (2009) Does labelling frequency affect N rhizodeposition assessment using the cotton-wick method? *Soil Biol Biochem* 41:2236–2243. <https://doi.org/10.1016/j.soilbio.2009.08.008>
- Mallik A (1998) Allelopathy and competition in coniferous forests. *Environ For Sci* 54:309–315. https://doi.org/10.1007/978-94-011-5324-9_33
- Millard P (1994) Measurement of the remobilization of nitrogen for spring leaf growth of trees under field conditions. *Tree Physiol* 14:1049–1054. <https://doi.org/10.1093/treephys/14.7-8-9.1049>
- Millard P, Gwen.-Aelle G (2010) Nitrogen storage and remobilization by trees : ecophysiological relevance in a changing world. *Tree Physiol* 30:1083–1095
- Molisch H (1937) Der Einfluss einer pflanze auf die andere, allelopathie. *Nature* 141:493
- Mommer L, Kirkegaard J, van Ruijven J (2016) Root–root interactions: towards a rhizosphere framework. *Trends Plant Sci* 21:209–217. <https://doi.org/10.1016/j.tplants.2016.01.009>
- Montesinos-Navarro A, Verdú M, Quejeto JA, Valiente-Banuet A (2017) Nurse plants transfer more nitrogen to distantly related species. *Ecology* 98:1300–1310. <https://doi.org/10.1002/ecy.1771>
- Moyer-Henry KA, Burton JW, Israel DW, Rufty TW (2006) Nitrogen transfer between plants: A ^{15}N natural abundance study with crop and weed species. *Plant Soil* 282:7–20
- Muller CH (1969) Allelopathy as a factor in ecological process. *Vegetatio* 18:348–357. <https://doi.org/10.1007/BF00332847>
- Neilsen D, Millard P, Herbert LC et al (2000) Remobilization and uptake of N by newly planted apple (*Malus domestica*) trees in response to irrigation method and timing of N application. *Tree Physiol* 21:513–521. <https://doi.org/10.1093/treephys/21.8.513>
- Paterson E, Gebbing T, Abel C et al (2006) Rhizodeposition shapes rhizosphere microbial community structure in organic soil. *New Phytol* 173:600–610. <https://doi.org/10.1111/j.1469-8137.2006.01931.x>
- Philipps DA, Fox TC, Six J (2006) Root exudation (net efflux of amino acids) may increase rhizodeposition under elevated CO_2 . *Glob Chang Biol* 12:561–567. <https://doi.org/10.1111/j.1365-2486.2006.01100.x>
- Pierik R, Mommer L, Voeseek LA (2013) Molecular mechanisms of plant competition: neighbour detection and response strategies. *Funct Ecol* 27:841–853. <https://doi.org/10.1111/1365-2435.12010>
- Rasmann S, Turlings TCJ (2016) Root signals that mediate mutualistic interactions in the rhizosphere. *Curr Opin Plant Biol* 32:62–68. <https://doi.org/10.1016/j.pbi.2016.06.017>
- Ribeiro Paula R, Bouillet JP, Ocheuze Trivelin PC et al (2015) Evidence of short-term belowground transfer of nitrogen from *Acacia mangium* to *Eucalyptus grandis* trees in a tropical planted forest. *Soil Biol Biochem* 91:99–108. <https://doi.org/10.1016/j.soilbio.2015.08.017>

- Rice EL (1985) Chemically mediated interactions between plants and other organisms. *Chem Mediat Interact between Plants Other Org* 19:81–105. <https://doi.org/10.1007/978-1-4757-9658-2>
- Robertson GP, Groffman PM (2007) Nitrogen transformations. In: *Soil microbiology, ecology and biochemistry*, Third Edit. Elsevier Inc., pp 341–364
- Rovira AD (1969) Plant root exudates. *Bot Rev* 35:35–57. <https://doi.org/10.1007/BF02859887>
- San Emeterio L, Damgaard C, Canals RM (2007) Modelling the combined effect of chemical interference and resource competition on the individual growth of two herbaceous populations. *Plant Soil* 292:95–103. <https://doi.org/10.1007/s11104-007-9205-9>
- Scandellari F, Ventura M, Gioacchini P et al (2010) Seasonal pattern of net nitrogen rhizodeposition from peach (*Prunus persica* (L.) Batsch) trees in soils with different textures. *Agric Ecosyst Environ* 136:162–168. <https://doi.org/10.1016/j.agee.2009.12.017>
- Schenck Zu Schweinsberg-Mickan M, Jörgensen RG, Müller T (2012) Rhizodeposition: its contribution to microbial growth and carbon and nitrogen turnover within the rhizosphere. *J Plant Nutr Soil Sci* 175:750–760. <https://doi.org/10.1002/jpln.201100300>
- Schöb C, Prieto I, Armas C, Pugnaire FI (2014) Consequences of facilitation: one plant's benefit is another plant's cost. *Funct Ecol* 28:500–508. <https://doi.org/10.1111/1365-2435.12185>
- Schreiner O, Reed HS (1907) The production of deleterious excretions by roots. *Bull Torrey Bot Club* 34:279–303. <https://doi.org/10.2307/2479157>
- Semchenko M, Saar S, Lepik A (2014) Plant root exudates mediate neighbour recognition and trigger complex behavioural changes. *New Phytol* 204:631–637. <https://doi.org/10.1111/nph.12930>
- Shamoot S, McDonald L, Bartholomew WV (1968) Rhizo-deposition of organic debris in soil. *Soil Sci Soc Am Proc* 32:817–820. <https://doi.org/10.2136/sssaj1968.03615995003200060031x>
- Simard SW, Perry DA (1997) Net transfer of carbon between ectomycorrhizal tree species in the field. *Nature* 388:579. <https://doi.org/10.1038/41557>
- Simard SW, Durall DM, Jones MD (1997a) Carbon allocation and carbon transfer between *Betula papyrifera* and *Pseudotsuga menziesii* seedlings using a ¹³C pulse-labeling method. *Plant Soil* 191:41–55. <https://doi.org/10.1023/A:1004205727882>
- Simard SW, Jones MD, Durall DM et al (1997b) Reciprocal transfer of carbon isotopes between ectomycorrhizal *Betula papyrifera* and *Pseudotsuga menziesii*. *New Phytol* 137:529–542. <https://doi.org/10.1046/j.1469-8137.1997.00834.x>
- Slankis V, Runeckles VC, Krotkov G (1964) Metabolites liberated by roots of white pine (*Pinus strobus* L.) seedlings. *Physiol Plant* 17:301–313. <https://doi.org/10.1111/j.1399-3054.1964.tb08162.x>
- Stagnari F, Maggio A, Galieni A, Pisante M (2017) Multiple benefits of legumes for agriculture sustainability: an overview. *Chem Biol Technol Agric* 4:1–13. <https://doi.org/10.1186/s40538-016-0085-1>
- Suriyagoda LDB, Lambers H, Renton M, Ryan MH (2012) Growth, carboxylate exudates and nutrient dynamics in three herbaceous perennial plant species under low, moderate and high phosphorus supply. *Plant Soil* 358:105–117. <https://doi.org/10.1007/s11104-012-1311-7>
- Taylor K, Rowland AP, Jones HE (2001) *Molinia caerulea* (L.) Moench. *J Ecol* 89:126–144. <https://doi.org/10.1046/j.1365-2745.2001.00534.x>
- Teste FP, Veneklaas EJ, Dixon KW, Lambers H (2015) Is nitrogen transfer among plants enhanced by contrasting nutrient-acquisition strategies? *Plant, Cell Environ* 38:50–60. <https://doi.org/10.1111/pce.12367>
- Tilman D (1990) Constraints and tradeoffs: toward a predictive theory of competition and succession. *Oikos* 58:3–15. <https://doi.org/10.2307/3565355>
- Tückmantel T, Leuschner C, Preusser S et al (2017) Root exudation patterns in a beech forest: dependence on soil depth, root morphology, and environment. *Soil Biol Biochem* 107:188–197. <https://doi.org/10.1016/j.soilbio.2017.01.006>
- Tukey HBJ (1966) Leaching of metabolites from above-ground plant parts and its implications. *Torrey Bot Soc* 93:385–401. <https://doi.org/10.2307/2483411>
- Vallis I, Haydock KP, Ross PJ, Henzell EF (1967) Isotopic studies on the uptake of nitrogen by pasture plants. *Aust J Agric Res* 18:865–877. <https://doi.org/10.1071/AR9730693>
- van Kessel C, Singleton PW, Hoben HJ (1985) Enhanced N-transfer from a soybean to maize by vesicular arbuscular mycorrhizal (VAM) fungi. *Ecology* 66:562–563. <https://doi.org/10.1111/jpp.79.2.562>
- Vernay A, Balandier P, Guinard L et al (2016) Photosynthesis capacity of *Quercus petraea* (Matt.) saplings is affected by *Molinia caerulea* (L.) under high irradiance. *For Ecol Manage* 376:107–117. <https://doi.org/10.1016/j.foreco.2016.05.045>
- Vernay A, Malagoli P, Fernandez M et al (2018) Improved *Deschampsia cespitosa* growth by nitrogen fertilization jeopardizes *Quercus petraea* regeneration through intensification of competition. *Basic Appl Ecol* 31:21–32. <https://doi.org/10.1016/j.baae.2018.06.002>
- Virtanen, Artturi I, Laine T (1939) LIII. Investigations on the root nodule bacteria of leguminous plants XXII. The excretion products of root nodules. The mechanism of N-fixation. In: *The mechanism of N-Fixation*. pp 412–427
- Welker JM, Gordon DR, Rice KJ (1991) Capture and allocation of nitrogen by *Quercus douglasii* seedlings in competition with annual and perennial grasses. *Oecologia* 87:459–466. <https://doi.org/10.1007/BF00320407>
- Whitney AS, Kanehiro Y, Sherman GD (1967) Nitrogen relationships of three tropical forage legumes in pure stands and in grass mixtures. *Agron J* 59:47–50. <https://doi.org/10.2134/agronj1967.00021962005900010014x>
- Wichern F, Eberhardt E, Mayer J et al (2008) Nitrogen rhizodeposition in agricultural crops: methods, estimates and future prospects. *Soil Biol Biochem* 40:30–48. <https://doi.org/10.1016/j.soilbio.2007.08.010>
- Wittwer SH, Teubner FG (1959) Foliar absorption of mineral nutrients. *Annu Rev Plant Physiol* 10:13–30. <https://doi.org/10.1146/annurev.pp.10.060159.000305>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.