

Primary culture of trout hepatocytes: a cellular system for studying induction of P450 enzymes and alteration of DNA by aquatic contaminants

Alain Devaux, Jean-François Masfaraud, Gilles Monod

▶ To cite this version:

Alain Devaux, Jean-François Masfaraud, Gilles Monod. Primary culture of trout hepatocytes : a cellular system for studying induction of P450 enzymes and alteration of DNA by aquatic contaminants. 40. Meeting of the European Tissue Culture Society, Institut National de la Santé et de la Recherche Médicale (INSERM). Rennes, FRA., 1993. hal-02775837

HAL Id: hal-02775837 https://hal.inrae.fr/hal-02775837

Submitted on 4 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOF¢TEC



THE 40th MEETING OF THE EUROPEAN TISSUE CULTURE SOCIETY 4-8 JULY 1993 - RENNES - FRANCE 40^e CONGRES DE L'EUROPEAN TISSUE CULTURE SOCIETY 4-8 JUILLET 1993 - RENNES - FRANCE CHAMBRE DE COMMERCE ET D'INDUSTRIE - 2 AVENUE DE LA PREFECTURE



PRIMARY CULTURE OF TROUT HEPATOCYTES : A CELLULAR SYSTEM FOR STUDYING INDUCTION OF P450 ENZYMES AND ALTERATION OF DNA BY AQUATIC CONTAMINANTS.

A. Devaux, J.F. Masfaraud[§] and G. Monod^{*}, Laboratoire des Sciences de l'Environnement, ENTPE, 69518 VAULX EN VELIN Cedex, [§]69007 LYON and ^{*}INRA, équipe d'Ecotoxicologie, 35042 RENNES Cedex France.

Xenobiotic metabolism is mainly carried out in the liver. Cytochrome P450 dependent monooxygenases play a major role in the biotransformation of lipophilic xenobiotics among which environmental pollutants are highly represented. Furthermore planar xenobiotics such as polyaromatic hydrocarbons (PAHs) and dioxines are potent *in vivo* inducers of P450 1A1. Recently induction of P450 1A1 has been proposed as a biomarker of exposure to dioxine-like compounds. P450 1A1 has also been demonstrated to transform PAHs into DNA damaging species.

In this work, the induction of P450 1A1 (measured through the increase of ethoxyresorufin *O*-deethylase (EROD) activity) by the model inducer betanaphthoflavone (BNF), and formation of DNA adducts after exposure to benzo(a)pyrene (B(a)P, a model PAH) were studied in rainbow trout *in vivo* and in primary culture of hepatocytes.

Induction experiment (100 mg BNF/kg body weight i.p. and 0,1mg BNF/ml culture medium, 24h after cell isolation) showed a similar pattern of EROD induction both *in vivo* and in hepatocytes, with an induction peak 72h after BNF administration (5-fold and 2.5-fold induction *in vivo* and in hepatocytes, respectively).

Hypothesizing the involvement of regulating compounds in induction of P450 1A1 by BNF, influence of glucocorticoids on EROD induction was studied in trout hepatocyte culture. A 5- to 6-fold potentiation of EROD induction by BNF was obtained in hepatocytes exposed to cortisol and dexamethasone.

Exposure to B(a)P (80 mg/kg i.p. *in vivo* and 0,252 µg/ml in cultured hepatocytes) resulted in similar pattern of DNA adducts *in vivo* and in hepatocytes.

These results showed that primary culture of trout hepatocytes seems to be a promising cellular system for studying the induction of P450 1A1 and alteration of DNA by aquatic contaminants and for evaluating the ecotoxicity potential of samples (water, sediments) from the aquatic environment.