



# Isolation and culture of cells from the pituitary of the rainbow trout: immunological and physiological characterization of the purified cells

Claudine Weil, Elisabeth Sambroni, Pierre-Yves Le Bail, Maurice Loir

## ► To cite this version:

Claudine Weil, Elisabeth Sambroni, Pierre-Yves Le Bail, Maurice Loir. Isolation and culture of cells from the pituitary of the rainbow trout: immunological and physiological characterization of the purified cells. 2. International Symposium on Fish Endocrinology, Jun 1992, Saint-Malo, France. 116 p., 1992. hal-02775907

**HAL Id: hal-02775907**

**<https://hal.inrae.fr/hal-02775907>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**2<sup>nd</sup>**

**INTERNATIONAL  
SYMPOSIUM  
on FISH  
ENDOCRINOLOGY**

*Abstracts*

**PALAIS DU GRAND LARGE**

**SAINT-MALO**

**JUNE 1 - 4 1992**



VILLE  
DE  
SAINT-MALO



ISOLATION AND CULTURE OF CELLS FROM THE PITUITARY OF THE  
RAINBOW TROUT : IMMUNOLOGICAL AND PHYSIOLOGICAL  
CHARACTERIZATION OF THE PURIFIED CELLS

Claudine WEIL, Elisabeth SAMBRONI, Pierre-Yves LE BAIL and  
Maurice LOIR

Laboratoire de Physiologie des Poissons, INRA, Campus de  
Beaulieu, 35042 Rennes Cedex, France

In order to study the possible paracrine interactions between gonadotrophs (GtH cells) and somatotrophs (GH cells), the isolation of enriched populations of each cell type was undertaken. The separation was based on the difference in density of the cells using Percoll as the density gradient-forming agent.

Using collagenase dispersed pituitary cells from spermiating rainbow trout, it was possible to obtain a 85% to 65%-enriched population of somatotrophs (density 1.081-1.063) containing 11% of gonadotrophs as assessed by their immunological identification. On the other hand, using the same technique, the purification of one population of GtH cells was not possible in spermiating males or early vitellogenic females. In both cases, GtH cells appear to have variable densities (varying from 1.081 to 1.040) which might correspond to different degrees of maturity as described in mammals.

The separated cells are able to be cultured and to respond to specific secretagogue. In the case of GH cell enriched population, somatostatine induced a decrease in growth hormone (GH) release while salmon gonadotropin releasing hormone (sGnRH) increased GtH release from the GtH cells contained in the different cell fractions.