



Development of a protein binding assay for teleost IGF: Relationships between GH and IGF in the serum of rainbow trout (*Oncorhynchus mykiss*)

Ping-De Niu, Michel Binoux, Jaume Perez-Sanchez, Pierre-Yves Le Bail

► To cite this version:

Ping-De Niu, Michel Binoux, Jaume Perez-Sanchez, Pierre-Yves Le Bail. Development of a protein binding assay for teleost IGF: Relationships between GH and IGF in the serum of rainbow trout (*Oncorhynchus mykiss*). 2. International Symposium on Fish Endocrinology, Jun 1992, Saint-Malo, France. 116 p. hal-02776403

HAL Id: hal-02776403

<https://hal.inrae.fr/hal-02776403>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



2nd

**INTERNATIONAL
SYMPOSIUM
on FISH
ENDOCRINOLOGY**

Abstracts

PALAIS DU GRAND LARGE

SAINT-MALO

JUNE 1 - 4 1992



MINISTÈRE
DE LA RECHERCHE
ET DE LA TECHNOLOGIE



VILLE
DE
SAINT-MALO



DEVELOPMENT OF A PROTEIN BINDING ASSAY FOR TELEOST IGF: RELATIONSHIPS BETWEEN GH AND IGF IN THE SERUM OF RAINBOW TROUT (*Oncorhynchus mykiss*).

Ping-De NIU, Michel BINOUX, Jaime PEREZ-SANCHEZ and Pierre-Yves LE BAIL

Laboratoire de Physiologie des Poissons, INRA, Campus de Beaulieu, 35042 Rennes FRANCE.

In rainbow trout, we have recently described the presence of a serum protein (tIGFBP) which specifically binds human IGF. Using a tIGFBP preparation, we have developed a binding assay to measure the plasma IGF level in different teleost species.

Before the assay, the binding protein of serum or tissue extracts was discarded using SP Sephadex G-25 in acidic conditions. After this treatment, contamination of the IGF fraction by IGFBP estimated by binding assay, was around 5 %, and was not detectable by a western ligand blot technic.

Using human IGF1 as standard and labelled hormone, sensitivity of the assay was 0.15 ng/ml (ED90) with an ED50 of about 1.3 ng/ml. Increasing concentrations of hIGF1 added in trout serum were fully recovered by the assay. hIGF2 crossreaction was partial and no significant displacement was observed with Insulin from different species or with other hormones. Inhibition curves obtained with serum IGF fractions from teleost and mammals are parallel to the standard curve which suggests that the binding sites of IGF are well conserved during evolution. No significant displacement was obtained with extracts from tissues such as spleen, liver, kidney, brain and muscle. Using this tIGF binding assay, IGF was measured in parallel with GH in serum from young trout sampled every 1.5 hour during one day. The daily profiles for both hormones, which appear pulsatile, are similar but not completely synchronous. Significant correlation was observed between GH and IGF levels but changes in IGF were delayed 1.5 hour. Analogous observations were obtained in catheterized adult trout. Although serum GH levels differ greatly between fish, less variability is found in IGF. This suggests that IGF secretion depends on the level of serum GH and that the receptivity to GH varies between fish. To verify this hypothesis, trout were submitted to starvation or to bovine GH treatment for four weeks. In starved fish, in which serum GH levels increased in parallel to a decrease in the number of liver GH receptors, the serum IGF level was significantly lower than in fed fish. In bGH injected fish, the serum IGF level was significantly higher than in non-injected fish.

These results are in accordance, with those obtained in mammals in which liver IGF1 secretion is controlled by GH and modulated by the nutritional status.