Influence of triploid status on salmon smoltification

Gilles Boeuf, H. Seddiki, Anne-Claire Le Roux, Armelle Severe, Pierre-Yves
Le Bail

► To cite this version:

Gilles Boeuf, H. Seddiki, Anne-Claire Le Roux, Armelle Severe, Pierre-Yves Le Bail. Influence of triploid status on salmon smoltification. 4. International Workshop on Salmonid Smoltification, Oct 1992, St Andrews, New Brunswick, Canada. 10.1016/0044-8486(94)90035-3 . hal-02777974

HAL Id: hal-02777974

<https://hal.inrae.fr/hal-02777974>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Influence of triploid status on salmon smoltification

Gilles Boeuf^{a,*}, Hachim Seddiki^a, Annick Le Roux^a, Armelle Severe^a,
Pierre-Yves Le Bail^b

^aIFREMER, BP 70, 29280 Plouzane, France

^bINRA, Campus de Beaulieu, 35042 Rennes, France

Abstract

Growth and survival in fresh water (FW), completion of smoltification, and subsequent growth in seawater (SW) of diploid (2n) and triploid (3n) Atlantic salmon were compared. 3n eggs were produced by thermal shock (10 min at 29°C) and verified by caryological examination of young embryos (86% survival) and, later on 9-month-old juveniles, by measurement of erythrocyte nuclear diameter (96% 3n). Parr were reared in four tanks (Le Conquet hatchery) under natural temperature and photoperiod (48°N). Survival from first-feeding (20 March) to smolting (mid-April of the next year) was 84.3% in the 2n and 81.4% in the 3n fish. At completion of smolting, the triploid salmon were significantly larger than diploid fish (171 ± 2 vs 163 ± 2 mm; $P < 0.01$) but both groups exhibited the same condition factor (1.04 ± 0.03). Gill microsomal Na^+, K^+ -ATPase activity showed similar patterns in both groups, increasing from mid-February, both peaking on 21 April at $30.1 \pm 2.2 \mu\text{mol PO}_4 \cdot \text{mg prot}^{-1} \cdot \text{h}^{-1}$. In contrast, 2n and 3n fish exhibited significant differences in plasma thyroid hormones (T_3 and T_4) and growth hormone (GH) levels. T_3 increased in 2n fish at the end of March, reaching $12.5 \pm 1 \text{ ng} \cdot \text{ml}^{-1}$, but in 3n fish remained low and varied little after the beginning of March ($1.5 \text{ ng} \cdot \text{ml}^{-1}$). T_4 surged in 2n salmon on 28 March ($19.5 \pm 1.5 \text{ ng} \cdot \text{ml}^{-1}$), but peaked later in 2n fish (15 April) and at a lower level ($10.2 \pm 1.2 \text{ ng} \cdot \text{ml}^{-1}$, $P < 0.01$). GH levels increased in 2n fish in late March, reaching $12.2 \pm 1 \text{ ng} \cdot \text{ml}^{-1}$ on 4 April, significantly ($P < 0.01$) higher than the 3n salmon ($8.0 \pm 0.5 \text{ ng} \cdot \text{ml}^{-1}$).

Both 2n and 3n groups were transferred directly to 35‰ SW on 18 April. Both groups showed good hypo-osmoregulatory ability. Plasma osmolality and chloride after 48 h in SW were $362 \pm 2 \text{ mOsm} \cdot \text{l}^{-1}$ and $148 \pm 2 \text{ mM} \cdot \text{l}^{-1}$ in 2n and 360 ± 2 and 149 ± 4 in 3n smolts. Mortality after 21 days in SW was 3% (2n) and 4% (3n).

Long-term results in net pens showed better performance in 2n than in 3n salmon in SW. In June, after 14 months in SW, survival of 2n salmon was 70% and only 45% for 3n fish. At this time, 2n and 3n salmon weighed 1.568 and 1.420 kg, respectively. Although triploid Atlantic salmon are used in Scotland, Canada and Tasmania, the seawater rearing of 3n salmon in France may not be practicable because of local environmental conditions. Triploids appeared able to adapt to SW but their growth and survival were inferior to those of 2n fish. The endocrinological differences between 2n and 3n fish may be responsible for relatively poor performance of 3n salmon in SW.

*Corresponding author.

Special Issue

Salmonid Smoltification IV

**Proceedings of a Workshop hosted by the Department of Fisheries and Oceans,
Biological Station, St. Andrews, New Brunswick, Canada**

**Held at the Harry B. Hachey Conference Centre, St. Andrews Biological Station,
19-23 October 1992**

Guest Editors:

RICHARD L. SAUNDERS

Department of Fisheries and Oceans, Biological Station, St. Andrews, N.B. E0G 2X0, Canada

TILLMANN J. BENFEY

Department of Biology, University of New Brunswick, Fredericton, N.B. E3B 6E1, Canada

TERRANCE M. BRADLEY

Department of Fisheries, Animal and Veterinary Sciences, University of Rhode Island, Kingston,
RI 02881, USA

JAMES DUSTON

Connors Bros. Limited, Aquaculture Division, Blacks Harbour, N.B. E0G 1H0, Canada

GIL J. FARMER

Department of Fisheries and Oceans, Fish Culture Section, P.O. Box 550, Halifax, N.S. B3J 2S7,
Canada

STEPHEN D. McCORMICK

Northeast Anadromous Fish Research Laboratory, U.S. Fish and Wildlife Service, P.O. Box 796,
Turners Falls, MA 01376, USA

JENNIFER L. SPECKER

Department of Zoology, University of Rhode Island, Kingston, RI 02881, USA

Section Editor, Physiology and Endocrinology:

NIALl BROMAGE

Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK