



HAL
open science

In vitro gonadal steroidogenesis in relation to sex inversion in the protandrous seabass *Lates calcarifer*

Yann Guiguen, Alexis Fostier, - Aquacop (pyf), Bernard Jalabert

► **To cite this version:**

Yann Guiguen, Alexis Fostier, - Aquacop (pyf), Bernard Jalabert. In vitro gonadal steroidogenesis in relation to sex inversion in the protandrous seabass *Lates calcarifer*. 2. International Symposium on Fish Endocrinology, Jun 1992, Saint-Malo, France. 116 p., 1992. <hal-02778209>

HAL Id: hal-02778209

<https://hal.inrae.fr/hal-02778209v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



2nd

**INTERNATIONAL
SYMPOSIUM
on FISH
ENDOCRINOLOGY**

Abstracts

PALAIS DU GRAND LARGE

SAINT-MALO

JUNE 1 - 4 1992



MINISTÈRE
DE LA RECHERCHE
ET DE LA TECHNOLOGIE



VILLE
DE
SAINT-MALO



IN VITRO GONADAL STEROIDOGENESIS IN RELATION TO SEX-INVERSION IN THE PROTANDROUS SEABASS *LATES CALCARIFER*.

Y. GUIGUEN¹, A. FOSTIER¹, AQUACOP², B. JALABERT¹.

¹Laboratory of Fish Physiology, INRA, Campus de Beaulieu 35042 Rennes France.

²Aquaculture Team of the Oceanologic Centre of the Pacific, IFREMER, BP 7004 Taravao Tahiti Polynésie Française.

The tropical seabass *Lates calcarifer* is an important commercial fish for aquaculture and fisheries in the Indo-Pacific area and it has been introduced in French Polynesia for breeding in sea-cages. Control of sex-inversion in this hermaphrodite species is required for the management of the spawners stock.

In Tahiti this protandrous fish underwent sex-inversion after the annual reproductive season. Histological studies of the gonads has been performed during a whole year sexual cycle. The main features of the various stages of the sex-inversion process were: degeneration of the male germinal tissue (early inverting stage), appearance of a peripheral female germinal tissue with persistence of degenerative male cells (mid inverting stage), and centripetal proliferation of the female germinal tissue with no more male cells (late inverting stage). Furthermore, this process required deep morphological changes in the gonads because of a strong dimorphism between testis and ovary.

In parallel to the histological analysis, (1,2,6,7-³H) Androstenedione (And) metabolism (incubation of 1 g of minced tissue, 1 h, t=29±1 °C, without cofactor) by the gonads was studied *in vitro* at various sexual stages including early and late inverting stages. Metabolites were identified by means of thin layer chromatography, high performance liquid chromatography and recrystallization to constant specific activity.

In the male gonad at the spermiation stage (testis full of spermatozoa) incubations gave the following products: Testosterone (4Androsten17βol-3one, T), 5βAndrostan3,17dione (5βAnd), Etiocholanolone (5βAndrostan3αol-17one, Et), 5βAndrostan3βol-17one (5βAnd3βol), 5βAndrostan17βol-3one (5βT), 5βAndrostan3α17βdiol (5βAnd3α17βdiol) and 11βhydroxy-androstenedione (4Androsten11βol-3,17dione, 11βAnd).

In the early inverting stage we identified 11βAnd, Adrenosterone (4Androsten3,11,17trione, Ad), 11ketotestosterone (4Androsten17βol-3,11dione, 11KT) 11βhydroxytestosterone (4Androsten11β,17βdiol-3one, 11βT), T, 5βAnd, 5βT, Et and 5βAnd3βol.

In the late inverting stage only T and 5βAnd have been identified, but the main metabolite (up to 50% of all the metabolites) was an unknown oestrogen like compound.

In the female gonad at the previtellogenesis stage (ovary mainly filled with previtellogenetic oocytes) we found T, 5βAnd, Et, 5βAnd3α17βdiol and oestrone (E1).

To summarize, 17βhydroxysteroid deshydrogenase (17βHSD), 5β reductase (5βR) and 3α hydroxylase (3αH) were found in all incubations, while aromatase was only detected in the female and 3βH and 11βH only in the male and the early inverting stage. In addition to this high 11βH activity in the early inverting stage, we characterized a high 11βHSD activity. These important enzymatic activities have previously been described in testicular regression stage of the protandrous *Amphiprion frenatus* (LATZ *et al.*, 1991), and *Pagellus acarne* (REINBOTH *et al.*, 1986). Furthermore, in gonochoristic fish gonads this 11βH activity has been considered to be a male characteristic, but physiological roles of 11 oxygenated androgens remain unclear especially during protandrous sex-inversion.

Attempts are now being made to characterize the principal unknown metabolite found in the late inverting stage with the view of testing its biological activity on the sex-inversion process.

LATZ, M., STAHLSCHMIDT-ALLNER, P., REINBOTH, R., 1991. In Scott, Sumpter, Kime and Rolfe Eds. Proceedings of the fourth international symposium on the reproductive physiology of fish. Norwich, U.K., 7-12 July 1991. 89-91.

REINBOTH, R., BECKER, B., LATZ, M., 1986. Gen. Comp. Endocrinol. 62, 335-340.