



HAL
open science

Fertilizing ability of frozen-thawed sperm of trout in relation with rearing temperature and dietary lipid origin : what about membrane fluidity and polar lipid composition

Catherine Labbé, Gérard Maise, K. Muller, A Zachowski, Sadasivam Kaushik, Maurice Loir

► To cite this version:

Catherine Labbé, Gérard Maise, K. Muller, A Zachowski, Sadasivam Kaushik, et al.. Fertilizing ability of frozen-thawed sperm of trout in relation with rearing temperature and dietary lipid origin : what about membrane fluidity and polar lipid composition. Workshop on Gamete and Embryo Storage and Cryopreservation in Aquatic Organisms, Mar 1992, Marly-Le-Roy, France. hal-02778512

HAL Id: hal-02778512

<https://hal.inrae.fr/hal-02778512>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

WORKSHOP ON GAMETE AND EMBRYO STORAGE AND CRYOPRESERVATION IN AQUATIC ORGANISMS; INJEP

Institut National de la Jeunesse et de l'Education populaire

Parc du Val Flory, 11 rue Paul Leplat

Tel : 33 (1) 39 68 49 11 : - : Fax : 33 (1) 39 16 57 79

78160 **MARLY le ROY/FRANCE**

(West of PARIS) 30/3 to 2/4 1992, sponsored by the CEC, Directorate for fisheries, IFREMER, Societé IMV, Societé l'AIR LIQUIDE and included in the "Reproductive Biology in Aquaculture" network.

Address for correspondance

R. BILLARD. Ichtyologie Museum d'Histoire Naturelle, 43, rue Cuvier F. 75231 PARIS cédex 05.

Tel : 33 (1) 40.79.37.45 - Fax : 33 (1) 40.79.37.71

PROGRAMME

ABSTRACTS

LIST OF PARTICIPANTS

Organizing Committee :R. BILLARD Laboratoire d'Ichtyologie, 43, rue Cuvier F. 75231 PARIS Cedex 05,
M. LOIR, G. MAISSE INRA Beaulieu F. 35042 RENNES, J. FAUVEL IFREMER BP 70, F. 29263 BREST
J. COSSON, CNRS F/06230 VILLEFRANCHE/MER,

Scientific Committee : M. ABRAHAM (Israel); R. BILLARD, H. BOULEKBACHE, J. COSSON, B. JALABERT, M. LOIR, G. MAISSE, (France); V. HILGE, W. HOLTZ, H. STEIN (Germany);
Y. NAGAHAMA (Japan); C. RICHTER, (NL); A. P. SCOTT, (UK)

**WORKSHOP ON GAMETE AND EMBRYO STORAGE AND
CRYOPRESERVATION IN AQUATIC ORGANISMS ;
MARLY LE ROI/France.
30 March - 2 April 1992**

General organization : R. BILLARD, J. COSSON, C. FAUVEL, M. LOIR, G. MAISSE.

Secretariat assistance and documentation : F. ANDRE, J. BARTHELEMY, J. FAURILLON, F. NADOT (Museum of Natural History), J. C. HUREAU (European Ichthyological Union).

We gratefully acknowledge the financial support we received from :

- The Commission of the European Community (DG 14) (Programme FAR)
200, rue de la Loi Bruxelles BELGIQUE. Tel : 3222351111

- IFREMER (Institut Français pour l'Etude de la Mer) rue J. J. Rousseau
92 Issy les Moulineaux FRANCE. Tel : 46 48 21 00

- Le Ministère de la Recherche et de la Technologie DAI (Bourses ACCES) 1, rue Descartes 75231 PARIS. Tel : 46 34 35 35

- La Société IMV (B. CASSOU) 10, rue Clémenceau 61302 L'AIGLE
Cédex FRANCE. Tel : 33 34 64 64 Fax : 33 34 11 98

- La Société L'AIR LIQUIDE (P. COFFRE et E. DELPOULLE) Parc G.
Eiffel 8, rue Gutenberg Bussy St Georges 77607 Marne La Vallée FRANCE.
Tel : 64 76 15 29

- Sepia International (A. BRUNEL) 14, rue G. Eiffel 78182 St Quentin
en Yvelines FRANCE. Tel : 64 76 15 29

FERTILIZING ABILITY OF FROZEN-THAWED SPERM OF TROUT IN RELATION WITH REARING TEMPERATURE AND DIETARY LIPID ORIGIN : WHAT ABOUT MEMBRANE FLUIDITY AND POLAR LIPID COMPOSITION

C. Labbe⁽¹⁾, G. Maise⁽¹⁾, K. Müller⁽²⁾, A. Zachowski⁽²⁾, S. Kaushik⁽³⁾ and M. Loir⁽¹⁾

(1) Laboratoire de Physiologie des Poissons, INRA, Campus de Beaulieu, 35042 RENNES Cedex (France) ; (2) Laboratoire de Biophysique Cellulaire, Institut de Biologie Physico-Chimique, 13 rue Pierre et Marie Curie, 75005 PARIS (France) ; (3) Laboratoire de Nutrition des Poissons, INRA, BP 3, 64310 SAINT PEE sur NIVELLE (France)

A factorial experiment was designed to investigate the influence of low rearing temperature (8 °C) and high rearing temperature (18 °C) with a linoleic fatty acid rich diet (corn oil) and a n-3 fatty acid rich diet (cod liver oil) on fertilizing ability of frozen-thawed trout sperm. The biochemical and biophysical modifications of the sperm plasma membrane induced by these rearing conditions were assessed. Juvenile trout (one year old and 100 g mean weight) were submitted to one of the temperatures and one of the diets. The experiment was two years during which they went twice through gonadal development and reached twice sexual maturity. Milt was collected throughout the first spawning season. A part was frozen (Maise *et al.* 1988, *Aquat. Living Res.* 1 45-51). Fatty acid analyses inside each phospholipid classes of the plasma membrane was performed on the remaining semen sample. Sperm from the second spawning season was collected once for biophysical analyses : membrane fluidity was recorded by electron spin resonance (ESR) using spin-labeled phospholipids with a short (C5) b-chain bearing a nitroxide radical (Seigneuret et Devaux 1984, *Proc. Natl. Acad. Sci. USA* 81 3751-3755).

Semen from the 18 °C adapted trout displayed a two fold greater fertilizing ability after freeze-thawing than the 8 °C ones although motility was the same before cryopreservation. The diet had no influence on sperm quality and the fertilization rates were not significantly modified by either linoleate or n-3 fatty acid enrichments. Fatty acid analysis of each phospholipid revealed that PC had the lower UFA/SFA ratio (= 1,4) with 36 % of 16:0 and PS the higher (= 4) with 26 % of 18:1 n-9 and 26 % of 22:6 n-3. The fatty acid composition of PE, PI (phosphatidyl inositol), PS and PC reflected well the features of the experimental diets mainly with 20:4 n-6 being two fold higher in corn oil-milt and 20:5 n-3 being two fold higher in fish oil-milt. The rearing temperature of the broodstock influenced mainly the 18:0 levels, higher in 18 °C-milt than in 8 °C-milt. In PE, 16:0 and 20:5 n-3 were higher in 8 °C-milt than in 18 °C-ones but in minor proportions. Neither diet nor rearing temperature of the broodstock seemed to affect the fluidity of the sperm plasma membrane as assessed with labeled phosphatidyl choline (PC), phosphatidyl ethanolamine (PE), phosphatidyl serine (PS) or sphingomyeline (SM). However, fluidity assessed with labeled cholesterol seemed slightly lower when mesured at 8 °C on 8 °C-milt than when mesured at 18 °C on 18 °C-milt. We estimated that 90 % of the membrane PS and 85 % of the membrane PE were localized on the inner leaflet of the membrane while 85 % of PC and 100 % of SM were on the outer leaflet. Labeled PC and SM are in a more packed environment than labeled PS leading to the observation that the outer membrane leaflet is more rigid than the inner one. These characteristics are not influenced by diet or temperature.

In conclusion, we pointed out that 18 °C-milt tolerance to cryopreservation is two fold better than 8 °C-milt although neither fluidity nor phospholipid fatty acid compositions displayed major differences between the two temperatures. When measured by ESR, no "homeoviscous adaptation" appeared between the two broodstock sperm. We propose that, at the opposite of metabolically active cells, the sperm cell, which has a very low metabolic activity, does not restructure its plasma membrane. Such a restructuring is energy-consuming and would be without adaptative advantage for the gamete. The biological factors involved in the improved freeze-thaw tolerance we observed for 18 °C milt are then not yet elucidated. We also described sperm plasma membrane with the outer leaflet more fluid, enriched in PE and PS, in MUFA and PUFA while the outer leaflet is more rigid, enriched in PC and in SFA. The high differences induced by diet on fatty acid composition did not affect fluidity or sperm freeze-thaw resistance.