



HAL
open science

Regulation of GtH secretion by GnRH and steroid hormones in male and female rainbow trout. An in vitro study

Claudine Weil, Odile Marcuzzi

► To cite this version:

Claudine Weil, Odile Marcuzzi. Regulation of GtH secretion by GnRH and steroid hormones in male and female rainbow trout. An in vitro study. 3. International Symposium on the Reproductive Physiology of Fish, Marine Sciences Research Laboratory., Aug 1987, St. John's, Newfoundland, Canada. hal-02781342

HAL Id: hal-02781342

<https://hal.inrae.fr/hal-02781342>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

REGULATION OF GtH SECRETION BY GnRH AND STEROID HORMONES IN MALE AND FEMALE RAINBOW TROUT.
AN IN VITRO STUDY

C. WEIL, O. MARCUZZI

Laboratoire de Physiologie des Poissons, INRA, Rennes, France

Male and female rainbow trout exhibit an annual sexual cycle characterized by variations of pituitary gonadotropin hormone (GtH) levels, of plasma GtH and steroid hormones levels and of pituitary responsiveness to gonadotropin-releasing hormone (GnRH). For this latter parameter maximal values were recorded at the time of maturation - ovulation or prespermiation - spermiation (Weil, 1981).

In the present work, we test *in vitro* the change in gonadotroph sensitivity to GnRH and its modulation by steroid hormones during the reproductive cycle.

Primary cultures of whole pituitary maintained in standardized conditions (Weil et al., 1986) were used. At definite stages of gametogenesis, cells dispersed with collagenase were preincubated for 3 days in control medium or in medium containing the main steroids involved in oocyte maturation 17 α hydroxy 20 β dihydroprogesterone (17,20-P) and spermiation 11-Ketotestosterone (11K-T) and 17,20-P. Cultures were then incubated with sGnRH during 24 hrs after which GtH released in the medium was measured.

Pituitary responsiveness to sGnRH was studied in female at the beginning of vitellogenesis (BV), at the SPGV stage - corresponding to oocytes with subperipheral germinal vesicle and the day of ovulation. Two doses of 17,20-P were tested : the first one corresponding to circulating levels at maturation (400 ng/ml), the second one to levels just prior maturation (20 ng/ml) when the germinal vesicle is in a peripheral position (PGV stage). In control cultures, pituitary responsiveness to GnRH is maximal at the time of ovulation. This might partly be due to high *in vivo* circulating levels of 17,20-P since a pretreatment with this steroid (maturation dose) increases the GnRH-induced GtH release of pituitary cultures from females at the BV stage. At the SPGV stage, only the PGV dose induces an increase likely to explain SPGV to PGV stages rise in plasma GtH levels (Weil, 1981). At the time of ovulation, both doses of 17,20-P induce a decrease in pituitary sensitivity to GnRH. This might account for the decline in plasma GtH levels following the injection of 17,20-P (Jalabert et al., 1976).

Males were studied at the beginning of spermatogenesis, at spermiation and prespermiation. 11K-T and 17,20-P were used at do-

ses corresponding to spermiation circulating levels, 50 and 20 ng/ml respectively. In control cultures, pituitary sensitivity to GnRH increase from beginning of spermatogenesis to spermiation. This could be related to the *in vivo* rising levels of 17,20-P and 11K-T since a pretreatment with these hormones increases the GnRH-induced GtH release of beginning of spermatogenesis cultures. At spermiation, no effect of steroid pretreatment is recorded whereas during prespermiation a slight decrease is linked with the presence of 17,20-P. This latter observation might explain the slight decline in plasma GtH observed at onset of spermiation (Sanchez-Rodriguez et al., 1978).


In conclusion, we demonstrate that the variation of circulating GtH levels may partly be due to a direct action of gonadal steroid hormones on pituitary gonadotrophs by modulating their responsiveness to GnRH.

References

- Jalabert, B., Bry, C., Breton, B. and Campbell, C. 1976. Action de la 17 α hydroxy-20 β dihydroprogesterone et de la progesté-
rone sur la maturation et l'ovulation *in vivo* et sur le niveau d'hormone gonadotrope plasmatique t-GtH. C.R. Acad. Sci. Paris. 283 : 1205-1208.
- Sanchez-Rodriguez, M., Escaffre, A.M., Marlot, S. and Reinaud, P. 1978. The spermiation period in the rainbow trout (*Salmo gairdneri*). Plasma gonadotropin and androgen levels, sperm production and biochemical changes in the seminal fluid. Ann. Biol. Anim. Biochim. Biophys. 18, 943-948.
- Weil, C. 1981. La fonction gonadotrope de l'hypophyse au cours du cycle sexuel chez deux poissons téléostéens la Carpe commune (*Cyprinus carpio*) et la truite arc-en-ciel (*Salmo gairdneri*) ; son contrôle par l'hypothalamus, les gonades et les facteurs externes. Thèse de Doctorat d'Etat, Paris VI.
- Weil, C., Hansen, P., Hyam, D., Le Gac, F., Breton, B. and Crim, L.W. 1986. Use of pituitary cells in primary culture to study the regulation of gonadotropin hormone (GtH) secretion in rainbow trout : setting up and validating the system as assessed by its responsiveness to mammalian and salmon gonadotropin releasing hormone. Gen. Comp. Endocr. 62, 202-209.

***Proceedings of the Third International Symposium
on the Reproductive Physiology of Fish***

St. John's, Newfoundland, Canada, 2-7 August 1987



Marine Sciences Research Laboratory

Memorial University of Newfoundland

Logy Bay, Newfoundland