Tentative estimation of chloroplast and non chloroplast proteins of the leaves of a plant biomass

Roger Douillard

► To cite this version:

Roger Douillard. Tentative estimation of chloroplast and non chloroplast proteins of the leaves of a plant biomass. 6. International congress on photosynthesis, Aug 1983, Bruxelles, Belgium. hal-02781574

HAL Id: hal-02781574

<https://hal.inrae.fr/hal-02781574>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

OFFPRINT FROM

Advances in Photosynthesis Research

Proceedings of the VIth International Congress on Photosynthesis, August 1-6,
1983, Brussels, Belgium

Edited by C. SYBESMA

ADVANCES IN AGRICULTURAL BIOTECHNOLOGY

The majority of the invited contributions are review papers, summarizing and putting into perspective the latest results of research in a particular area; the other contributions comprise research papers giving details on specific aspects. "Advances in Photosynthesis Research", therefore, is an important document, containing the latest high-level information about photosynthesis in its broadest sense. The four volumes are of considerable value as a reference source for scientists active in the field and others wishing to be informed about recent developments. Furthermore, these volumes will prove to be important as a means of background study for those entering this exciting and promising field of investigation.

1984. 3480 pp. in 4 volumes

ISBN 90-247-2946-7

Dfl. 1.250,00/US\$ 478.00/£ 317.50

Inventaire

N°

PUB 133

MARTINUS NIJHOFF/DR W. JUNK PUBLISHERS
A Member of the Kluwer Academic Publishers Group
P.O. Box 566, 2501 CN The Hague, The Netherlands

NEW AND FORTHCOMING PUBLICATIONS

Photosynthesis Research

an international journal

ISSN 0166-8595

Editor-in-Chief

DR. MARCELLE

*Laboratory of Plant Physiology
Research Station of Gorsem
Brede Akker 3
B-3800 Sint Truiden, Belgium
Tel.: 011-682019*

Associate Editor

DR. GOVINDJEE

*Department of Plant Biology
UIUC
289 Morrill Hall
505 South Goodwin Avenue
Urbana, Illinois 61801 (USA)
Tel.: (217) 333-1794*

The consulting Editorial Board consists of approximately 30 members

Aims and Scope

Photosynthesis Research aims to provide a forum for original papers dealing with photosynthesis without imposing an artificial division between fundamental and applied research. Our aim is to cover *all aspects of photosynthesis research* starting from the primary reactions and going as far as the energy from biomass. Topics acceptable for publication could include primary reactions, electron transport, photophosphorylation, carbon assimilation, regulatory phenomena, environmental and ecological aspects of photosynthesis, photorespiration, CO₂ fixation in CAM plants, and bacterial and algal photosynthesis, although one should not feel restricted just to these topics.

In the case of artificial systems, the authors should be able to judge for themselves whether to publish in a photosynthesis or photobiology journal. In the case of doubt, the Editorial Board can help the author in making this decision.

Papers will be considered for publication at all levels of plant organization: sub-cellular, cellular, whole plant and canopy levels.

Manuscripts submitted for publication are always reviewed by two referees, not necessarily members of the Editorial Board, and chosen for their competence in the field covered by the paper. The appeal to people who are not members of the Editorial Board is necessary in order to cover the whole field of photosynthesis.

Call for papers

Authors who wish to contribute to Photosynthesis Research are invited to send their manuscripts or requests for "Instructions to Authors". Manuscripts from North America should be sent to Dr. Govindjee, from all other areas to Dr. Marcelle.

Subscription information

Vol. 5, 1984 (4 issues): For institutions, Dfl. 165.00 (US\$ 66.00), plus Dfl. 25.00 (US\$ 10.00) for postage and handling. For individuals, Dfl. 100.00 (US\$ 40.00) including postage and handling.

Available at the end of 1984

Photosynthesis During Leaf Development

Edited by
Z. SESTÁK

This unique monograph was written by nine scientists from the Department of Physiology of Photosynthesis and Water Relations of the Institute of Experimental Botany, Czechoslovak Academy of Sciences in Prague. They based it on a choice of papers from the vast literature, which was not always easy to assess, and on the 20 years pioneering research experiences of their laboratories.

MARTINUS NIJHOFF/DR W. JUNK PUBLISHERS

P.O. Box 566, 2501 CN The Hague, The Netherlands/190 Old Derby Street, Hingham MA 02043, U.S.A.

TENTATIVE ESTIMATION OF CHLOROPLAST AND NON CHLOROPLAST PROTEINS OF THE LEAVES OF A PLANT BIOMASS

R. DOUILLARD

1. INTRODUCTION

The membranous proteins of lamellae and the soluble proteins of chloroplast stroma account for nearly all the proteins in leaves (Joyard, 1979). Agriculture productivity relies on these proteins in different ways. First, the biosynthesis of sugars from carbon dioxide, water and light, depends on them since they are building blocks of the photosynthetic apparatus. Second, they are part of several crops such as vegetables, fodder or leaf protein concentrate; the quality and value of these crops depend obviously on leaf proteins. It is thus clear that leaf protein composition is an essential data in plant physiology and ecology or in plant breeding and agriculture. In this communication a model is presented which takes into account RuBPcase turnover number and content in leaves of C₃ and C₄ type plants and relates leaf protein weight to net and gross primary production of a canopy.

2. MODEL DESCRIPTION

The principle of the derivation relating the number N of RuBPcase molecules of a canopy to the carbon weight it may assimilate during a year (CW) has been given by Ellis (1979):

$$CW = N \cdot \tau \cdot TN \cdot 12 \quad (1) \quad (\text{Abbreviations in table I})$$

12 is the atomic weight of carbon.

TABLE I. Abbreviations and numerical values

CW :	annual gross fixation of carbon by a plant biomass ; for land plants of the Earth, CW is estimated to equal two times the net fixation (Box, 1977), that is $2 \times 53 \cdot 10^{15}$ g (Woodwell, 1978).
LP :	leaf protein weight of a plant biomass.
MRuBPcase :	molecular weight of RuBPcase, estimated at $5.5 \cdot 10^5$ daltons (Yeoh et al., 1982).
N :	number of RuBPcase molecules of a plant biomass.
RuBPcase :	ribulose 1,5-bisphosphate carboxylase/oxygenase.
TN :	turnover number of RuBPcase ($\text{kat} \cdot \text{mole}^{-1}$) ; see table III.
ρ :	ratio of leaf protein weight to RuBPcase weight in a plant biomass ; see table II.
τ :	duration of enzyme activity in one year.
i :	index defining an homogeneous type of plant in a specified climatic area.
3 and 4 :	indexes corresponding to C ₃ and C ₄ plants.

For the whole world N may thus be calculated as follows :

$$N = \frac{1}{12} \sum_{i=1}^I \frac{CW_i}{\tau_i \cdot TN_i} \quad (2)$$

Sybesma, C. (ed.), *Advances in Photosynthesis Research*, Vol. IV. ISBN 90-247-2945-9.

© 1984 Martinus Nijhoff/Dr W. Junk Publishers, The Hague/Boston/Lancaster.

Printed in The Netherlands.

IV.2.166

Each i defines an homogeneous type of plant in a specified climatic area. Leaf protein weight is thus given by :

$$LP = \frac{MRuBPcase}{12} \sum_i \frac{CW_i}{\tau_i \cdot TN_i} \rho_i \quad (3)$$

where it is assumed, according to Yeoh et al. (1982) that RuBPcase molecular weight is rather constant among various plant species. Formula (3) raises nevertheless the question of the choice of plant types and of climatic areas. An example is shown below.

TABLE II. Protein (LP) and RuBPcase content of leaves; DM : dry matter of leaves ; ρ = LP/RuBPcase. (a) : Singer et al. (1952) ; (b) : Pheloung, Brady (1979).

Plant	LP (% DM)	ρ	RuBPcase (% LP)	ref.
C₃ Plants				
Cucumis anguria	28	7	14	a
Lycopersicum esculentum	31	10	10	a
Pisum sativum	25	9	11	a
Xanthium pennsylvanicum	40	11	9.2	a
Triticum aestivum L.	48	7	14	b
Phalaris tuberosa L.	42	6.3	16	b
Bromus uniloides H.B.K.	33	8	12	b
C₄ Plants				
Cynodon dactylon L. Pers.	30	22	4.5	b
Chloris gayana Kunth	26	36	2.8	b
Paspalum dilatatum Poir.	27	23	4.3	b
Pennisetum clandestinum Hochst.	26	29	3.4	b

3. LEAF PROTEIN WEIGHT OF THE EARTH LAND PLANTS

In this example, a single climatic area is considered but a distinction is made between C₃ and C₄ plants. Thus :

$$LP = \frac{MRuBPcase}{12 \cdot \tau} \left(\frac{CW_3}{TN_3} \cdot \rho_3 + \frac{CW_4}{TN_4} \cdot \rho_4 \right) \quad (4)$$

As shown in Tab II, ρ_3 ranges from 6 to 11 and ρ_4 from 22 to 36. ρ_4 is thus about 3 times higher than ρ_3 . On the other hand, the turnover number (TN) is limited in vivo by the non saturating carbon dioxide amount (Ellis, 1979 ; Wildner, 1981 ; Tab. III). In fact, this amount is much higher in the bundle sheet cells of C₄ type plants, where RuBPcase works, than in C₃ type plant cells (Ku, 1979). TN₄ is thus clearly higher than TN₃. For these reasons, it is admissible that the ratios ρ_3/TN_3 and ρ_4/TN_4 have similar values. Since $CW_3 + CW_4 = CW$, the relation (4) may be written as follows :

$$LP = \frac{MRuBPcase}{12 \cdot \tau} CW \frac{\rho_3}{TN_3} \quad (5)$$

CW, the gross primary fixation, equals at earth scale, about two times the net primary fixation of carbon dioxide by land plants (Box, 1977). The difference

between gross and net productivity is accounted for by dark respiration. Dark respiration estimated in various conditions and for numerous plants ranges indeed from 20 to 70 % of gross primary photosynthetic productivity (Kira, 1975; Yamaguchi, 1978 ; Ruget, 1981 a and b). Photorespiration is neglected in this estimation.

τ , duration of enzyme activity during a year greatly depends on the plant and on the climatic area. It is estimated to be 0.3 year or $9.5 \cdot 10^6$ sec.

The amount of leaf protein in the land plants of the Earth is thus (data of Tab. I, II and III) :

$$LP = \frac{5.5 \cdot 10^5}{12 \cdot 9.5 \cdot 10^6} \cdot 2.53 \cdot 10^{15} \cdot \frac{6 \text{ to } 11}{1 \text{ to } 3.3} \text{ g}$$

$$LP = 9.3 \text{ to } 56 \cdot 10^{14} \text{ g}$$

TABLE III. Turnover number (TN) of RuBPCase ($\text{kat} \cdot \text{mole}^{-1}$), assuming a molecular weight of 550 000 daltons.

Measurement conditions	Plant	TN	Reference
In vitro		18 - 37	Ellis, 1977
"	Spinach	14 - 16	Hall et al., 1981
"	"	17 - 32	Berhow et al., 1982
"	"	8 - 9	Johal, Bourque, 1979
"	Barley	15 - 32	Berhow et al., 1982
Saturating substrate concentrations		17	Ellis, 1979
Natural conditions		3.3	Ellis, 1979
		1 to 3.3	Wildner, 1981

4. CONCLUDING REMARKS

The present model of plant productivity allows the connection of data on the amount and turnover number of RuBPCase of photosynthetic apparatus and data on carbon dioxide assimilation by a canopy. Such a model should allow a better understanding of the mechanisms and of the limitations of carbon dioxide assimilation by a canopy. It could also be used as a model for leaf protein productivity.

World production of cereals was $17 \cdot 10^{14}$ g in 1981 (FAO, 1982). Their protein content is about 8% (Demarquilly et al., 1978). World cereal protein production was thus $14 \cdot 10^{13}$ g in 1981. The amount of leaf protein on the Earth is thus about ten times higher than the yearly cereal protein production.

The estimate of leaf protein in a canopy allows the evaluation of other leaf components and specially those of chloroplasts as far as their ratio to leaf proteins is known. Thus the world amount of RuBPCase may be evaluated assuming either a C_3 type or a C_4 type carbon dioxide assimilation. According to the above estimation of leaf proteins and to the data of table II, the estimate ranges from 8.6 to $90 \cdot 10^{13}$ g for a C_3 type or from 2.6 to $25 \cdot 10^{13}$ g for a C_4 type. These values are higher than the $4 \cdot 10^{13}$ g computed by Ellis (1979) who

IV.2.168

used net instead of gross primary productivity and estimated τ at 0.5 year (Personal communication).

REFERENCES

- Berhow MA, Saluja A and Mc Fadden BA (1982) Rapid purification of D-Ribulose-1,5-bisphosphate carboxylase by vertical sedimentation in a reoriented gradient, *Plant Sci. Lett.* 27, 51-57.
- Box E and Lieth H (1977) Geographical dimensions of terrestrial net and gross primary productivity : a first attempt. European seminar on biological solar energy conversion systems. Grenoble, France.
- Demarquilly C, Andrieu J and Sauvant D (1978) Tableaux de la valeur nutritive des aliments. In INRA ed. Alimentation des ruminants, pp 519-555. Versailles, France.
- Ellis RJ (1979) The most abundant protein in the world, *Trends biochem. Sci.* 4, 241-244.
- FAO (1982) Production yearbook 35, FAO p 85. Rome, Italy.
- Hall NP, Mc Curry SD and Tolbert NE (1981) Storage and maintaining activity of ribulose bisphosphate carboxylase/oxygenase, *Plant Physiol.* 67,1220-1223.
- Johal S and Bourque DP (1979) Crystalline ribulose-1,5-bisphosphate carboxylase/oxygenase from spinach, *Science* 204, 75-76.
- Joyard J (1979) L'enveloppe des chloroplastes. Thèse de doctorat es sciences naturelles. Université scientifique et médicale de Grenoble.
- Kira T (1975) Primary production of forests. In Cooper JP, ed. Photosynthesis and productivity in different environments, pp 5-40. London, England : Cambridge University Press.
- Ku MSB, Schmitt MR and Edwards GE (1979) Quantitative determination of RuBP Carboxylase-Oxygenase protein in leaves of several C₃ and C₄ plants, *J. exp Bot.* 30, 89-98.
- Pheloung P and Brady CJ (1979) Soluble and fraction 1 protein in leaves of C₃ and C₄ grasses, *J. Sci. Food Agric.* 30, 246-250.
- Ruget F (1981 a) Respiration de croissance et respiration d'entretien : méthodes de mesure, comparaison des résultats, *Agronomie* 1, 601-610.
- Ruget F (1981 b) Respiration de croissance et d'entretien du Maïs dans différentes conditions artificielles de culture, *Agronomie* 1, 845-852.
- Singer SJ, Eggman L, Campbell JM and Wildmänn SG (1952) The proteins of green leaves IV a high molecular weight protein comprising a large part of the cytoplasmic proteins, *J. Biol. Chem.* 97, 233-239.
- Wildner GF (1981) Ribulose-1,5-bisphosphate carboxylase-oxygenase aspects and prospects, *Physiol. Plant.* 52, 385-389.
- Woodwell GM (1978) The carbon dioxide question, *Sci. Am.* 238, 34-43.
- Yamaguchi J (1978) Respiration and the growth efficiency in relation to crop productivity, *J. Fac. Agric. Hokkaido Univ.* 59, 59-129.
- Yeoh HH, Stone NE and Watson L (1982) Taxonomic variation in the subunit amino acid composition of RuBPCarboxylases from grasses, *Phytochem.* 21,71-80.

ACKNOWLEDGEMENTS

The author wishes to express his gratitude to C. Varlet-Grancher for invaluable help in the preparation of this communication.

Author address : R. Douillard, Laboratoire de Biochimie et Technologie des Protéines, INRA, La Géraudière, 44072 Nantes Cédex, France.