



**HAL**  
open science

## Characterization and GtH regulation of microsomal ovarian aromatase activity in rainbow trout

A. de Mones, Alexis Fostier

► **To cite this version:**

A. de Mones, Alexis Fostier. Characterization and GtH regulation of microsomal ovarian aromatase activity in rainbow trout. 3. International Symposium on the Reproductive Physiology of Fish, Marine Sciences Research Laboratory., Aug 1987, St. John's, Newfoundland, Canada. hal-02783867

**HAL Id: hal-02783867**

**<https://hal.inrae.fr/hal-02783867>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A. de MONES and A. FOSTIER

Laboratoire de Physiologie des Poissons - I.N.R.A. - Rennes, France

### Introduction

The maturational gonadotropin (GtH) has been claimed to depress aromatase activity in fish ovary, especially at the end of the sexual cycle. These conclusions were based on *in vivo* analysis of plasma oestradiol (E2) levels, *in vitro* measurements of E2 output (Fostier & Jalabert, 1986) and *in vitro* studies of androstenedione metabolism by ovarian tissues (Sire & Depeche, 1981).

To confirm GtH action, we partially isolated aromatase activity by subcellular fractionation for further characterization and kinetic studies.

### Experimental

Microsomes were obtained by differential centrifugation of homogenates from preovulatory or vitellogenic (oocyte mean diameter: 2.5 mm) ovaries. They were incubated 10 min. with NADPH and 3H androstenedione ( $\Delta 4$ ) or testosterone (T). Oestradiol and oestrone production was measured after extraction and chromatography. Oestrogens identity was checked in two cases by crystallization to constant isotopic ratio.

To study aromatase activity regulation, ovaries were preincubated with or without GtH (60 ng/ml) for 24 h before their processing for subcellular fractionation. Each set of experiment was performed with the pooled ovaries of one (preovulatory) or two (vitellogenic) animals. Kinetic constants were estimated from experimental data by a non-parametric fit to the Michaelis-Menten equation (Eisenthal & Cornish-Bowden, 1974) and compared using a Mann-Whitney test.

### Results

The aromatase specific activity was compared in the various fractions collected during subcellular fractionation. It was mainly located in the microsomal pellet (195000g pellet: 92 fmoles oestrogens/min. mg; total homogenate: 0.2 fmoles/min.mg). Aromatization was therefore studied in the microsomes.

The activity was linear with time during at least 10 minutes. Specific activity was not affected by microsomal protein concentration between 1 and 5 mg. Androstatrienedione, known as an aromatase inhibitor in mammals, was also a powerful inhibitor of oestrogens synthesis in our preparations (I 50: 90 nM, for a 200 nM  $\Delta 4$  concentration).

Kinetics of aromatization was investi-

gated in microsomes issued from GtH treated or control ovaries. In most cases, the Michaelis constant (Km) was not affected while Vmax was lowered (table) by GtH treatment.

Table: Effect of GtH pretreatment (60ng/ml) of trout ovaries on their microsomal aromatase activity.

\*: significantly different from the control

Sexual cycle period	substrate	Vmax (pmoles/min-animal)	
		control	+GtH
preovulation		8	8
		20	7 *
	$\Delta 4$	11	7 *
		4	3 *
vitellogenesis		46	47
		39	25 *
	T	46	31 *

### Conclusion

As in mammals, trout ovarian aromatase appears to be mostly located in microsomes. Its partial isolation provides a new tool to study the regulation of its activity.


We confirm GtH inhibitory effect on ovarian aromatase activity during the second half of the sexual cycle.

### References

- Eisenthal, R. & A. Cornish-Bowden, 1974. The direct linear plot. A new graphical procedure for estimating enzyme kinetic parameters. *Biochem. J.*, 139: 715-720
- Fostier, A. & B. Jalabert, 1986. Steroidogenesis in rainbow trout (*Salmo gairdneri*) at various preovulatory stages: changes in plasma hormone levels and *in vivo* and *in vitro* responses of the ovary to salmon gonadotropin. *Fish Physiol. Biochem.*, 2: 87-99
- Sire, O. & J. Depeche, 1981. *In vitro* effect of a fish gonadotropin on aromatase and 17 $\beta$ -hydroxysteroid dehydrogenase activities in the ovary of the rainbow trout (*Salmo gairdneri* R.). *Reprod. Nutr. Develop.*, 21: 715-726

***Proceedings of the Third International Symposium  
on the Reproductive Physiology of Fish***

***St. John's, Newfoundland, Canada, 2-7 August 1987***



***Marine Sciences Research Laboratory***

***Memorial University of Newfoundland***

***Logy Bay, Newfoundland***