

A hydraulic perspective on the interspecific diversity of drought resistance strategies in coexisting tropical rainforest tree species of French Guiana

Camille Ziegler, Sabrina Coste, Sylvain Delzon, Clement Stahl, Damien Bonal

▶ To cite this version:

Camille Ziegler, Sabrina Coste, Sylvain Delzon, Clement Stahl, Damien Bonal. A hydraulic perspective on the interspecific diversity of drought resistance strategies in coexisting tropical rainforest tree species of French Guiana. 3. Xylem International Meeting, 2017, bordeaux, France. hal-02784974

HAL Id: hal-02784974 https://hal.inrae.fr/hal-02784974

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A hydraulic perspective on the interspecific diversity of drought resistance strategies in coexisting tropical rainforest tree species of French Guiana

Camille Ziegler^{1,3*}, Sabrina Coste^{2,3}, Sylvain Delzon⁴, Clément Stahl³, and Damien Bonal¹

Abstract:

- The impressive tree species diversity of the Amazon rainforest has shown an associated wide-range of trait responses related to water use efficiency but little is known about embolism resistance traits in tropical tree species.
- Predicting tree species' responses to an increase in frequency and severity of drought events is therefore challenging. Experimental data is needed in order to deepen our understanding on the ecophysiological mechanisms involved in tolerance and avoidance strategies of tree species in this ecosystem.
- In this ongoing study, we characterize leaf and stem hydraulic traits, i.e. predawn (Ψ_{pd}) and minimum (Ψ_{min}) leaf water potential and leaf water potential at turgor loss point (π_{tlp}) as well as embolism resistance traits for a broad range of co-occurring canopy tree species.
- Preliminary results point out a broad range of interspecific variability in branch vulnerability to embolism. Further work will include predawn and minimum water potential measurements during the dry season to determine the magnitude of experienced stress and calculate hydraulic security margins. This will fuel discussions on the ecological relevance of these traits in predicting a species' drought-resistance.

¹Université de Lorraine, INRA, UMR Écologie et Écophysiologie Forestière, 54280 Champenoux, France

²Université de Guyane, UMR Écologie des Forêts de Guyane, 97379, Kourou, French Guiana

³INRA, CNRS, UMR Écologie des Forêts de Guyane, 97379 Kourou, French Guiana

⁴INRA, University of Bordeaux, UMR BIOGECO, 33615 Pessac, France

^{*}Author for correspondence: Camille.Ziegler@ecofog.gf