



**HAL**  
open science

## Tree, Sex and Size: Ecological determinants of male versus female fecundity in three *Fagus sylvatica* stands

Sylvie Muratorio, Julie Gauzere, Aurore Bontemps, Jean-François Rey,  
Etienne Klein

### ► To cite this version:

Sylvie Muratorio, Julie Gauzere, Aurore Bontemps, Jean-François Rey, Etienne Klein. Tree, Sex and Size: Ecological determinants of male versus female fecundity in three *Fagus sylvatica* stands. 2. Joint Congress on Evolutionary Biology (EVOLUTION 2018), Aug 2018, Montpellier, France. 2018. hal-02785554

**HAL Id: hal-02785554**

**<https://hal.inrae.fr/hal-02785554v1>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Tree, Sex and Size: Ecological determinants of male versus female fecundity in three *Fagus sylvatica* stands

Sylvie Oddou-Muratorio\*, Julie Gauzere\*, Aurore Bontemps\*, Jean-François Rey # & Etienne K. Klein#


\* INRA, URFM and # INRA, BioSp 84000 Avignon, France



Inter-individual variations in female (♀) and male (♂) fecundities have major consequences on population evolutionary potential, through genetic drift and selection. In plants, basic fecundities are classically estimated before dispersal through the resource allocated to ♀ (i.e. the biomass/number of ovules, seeds, ovuliferous flowers or fruits) and ♂ (i.e. the biomass/number of pollen grains or staminate flowers) functions. Alternatively, paternity and parentage reconstruction can provide fitness-based estimates of effective fecundity and reproductive success, though the numbers of offspring assigned to a given individual. However, we miss a conceptual framework linking resource-based and fitness-based estimates of fecundity to reproductive success, and tools to obtain comparable estimates of ♀ and ♂ effective fecundities. The present study aims to fill this gap, first by **providing a new method to estimate together ♀ and ♂ effective fecundities accounting for a spatial configuration effect** (implemented in the MEMM program). Then we apply this method to **study the variation in ♀/♂ fecundities and gender in relation to plant size and competition in a major monoecious tree species, the European beech.**


## Context and objectives

### Bridging the gap between resource-based and fitness-based estimates of fecundity and gender in plants



- **Basic fecundity and phenotypic gender** encompass only the production of pollen and seed and are estimated before dispersal by the number/biomass of seed/pollen (**resource-based estimates**).
- **Reproductive success** is driven by basic fecundity together with additional factors, including spatial processes and individual effects on propagule success.
- **Reproductive success and functional gender** are **observed after dispersal** using **parentage/paternity analyses**, which are potentially biased by spatial and sampling effects.
- **Spatially Explicit Mating Models (SEMM)** get rid of these confounding effects to retrieve the **effective fecundity and gender**, which accounts for individual effects on propagule success.

### Estimating ♀ and ♂ individual fecundities with MEMM




**IMPLEMENTATION** : All parameters and latent variables were estimated in a Bayesian framework using the **MEMM program** with 2 options :

DATA: Offspring = established seedlings → MEMM Option 2 → ESTIMATED Parameters:  $m_p, s, \delta_p, b_p, \sigma^2_\sigma + \{F\sigma_k\}$  and Latent variables:  $\{F\sigma_k\}$

DATA: Offspring = Seeds → MEMM Option 1 → ESTIMATED Parameters:  $m_p, s, \delta_p, b_p, \sigma^2_\sigma + \{F\sigma_k\}$  and Latent variables:  $\{F\sigma_k\}$

The MEMM program (Mixed Effect Mating Model) can be downloaded at : <https://gitlab.paca.inra.fr/frey/MEMMseedlings.git>

### Sampling three *F. sylvatica* plots along an altitudinal gradient




**Previous result** : Pollen immigration increased with altitude: protogyny combined with temperature variation may result in oriented pollen flow from bottom to top (Gauzere et al. Mol Ecol. 2013)

### Result: ecological drivers of fecundity and gender

	Ecological driver	♀ fecundity	♂ fecundity	Femaleness
Individual size	Maximum stem diameter	↗		
	Sum of stems' diameter		↗	
	Tree stature (dominated, dominant)		↗	
Competition	Conspecific competition	↘	↘	↘
	Total competition	↘		↘

- The effects of size and competition were consistent across the three plots.
- Both ♀ and ♂ effective fecundities were resource-limited in the studied species.
- Size effects were sex-specific and consistent with the higher unit cost of ♀ as compared to ♂ reproduction, and with the limitation of ♂ fecundity by mating opportunities.
- Effective gender was independent of size, but femaleness decreased under limited resources (i.e. with increasing competition and density), consistent with the higher unit cost of ♀ reproduction

### Result: ♀ and ♂ effective fecundities and femaleness



- Both ♂ and ♀ fecundities distribution were L-shaped (Y-scale is relative, and vary among plots)
- The variance in ♀ fecundities ( $\sigma^2_\sigma$ ) tended to be higher than the variance in ♂ fecundity ( $\sigma^2_\sigma$ )
- The variance in fecundities tended to be higher at high elevation (significant for  $\sigma^2_\sigma$ )
- The three plots were male-biased, without significant correlation between ♂ and ♀ fecundity

### Conclusions

- SEMM- and resources-based estimates of fecundity are complementary: the latter estimate the resources allocated to reproduction while the former can only estimate a relative amount of pollen or seed produced. Moreover, SEMM-estimates account for individual effects that act independently on location to modify the success of mating.
- In the monoecious tree *Fagus sylvatica*, both ♂ and ♀ fecundities increased with tree size and decreased with competition in the neighborhood (with sex-specific strategies to deal with limited resources).
- This study suggests that the evolutionary potential of *Fagus sylvatica* population is enhanced by low variance in fecundity at low elevation, and high pollen flow at high elevation.

