



HAL
open science

Guinea fowl domestication : new insights from the first whole genome assembly and the pool sequencing of wild and domestic populations

Simon Boitard, Alain Vignal, Noémie Thébault, Michèle Tixier-Boichard, Xavier X Rognon, Guiguigbaza Kossigan-Dayo, Chia Valentine Yapi-Gnaore, Issaka Youssao Abdou Karim, Wes Warren

► **To cite this version:**

Simon Boitard, Alain Vignal, Noémie Thébault, Michèle Tixier-Boichard, Xavier X Rognon, et al.. Guinea fowl domestication : new insights from the first whole genome assembly and the pool sequencing of wild and domestic populations. II Joint Congress on Evolutionary Biology, Aug 2018, Montpellier, France. hal-02785577

HAL Id: hal-02785577

<https://hal.inrae.fr/hal-02785577>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Guinea fowl domestication : new insights from the first whole genome assembly and the pool sequencing of wild and domestic populations.

Simon Boitard¹, Noémie Thebault¹, Guiguigbaza Kossigan Dayo², Valentine Yapi-Gnaore², Issaka Youssao³, Michèle Tixier-Boichard⁴, Wes Warren⁵, Alain Vignal¹, Xavier Rognon⁴

(1) GenPhySE, Castanet-Tolosan, France. (2) CIRDES, Bobo-Dioulasso, Burkina-Faso. (3) Université d'Abomey-Calavi, Cotonou, Bénin. (4) GABI, Jouy-en-Josas, France (5) McDonnell Genome Institute, St. Louis, Missouri.

II Joint Congress on Evolutionary Biology – Montpellier 2018



- **Village poultry in Africa**, intensive livestock farming for meat production in **Europe** (mostly France and Italy).

- **Natural range in sub-Saharan Africa**, 8 subspecies.
- **Domestication** may have occurred **about 2,000 years BP** in Mali and Sudan, archaeozoological and artistic data (Larson and Fuller, 2014).
- Food supply, farm keeper and parasite / snake hunter.
- **Diffused** in the Mediterranean world during the Roman Empire, and throughout the world **from the 16th century**.
- **Genetic diversity** studied from mtDNA (Walker *et al*, 2004; Adeola *et al*, 2015) or microsatellites (Kayang *et al*, 2010; Weimann *et al*, 2016).


- **Sequencing** of a male sample on Illumina HiSeq 2500, at about 70X coverage (3 and 8kb paired-end reads).
- **Assembled** using ALLPATHS2 (Maccallum *et al*, 2009)
 - Male genome of **1.04Gb**
 - **2,739 scaffolds (N50 7.8Mb)**
- **Alignment to the chicken genome** (Galgal5) + known **cytogenetic rearrangements** (Shibusawa *et al*, 2002)
 - 29 autosomes + Z + mtDNA + linkage group LGE64.
- Assembly quality comparable to that of other Galliformes (chicken, quail, common duck).

Pool-sequencing of 12 samples


- **Africa, wild** : **AFS-w** (South Africa, $n = 3$, 14X), **KOF-w**, **YAB-w** (Burkina Faso, $n = 8$, 20X).
- **Africa, domestic** : **SDA-t**, **SKO-t**, **DOR-t** (Burkina Faso, $n = 5$, 10X), **BEN-t** (Bénin, $n = 15$, 16X).
- **Europe, domestic** : **GOD-t**, **HAR-t** (Hungaria, $n = 30$, 17X).
- **Europe, selected** : **BEG-s** (France, $n = 12$, 15X), **GAL-s** (France, $n = 29$, 37X), **GRI-s** (France, $n = 20$, 38X).

→ **80 million SNPs** with allele frequency called in all pools and $MAF \geq 0.05$.

Genetic structure reflects sub-species and domestication




Reduced diversity in domestic populations



Population tree confirms genetic structure and diversity











Whole Genome tree



- Estimated by FLK (Bonhomme *et al*, 2010) from allele frequencies at ≈ 8.9 million SNPs.
- Domestic samples from Burkina-Faso merged (**BUR-t**), AFS-w used as outgroup.

- **Selective sweep** signatures detected **within each population** using Pool-HMM (Boitard *et al*, 2012; 2013).
→ thousands of candidate regions.
- 5 regions related to **domestication** : detected in 6 out of 7 domestic populations, undetected in wild populations.
- 31 regions specific to **Europe**, 64 to **intensive meat production**.
- Combining candidates from the 3 categories leads to 0.6% of the genome, including 223 genes (**2-fold enrichment**).

Domestication candidate region



Candidate genes and mutations

- Within the 5 domestication regions, **30 genes in total** (from 1 to 16 per region).
- **29** of these genes **include candidate variants**:
allele frequency ≥ 0.75 in all domestic populations and ≤ 0.25 in all wild populations.
- **2** of these genes include **1 candidate variant with functional effect (missense)** : *PAPPA2* and *EDNRB*.

Example 1 : *PAPPA2*

- **Only gene** in a **candidate region** of 21 kb, at position 7.3 Mb on chromosome 7.
- *PAPPA2* (Pregnancy-Associated Plasma Preproprotein-A2) is a **modulator of IGF-I** (Fujimoto *et al*, 2017).
- Mutations associated to **body size** in at least 3 vertebrate species: mice (Christians *et al*, 2006), humans (Lango Allen *et al*, 2010) and cows (Bouwman *et al*, 2018).

Example 2 : *EDNRB*

- Expressed in **melanocytes deriving from the neural crest** → potential domestication gene (Wilkins *et al*, 2014).
- Mutations associated to **white spotted patterns** in chickens (Kinoshita *et al*, 2014) and **pigmentation disorders** in other species (Dang *et al*, 2011; Ayala-Valdovinos *et al*, 2016).
- In our African samples, **strong pigmentation differences between wild** (left) **and domestic** (right) animals.



- **White spotted** patterns found among **village chickens in Benin**, said to be associated with luck and wealth (Chrysostome *et al*, 2013).
- **Selection for tameness via *EDNRB*?**
- **11 other genes in the candidate region** of *EDNRB*, 175 kb long at 11.1-11.3 Mb on chromosome 8.
- **Two of them associated to behavioral traits:** *VAMP7* (mice, Danglot *et al*, 2012) and *TMLHE* (humans, Ziats *et al*, 2015; quails, Fariello *et al*, 2017).

- **Genome assembly of the Guinea fowl.**
- First **study of genetic diversity** in this species **based on genome-wide data.**
- **Domestic** populations **cluster together**, and have **reduced genetic diversity.**
- **Domestication** likely occurred **in Western Africa.**
- Domestication **candidate genes** and mutations, related to body size, pigmentation or behaviour.

Acknowledgements

Sample and data collection:

- **Alain Vignal**, GenPhySE, Castanet-Tolosan, France
- **Michèle Tixier-Boichard, Xavier Rognon**, GABI, Jouy-en-Josas, France
- **Guiguibaza Kossigan Dayo, Valentine Yapi-Gnaore**, CIRDES, Bobo-Dioulasso, Burkina-Faso
- **Issaka Youssao**, Université d'Abomey-Calavi, Cotonou, Bénin
- **Cécile Berthouly-Salazar**, DIADE, Montpellier, France
- **Judit Barna**, Research Centre for Farm Animal Gene Conservation, Gödöllő, Hungary
- **Daniel Guémené**, SYSAAF, Nouzilly, France
- **GALOR SAS**, subsidiary of Groupe Grimaud, Amboise, France

Genome sequencing, assembly and annotation:

- **Alain Vignal, Noémie Thebault**, GenPhySE
- **Wes Warren**, McDonnell Genome Institute, St. Louis, Missouri
- **Francoise Thibaud-Nissen**, NCBI, Bethesda, Maryland

Computing Facilities:

- Genotoul bioinformatics platform Toulouse Midi-Pyrénées.

Funding:

- ITAVI SeqVol program
- INRA, Animal Genetics division
- AgroParisTech