



OH1 from Orf virus: a new tyrosine phosphatase that displays distinct structural features and triple substrate specificity

Gwenaëlle André-Leroux

► To cite this version:

Gwenaëlle André-Leroux. OH1 from Orf virus: a new tyrosine phosphatase that displays distinct structural features and triple substrate specificity. Conférences du CECAM, 2017, Evry, France. hal-02785772

HAL Id: hal-02785772

<https://hal.inrae.fr/hal-02785772>


Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OH1 from Orf virus: a new tyrosine phosphatase


Distinct structural features & triple substrate specificity



Gwenaëlle André-Leroux
MaIAGE, INRA Jouy-en-Josas
Mathématique et Informatique Appliquées du Génome à l'Environnement

Workshop: « Modelling bio-molecular interactions: A multi-scale approach joining theory and experiments »


OH1 from Orf virus: a new tyrosine phosphatase



Outline

- Orf virus introduction
- OH1 structural characterization *in silico/in vitro*
- OH1 functional characterization *in silico/in vitro*
- Conclusions & perspectives


OH1 from Orf virus: a new tyrosine phosphatase



Outline

- Orf virus introduction
- OH1 structural characterization *in silico/in vitro*
- OH1 functional characterization *in silico/in vitro*
- Conclusions & perspectives

Orf virus





Orf Poxvirus family
Like Vaccinia & Variola virus

dsDNA virus
Large 200-300 nm
Envelope

Orf = old english for « rough »



Orf virus




Orf Poxvirus family
Like Vaccinia & Variola virus

Orf = old english for « rough »

dsDNA virus
Large 200-300 nm
Envelope



Bovine papular stomatitis virus
Parapoxvirus of red deer in New Zealand
Pseudocowpox virus



Vaccinia virus
Variola virus
Cowpox virus

Orf virus



- Is responsible for Ecthyma disease primarily in sheep and goats
- Is contagious pustular dermatitis
- Replicates in keratinocytes and epithelial cells of the oral mucosa.
- Is reported worldwide
- Is purulent for 7 weeks
- Prevents lambs from sucking and cause weight loss
- Causes mastitis to mother ewes
- In its severe form on immune compromised animals can lead to death

Orf virus



- Is responsible for Ecthyma disease primarily in sheep and goats
- Is contagious pustular dermatitis
- Replicates in keratinocytes, the immune sentinels of skin.
- Is reported worldwide
- Is purulent for 7 weeks
- Prevents lambs from sucking and cause weight loss
- Causes mastitis to mother ewes
- In its severe form on immune compromised animals can lead to death

Highly contagious by nature

Zoonotic through direct contact

Live vaccine that can cause reinfection



Vaccine: immunity for 6-8 months

Secondary infections

Economical impact +++

Orf virus : new strain isolated in Uruguay in 2015

Strain UY1107 was fully sequenced : shows 90 essential and conserved genes

➡ Among them, a putative virulence factor was identified, by analogy with Vaccinia and Variola viruses


➡ Codes for a putative tyrosine phosphatase

➡ Deposited at genbank ID: KY651216

➡ Hypothesis: KY651216 is a Tyr-phosphatase responsible for dephosphorylation of phosphorylated substrates
Signaling mediators: Tyrosine phosphatase/kinase are molecular switches that turns on/off substrates

OH1 from Orf virus: a new tyrosine phosphatase

Distinct structural features & triple substrate specificity




Outline

- Orf virus introduction
- OH1 structural characterization *in silico/in vitro*
- OH1 functional characterization *in silico/in vitro*
- Conclusions & perspectives

Putative tyrosine phosphatase

- ➡ High 40% similarity with Vaccinia and Variola Tyr-phosphatase
- ➡ Strict conservation of the active site of a Tyr-phosphatase


Active site Aspartate loop



Putative tyrosine phosphatase


- ➡ High 40% similarity with Vaccinia and Variola Tyr-phosphatase
- ➡ Strict conservation of the active site of a Tyr-phosphatase

Ser to Cys substitution within helix α_1



Putative tyrosine phosphatase


- Homology modeling of Tyrosine phosphatase with 3D Vaccinia and Variola VH1 as 3D templates: Chains A then B.
- Reconstruction of the dimer by superposition onto the dimer template.
- Minimisation with Charmm to relax the side chains



Putative tyrosine phosphatase

➡ The main differences are located at the interface


➡ Helix α_1 at the N-terminal end is largely engaged in domain swapping



Putative tyrosine phosphatase




➡ The main differences are located at the interface

➡ Helix α_1 at the N-terminal end is largely engaged in domain swapping



Orf virus : tyrosine phosphatase

- Protein purified to homogeneity Histrap and size-exclusion chromatography wt and C112S
- Crystallisation screening using sitting drop vapor diffusion for both



Cristal conditions
200 mM NH4SO4,
100mM NaAc, pH 4,6
PEG MME 2K 30%

Beamline Proxima1
2 sets of data collection



Data statistics 5NCR
P₂₁
a, b, c (Å) 49.56 63.55 55.39
 α , β , γ (°) 90.00, 97.07 90.00

Refinement of the best at 1.89 Å resolution
Molecular replacement using Phaser
Model building with Coot
Refinement cycles within Refmac CCP4 package and Buster
Rwork/Rfree 18.72/21.69



Crystal grew after 3 days for C112S
No crystal after 3 months for wt

Pdb deposition 5NCR

Orf virus : tyrosine phosphatase

Solved structure reveals:

- ➡ Conserved fold of Tyr phosphatase DUSP- Dual Specificity Phosphatase
- ➡ Conserved active site
- ➡ Covalent dimer
- ➡ Ser to Cys15 substitution engages a disulfide bridge




- ➡ Covalent dimer
- ➡ 7 residues insertion results in one extra α_5 helix turn
- ➡ Extra interaction at Pro 141




Orf virus : tyrosine phosphatase

Solved structure reveals a dimer organization distinct from Vaccinia phosphatase



➡ The interface engages the same cluster of helices α_1 , α_5 et α_6



Interface of 678 Å²



Interface of 978 Å²




Pdb id 3cm3 Koksai et al, JBC 2009




Orf virus : tyrosine phosphatase

Solved structure reveals a dimer organization distinct from Vaccinia phosphatase

➡ The interface engages the same cluster of helices α_1 , α_5 et α_6






OH1 α_{1A} takes the position of VH1 α_{1B}
OH1 α_{1B} takes the position of VH1 α_{1A}



Orf virus : tyrosine phosphatase

Solved structure reveals a dimer organization distinct from Vaccinia phosphatase

➡ Incompatible with domain swapping




Orf virus : tyrosine phosphatase


Solved structure reveals a dimer organization distinct from Vaccinia phosphatase

➡ Not a crystallization bias because

- Production of recombinant OH1



- Size exclusion chromatography




OH1 is a dimer in solution

C15S shows also a dimer (data not shown)

C15S is not mandatory for a dimerization

- Ultra Analytical Centrifugation



40 kDa = 2*20 kDa

Detection of Parapox specific features of OH1


Cys 15 and 6 residues insertion conserved but only in Parapoxviruses

Phylogenetic studies explore how parapoxvirus associates N-terminal Ser15 to Cys substitution


- Multiple sequence alignment of 64 phosphatases from Chordopox family
- Tcofee
- Position with >50% gaps filtered
- Correlation techniques based on mutual informations

Ala40 locates in the vicinity of the active site

Tyr152 could stabilize helix α_1




OH1 from Orf virus: a new tyrosine phosphatase



Highlights

- OH1 displays distinct structural features compared to VH1 phosphatases
- Orf virus OH1 is a covalent dimer involving the N-terminal Cys15
- Orf virus OH1 possibly depicts the structure of Parapoxvirus genus phosphatases

OH1 from Orf virus: a new tyrosine phosphatase




Outline

- Orf virus introduction
- OH1 structural characterization *in silico/in vitro*
- OH1 functional characterization *in silico/in vitro*
- Conclusions & perspectives

Tyrosine phosphatase

➡ OH1 shows a phosphatase activity *in vitro*



Monitoring of dephosphorylation by absorbance at 405nm of pNP

Phosphatase activity assayed at 37°C using the artificial substrate pNPP

Michaelis-Menten Kinetics constants

OH1 wt

$$K_m = 2.6 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$


OH1 C15S

$$K_m = 2.0 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$

Tyrosine phosphatase

➡ OH1 shows a phosphatase activity *in vitro*



Phosphatase activity assayed at 37°C using the artificial substrate pNPP

Michaelis-Menten Kinetics constants

OH1 wt

$$K_m = 2.6 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$

OH1 C15S



$$K_m = 2.0 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$

➡ OH1 shows a DUSP –Dual Specificity Phosphatase *in vitro*

Active on p-Tyr containing protein


Active on p-Thr 17mer peptide



Dose dependent manner

Tyrosine phosphatase

➔ OH1 shows a phosphatase activity *in vitro*



Phosphatase activity assayed at 37°C using the artificial substrate pNPP

Michaelis-Menten Kinetics constants

OH1 wt

$$K_m = 2.6 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$

OH1 C15S



$$K_m = 2.0 \text{ mM}$$

$$K_{\text{cat}} = 0.08 \text{ s}^{-1}$$

➔ OH1 shows a DUSP –Dual Specificity Phosphatase *in vitro*



Active on p-Tyr containing protein

Active on p-Thr 17mer peptide




Dose dependent manner

➔ OH1 accommodates p-Tyr in the active site




A new tyrosine phosphatase

➔ OH1 highlights a new activity *in vitro*: a triple specificity





Phosphoinositides can be dephosphorylated

PtdIns3P and PtdIns5 (3,5)P2 can be dephosphorylated



➔ Docking evidences PdtIns bind in the pocket



The phospho-sugar part can be accommodated

The phospho-group in 5' binds into the active pocket

Relevant physiologically?

A new tyrosine phosphatase

➔ Why this phospholipid specificity should be relevant?

PtdIns

- Belong to phosphatidyl glycerides
- Compose the cytosolic side of eukaryotic cell membrane
- Regulate membrane homeostasis
- Are signaling lipids



OH1 could extract and recycle host phospholipids to form viral envelop during virion maturation

- Viral envelop allows to:
- Escape from the cell,
 - Stay stable
 - Get protected from immune system
 - Reinfect cell

Evolution in favor of enveloped virus : Influenza, HIV, Ebola Virus






Figure 11-3
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W.H. Freeman and Company




OH1 from Orf virus: a new tyrosine phosphatase



Highlights

- OH1 displays distinct structural features compared to VH1 phosphatases
- Orf virus OH1 is a covalent dimer involving the N-terminal Cys15
- Orf virus OH1 possibly depicts the structure of Parapoxvirus genus phosphatases
- Orf virus OH1 is a dual specificity phosphatase that presents activity towards PInsP *in vitro*

OH1 from Orf virus: a new tyrosine phosphatase




Outline

- Orf virus introduction
- OH1 structural characterization *in silico/in vitro*
- OH1 functional characterization *in silico/in vitro*
- Conclusions & perspectives

Interaction with STAT1


➔ Hypothesis: by analogy with Vaccinia and Variola viruses, OH1 could dephosphorylate STAT1



STAT : Signal Transducers and Activators of Transcription

- STAT1 is activated upon phosphorylation of its tyrosine Tyr701 -> STAT1 (P)
- STAT1 (P) proteins homodimerize
- STAT1 (P) homodimer is now capable to translocate to the nucleus
- STAT1 (P) transactivates Stat-responsive genes
- STAT1 (P) is involved in cardiac hypertrophy, apoptosis, and inflammation.

OrfV dephosphorylates STAT1



Format: Abstract ▾

Send to ▾

Virus Res. 2015 Oct 2;208:180-8. doi: 10.1016/j.virusres.2015.06.014. Epub 2015 Jun 22.

Orf virus inhibits interferon stimulated gene expression and modulates the JAK/STAT signalling pathway.

Harvey R¹, McCaughan C², Wise LM³, Mercer AA⁴, Fleming SB⁵.



Author information

Abstract

Interferons (IFNs) play a critical role as a first line of defence against viral infection. Activation of the Janus kinase/signal transducer and activation of transcription (JAK/STAT) pathway by IFNs leads to the production of IFN stimulated genes (ISGs) that block viral replication. The Parapoxvirus, Orf virus (ORFV) induces acute pustular skin lesions of sheep and goats and is transmissible to man. The virus replicates in keratinocytes that are the immune sentinels of skin. We investigated whether or not ORFV could block the expression of ISGs. The human gene GBP1 is stimulated exclusively by type II IFN while MxA is stimulated exclusively in response to type I IFNs. We found that GBP1 and MxA were strongly inhibited in ORFV infected HeLa cells stimulated with IFN-γ or IFN-α respectively. Furthermore we showed that ORFV inhibition of ISG expression was not affected by cells pretreated with adenosine N1-oxide (ANO), a molecule that inhibits poxvirus mRNA translation. This suggested that new viral gene synthesis was not required and that a virion structural protein was involved. We next investigated whether ORFV infection affected STAT1 phosphorylation in IFN-γ or IFN-α treated HeLa cells. We found that ORFV reduced the levels of phosphorylated STAT1 in a dose-dependent manner and was specific for Tyr701 but not Ser727. Treatment of cells with sodium vanadate suggested that a tyrosine phosphatase was responsible for dephosphorylating STAT1-p. ORFV encodes a factor, ORFV057, with homology to the vaccinia virus structural protein VH1 that impairs the JAK/STAT pathway by dephosphorylating STAT1. Our findings show that ORFV has the capability to block ISG expression and modulate the JAK/STAT signalling pathway.

Copyright © 2015 Elsevier B.V. All rights reserved.

Interaction with STAT1





Harvey et al, 2015

Modeling of OH1/STAT1 direct interaction


Interaction with STAT1

➔ Running hypothesis: OH1 dephosphorylates STAT1 by a direct interaction

- Modeling of the missing loop and phosphorylation at Tyr 701
- Docking of Protein/Protein interaction using HADDOCK



OH1 from Orf virus: a new tyrosine phosphatase




Perspectives


- OH1 /STAT1 needs further investigations *in silico*
- Pull-down *in vitro* experiments show that OH1 interacts directly with STAT1
- Pull-down *in vitro* experiments show that OH1 interacts with GAPDH - Glyceraldehyde-3-phosphate dehydrogenase
- GAPDH is a house-keeping protein involved in glycolysis and modulation of the organization of the cytoskeleton
- GAPDH could not only be a metabolic enzyme but in immunity also



Acknowledgements





Virology




Dario Porley
Mabel Berois

Crystallization





Danilo Segovia
Frederik Saul
Patrick Weber
Ahmed Haouz

Biochemistry



Danilo Segovia
Mariano Martinez
Andrea Villarino


Crystallography, modeling & phylogeny



Danilo Segovia
Dario Porley
Mahendra Mariadassou

OH1 from Orf virus: a new tyrosine phosphatase

Distinct structural features & triple substrate specificity



Thank you very much for your kind attention