



**HAL**  
open science

## **WOODNET: Connectivity patterns and processes along a gradient of European landscapes with woody vegetation and spatial heterogeneity. WP 2: Landscape legacies and species distribution**

Audrey Alignier, Déborah Closset-Kopp, Guillaume Decocq, Aude Ernoult, Cendrine Mony

### ► To cite this version:

Audrey Alignier, Déborah Closset-Kopp, Guillaume Decocq, Aude Ernoult, Cendrine Mony. WOODNET: Connectivity patterns and processes along a gradient of European landscapes with woody vegetation and spatial heterogeneity. WP 2: Landscape legacies and species distribution. WOODNET meeting, Nov 2019, Louvain-la-Neuve, Belgium. pp.39. hal-02786232


**HAL Id: hal-02786232**

**<https://hal.inrae.fr/hal-02786232>**

Submitted on 4 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Louvain meeting  
November, 6-8, 2019

France

Spain

Belgium



WOODNET: Connectivity patterns and processes along a gradient of European landscapes with woody vegetation and spatial heterogeneity

[woodnetweb.wordpress.com/project-fr/](http://woodnetweb.wordpress.com/project-fr/)

WP2:

Landscape legacies and species distribution

A. Alignier, D. Closset-Kopp, G. Decocq, A. Ernoult, C. Mony


## WP2 sub-tasks

- Task 2.1: Drivers of actual local species assemblages in hedgerows
- Task 2.2: Disentangling dispersal from recruitment limitation in hedgerow corridors
- Task 2.3: Measuring landscape connectivity: a landscape genetic approach


## Task 2.1. Understanding the main drivers of connectivity dynamics

**Objective:** Quantify the respective role of habitat quality and connectivity (using past and present landscape composition and configuration as proxies) in explaining local biodiversity in hedgerows

Connectivity measures  
over time



Local variables of hedgerow  
quality



Plant and carabid assemblages in  
woodlands



**Specific structure**

**Functional structure**

## Task 2.1. Understanding the main drivers of connectivity dynamics

**Objective:** Quantify the respective role of habitat quality and connectivity (using past and present landscape composition and configuration as proxies) in explaining local biodiversity in hedgerows

- Detect plant/carabid\* species richness response to changes over time of both local and connectivity variables

Comparison between forest specialists and generalists

- Detect plant trait syndromes to connectivity dynamics

Different responses depending on plant traits

- Generalize patterns across taxonomic groups (birds)

## Task 2.1. Understanding the main drivers of connectivity dynamics

### Methods


#### 1- Assessing connectivity changes over time

- Mapping from aerial photographs: landcover and hedgerows
- Selection of 6 dates: 1952, 1974, 1985, 2000, 2006 and 2016 in relation with various policies (Common Agricultural Policy, land reallocation program)


## Task 2.1. Understanding the main drivers of connectivity dynamics

Land-use maps (*source: Zone Atelier Armorique*)

**Rennes**




Hedgerows and woodlots maps (*source: Kermap*)




# Task 2.1. Understanding the main drivers of connectivity dynamics

Rennes

## Landscape changes in the study area



Increase of croplands  
Decrease of grasslands



Increase of crop field size (reallotment)  
Decrease of woodlot areas



# Task 2.1. Understanding the main drivers of connectivity dynamics

## Amiens

1949

Bocage



Openfield



1974

Bocage



Openfield




1984

Bocage



Openfield



Bocage



Openfield



# Task 2.1. Understanding the main drivers of connectivity dynamics

## Amiens

2001

Bocage



Openfield



- forest
- crops
- orchard
- urban area
- grassland
- water

2013

Bocage



- forest
- crops
- orchard
- urban area
- grassland
- water

Openfield



0 1 2 km

### Methods



#### 1- Assessing connectivity changes over time

- Mapping from aerial photographs: landcover and hedgerows
- Selection of 6 dates: 1952, 1974, 1985, 2000, 2006 and 2016 in relation with various policies (Common Agricultural Policy, land reallocation program)
- Measuring connectivity indices for each date

## Task 2.1. Understanding the main drivers of connectivity dynamics

### Patch isolation measures

- Euclidean distances
- Resistance distances



$$dF_k^* = \frac{\sum_{i=1, i \neq k}^{n-1} P_{ik}^*}{\sum_{i=1}^n \sum_{j=1, i \neq j}^n P_{ij}^*}$$

$p^*_{ij}$  is the maximum product probability of all possible paths between 2 forest fragments,  $i$  and  $j$ , in the landscape

$dF^*_k$  assessed the percentage of dispersal flux among all forest fragments

## Task 2.1. Understanding the main drivers of connectivity dynamics

### Methods

#### 2- Analyze plant (and bird) assemblages

##### 2a - Floristic surveys in woodlands


- Rennes: 50 woodlands sampled (PhD thesis L. Uroy)  
Woodland standardized in age, management, size

Protocol: 8 quadrats (10 x 5m for a total of 200-500m<sup>2</sup>)

- Amiens: 91 woodlands sampled (PhD thesis A. Jamoneau)  
n=62/29 in the bocage/openfield 5\*5km window

Protocol: exhaustive survey of vascular plant species over the whole patch area + 1 quadrat of 1000 m<sup>2</sup> in 30 patches (15 in each window)

- + Rennes: 30 woodlands sampled for birds (additional work to Woodnet: S. Croci & A. Butet)



Location of sampled woodlands (in green)

# Task 2.1. Understanding the main drivers of connectivity dynamics

## Methods

### 2- Analyze plant (and bird) assemblages

#### 2b - Floristic surveys in hedgerows

- Rennes: 30 hedgerows (see WP1)
- Amiens: 49 hedgerows (see WP1)  
n=31/18 in bocage/openfield window

# Task 2.1. Understanding the main drivers of connectivity dynamics

## Methods

### 2- Analyze plant (and bird) assemblages


- Ecological and functional traits databases

Establishment of specialist/generalist lists

Establishment of functional traits list for plants (LEDA, TRY databases)

Dispersal traits

Same work has to be done for birds



### Methods

### 3- Determine hedgerow ancienty



We assembled a chronosequence of 99 dated (1725-2008) hedgerows, which were surveyed for forest plant species and a number of local-proximal descriptors.





# Task 2.1. Understanding the main drivers of connectivity dynamics

## Preliminary results – Analysis of woodland assemblages

### Rennes

Herbaceous species: 0-28 species per woodland

Most abundant species:



## Task 2.1. Understanding the main drivers of connectivity dynamics

### Preliminary results – Analysis of woodland assemblages

#### Amiens


Herbaceous species: 11-83 species per woodland

Most abundant species: *Arum maculatum*, *Ranunculus ficaria*, *Geum urbanum*, *Rubus fruticosus* agg., *Glechoma hederacea*, *Hedera helix*, *Urtica dioica*, *Galeopsis tetrahit*, *Stachys sylvatica*, *Adoxa moschatellina*

## Task 2.1. Understanding the main drivers of connectivity dynamics

### Preliminary results – Analysis of woodland assemblages


#### Rennes



## Task 2.1. Understanding the main drivers of connectivity dynamics

### Preliminary results – Analysis of woodland assemblages

#### Rennes



## Task 2.1. Understanding the main drivers of connectivity dynamics

Amiens

### Preliminary results – Analysis of hedgerow assemblages


GLM:  $R^2 = 0.62$ ;  $R^2$  adjusted = 0.59

	Coefficient	S. E	Student (t)	(p)
Richness ( $X_i = 0$ )	79.95	16.245	4.922	(***) <0.001
Ancienty	-0.029	0.009	-3.407	(***) <0.001
Length	0.006	0.009	0.655	(n.s.) 0.514
Width	1.461	0.873	1.674	(n.s.) 0.097
Height	0.518	0.209	2.478	(*) 0.015
Height heterogeneity	-0.113	0.144	-0.780	(n.s.) 0.437
Connectivity	11.007	2.661	4.136	(***) <0.001
Forest area	0.000	0.000	1.258	(n.s.) 0.211
Land Use Intensity	-2.137	0.539	-3.966	(***) <0.001

Hedgerow age and length interacted to increase forest plant species richness.

Hedgerows attached to woodlands hosted systematically more species = +11

Taller and wider hedgerows with a lower intensity of adjacent land uses hosted more forest plant species.



## Task 2.1. Understanding the main drivers of connectivity dynamics

### Difficulties

Not enough sample sites for carabids in Rennes (10 woodlands sampled for WP1)

Not the same protocole between Amiens and Rennes for plant sampling

➡ need to check that they are comparable

### Planning for the end of the sub-task

November 2019 - February 2020: Statistical analysis and writing

Submission deadline: March 2020

Paper 1: Effect of connectivity dynamics on functional structure of forest plants  
(Amiens and Rennes sites)

## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

**Objective:** Disentangling the effect of dispersal from recruitment limitation of forest species in hedgerows

➤ **Dispersal limitation**

Seed arrival in the site

➤ **Recruitment limitation**

Abiotic factors (habitat quality)

Biotic factors (competition with existing vegetation)

➤ **Seed germination ability**

Seed germination power

## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

### Methods

#### 1 - Plant species selection

- Seed collection on mature individuals (June-September 2017) in spatially distant populations over the ZA Armorique and in the Amiens Region + Transplants collection (one population only)
- 10 to 12 populations sampled
- Same plant species list between the two sites (10 for Rennes *versus* 15 for Amiens)



## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

### Methods

#### 1 - Plant species selection

##### Specialist plant species:

*Hyacinthoides non-scripta* (A,R)

*Melica uniflora* (A,R)

*Carex sylvatica* (A,R)

*Stachys sylvatica* (A,R)

*Viola reichenbachiana* (A,R)

*Lamium galeobdolon* (A,R)

*Oxalis acetosella* (A)

*Stellaria holostea* (A)

*Poa nemoralis* (A)

*Milium effusum* (A)

*Galium odoratum* (A)

##### Generalist plant species:

*Potentilla sterilis* (R)

*Veronica hederifolia* (A,R)

*Circaea lutetiana* (A,R)

*Lapsana communis* (A,R)

*Senecio ovatus* (A)

*Aegopodium podagraria* (A)

*Fragaria vesca* (A)



# Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

## Methods

### 2 - Site selection

- 5 hedgerows and 1 woodland *per* site
- Criteria for hedgerows standardization (age < 20 yrs, tree composition, structure, no management over the study period...)



**Amiens**



# Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors


## Methods

### 3 - Experimental design


Species sowing



With vegetation




Without vegetation




Species transplantation



With vegetation



Without vegetation



1 control



Greenhouse

- 4 replicates per hedgerow
- 25 seeds per quadrat
- Each quadrat = 20x20cm

- 2 replicates per hedgerow
- 5 transplants per quadrats
- Each quadrat= 20x20cm

- 100 seeds

In total, 1200 seeds sown *per species*



## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

Amiens



Seed sowing and transplantation

Seed germination in greenhouse

## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

### Methods

#### 4 - Surveys

3 years after launching (April 2018)

Number of seeds germinated (greenhouse)

April, June & October 2018-2020

Number and size of the germinated seeds

April, June & October 2019-2020


Number and size of the transplants

April, June & October 2019-2020

+ environmental measurements: Hedgerow structure, soil pH, LAI (Leaf Area Index)


## Preliminary results (sowing)

- Very low germination *in situ* compared to greenhouse
- *Circea lutetiana* had very low germinative power




## Preliminary results (sowing)

- Overall, more germination in woodlands than in hedgerows
- Better seed germination rates in « without vegetation » quadrats (excluding competition)



## Preliminary results (transplants)


- Highest survival of the transplants in the first year of the study
- Good persistence (> 40% for 6 / 9 species)





## Preliminary results (transplants)


- Highest survival rates in « without vegetation » quadrats (excluding competition)



# Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

## Preliminary results


Amiens



# Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

## Preliminary results

Amiens



## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

Amiens

### Preliminary results

Proportion of individuals which successfully **germinated** or **persisted** within a given 20cm × 50cm sub-quadrat (i.e. the response variable)

- Habitat abiotic conditions (hedgerows vs. forest)
- Competition (disturbed vs. un-disturbed)
- Species ID as random variable

Symbol	Candidate Model	AIC
MG1	$Y \sim \text{competition (Y/N)}$	220.57
MG2	$Y \sim \text{habitat (H/F)}$	154.67
<b>MG3</b>	<b><math>Y \sim \text{competition (Y/N) + habitat (H/F)}</math></b>	<b>147.95</b>
MG4	$Y \sim \text{competition (Y/N) * habitat (H/F)}$	164.02
ME1	$Y \sim \text{competition (Y/N)}$	544.27
ME2	$Y \sim \text{habitat (H/F)}$	543.25
ME3	$Y \sim \text{competition (Y/N) + habitat (H/F)}$	523.11
<b>ME4</b>	<b><math>Y \sim \text{competition (Y/N) * habitat (H/F)}</math></b>	<b>517.65</b>

Best candidate model for germination

Best candidate model for persistence

## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

### Preliminary conclusions

- Some forest species succeeded in germinating into hedgerows (i.e. 7/10 species in Rennes; *Veronica hederifolia* & *Stellaria holostea* in Amiens) → **There is dispersal limitation**
- Most of the forest herb species are limited by the hedgerows' habitat conditions → **There is recruitment limitation in hedgerows**
- There is more germination and transplants survival in “without vegetation” quadrats → **There is a competition effect in hedgerows**
- 9/13 studied forest plant species in Amiens and 8/10 in Rennes succeeded to establish in hedgerows → **There is a low persistence rate in hedgerows**

## Task 2.2. Disentangling dispersal from recruitment limitation in hedgerow corridors

### Difficulties

Low germination power of seeds in the hedgerows (dry summer, high litter mater; germination ability for Amiens as 6/17 species succeeded to germinate in the greenhouse)

Important delay in plant responses that need to have a long-duration experiment (3 years) and surveys after the end of the Woodnet project

### Planning for the end of the sub-task

June 2020: Last sampling campaign

Submission deadline: Fall 2020

Paper 2: Disentangling dispersal, establishment and competition effects on forest plants (Amiens and Rennes sites)

# Publications and data valorisation (1/2)

## Symposium

- Special session: Bergès & Mony 2018. Current challenges in landscape ecology: habitat amount, landscape connectivity, landscape history.
- Closset-Kopp & Decocq G., 2018. Connectivity between forest patches in changing agricultural landscapes: times also matters! (SFE Rennes)
- Mony et al., 2018. Biodiversity response to landscape connectivity dynamics (SFE Rennes)
- Uroy et al., 2019. Trame verte et bleue et biodiversité: une vue d'ensemble des outils et méthodes pour caractériser la connectivité paysagère et son effet sur les communautés végétales. Colloque »La cartographie de la flore, un outil au service des politiques publiques de la biodiversité », Nantes.
- Closset-Kopp et al., 2018. Are rural hedgerows effective corridors for forest plant species? Yes, but this is a matter of time and spatial connectedness! IAVS 2018. Montana

## Publications

- Paper 1 on going
- Historical continuity and spatial connectivity ensure hedgerows are effective corridors for forest plants: evidence from the species-time-area relationship. Lenoir et al. submitted



### Collaboration

Collaboration with K. Litza from M. Diekmann's group – Univ. Bremen, Germany  
(meta-analysis on forest plant species throughout Europe)

