



HAL
open science

Predictive diffusion of palms pests *Rhynchophorus ferrugineus* & *Paysandisia archon* and perspectives about biological control

Robert Castellana, Elisabeth Tabone, Mathilde Capelli, Catherine Gignoux,
M. Cristofaro, Hervé Pietra

► To cite this version:

Robert Castellana, Elisabeth Tabone, Mathilde Capelli, Catherine Gignoux, M. Cristofaro, et al.. Predictive diffusion of palms pests *Rhynchophorus ferrugineus* & *Paysandisia archon* and perspectives about biological control. Annual Meeting of the European Network of Palm Scientists (EUNOPS), May 2019, Paris, France. 2019. hal-02786340

HAL Id: hal-02786340

<https://hal.inrae.fr/hal-02786340v1>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Predictive diffusion of palms pests *Rhynchophorus ferrugineus* & *Paysandisia archon* and perspectives about biological control

Castellana R. (Phoenix Project), Tabone E. & Capelli M. (Inra Antibes), Gignoux C. (Fredon Corse), Cristofaro M. (Enea Italy) & Pietra H. (Sauvons Nos Palmiers)

More than 35 species of palms are actually infested by *Rhynchophorus ferrugineus* & *Paysandisia Archon*, according to the observations of our network of French & Italian Botanical Collections of ca 100 ornamental palm species. Predictive diffusion of these pests may also concern in the future the 2500 species of wild populations of palms in Asia, South America and Africa. The adaptation of these pests to new environments and climatic conditions, and the difficulties to detect early symptoms of infestation, were determining a failure of all eradication and suppression conventional control programs like Integrated Pest Management, based mainly upon the use of chemical or agronomic approaches. Our French & Italian Network is associated with the researches about biological control by Inra (France) & Enea (Italy).




Figure 1: Phases of the cycle of *Paysandisia archon* from left to right: eggs, larva, cocoon, adults (male above and female below). (Picture kindly provided by Jean-Benoît Peltier, INRA, France)

Our observations about infestation concerning ca 100 ornamental palms *Paysandisia archon* (PA) versus *Rhynchophorus ferrugineus* (RPW)

- | | |
|--|--|
| Acoelorrhaphe wrightii | Brahea armata & edulis |
| Arenga engleri | Butia capitata & yatay |
| Brahea aculeata, armata, dulcis, edulis | Chamaerops humilis |
| Chamaerops humilis | Howea fosteriana |
| Howea fosteriana | Jubaea chilensis |
| Jubaea chilensis | Livistona australis, chinensis, decora |
| Livistona australis, chinensis, decora | (decipiens), saribus |
| (decipiens), saribus | Nannorrhops arabica |
| Nannorrhops arabica | Phoenix dactylifera, reclinata, robellini, |
| Phoenix dactylifera, reclinata, robellini, | theophrasti |
| theophrasti | Sabal bermudana, mauritiiformis, minor, |
| Sabal bermudana, mauritiiformis, minor, | palmetto |
| palmetto | Serenoa repens |
| Serenoa repens | Syagrus romanzoffiana |
| Syagrus romanzoffiana | Trachycarpus fortunei |
| Trachycarpus sp: 10 sp/15 | Trithrinax campestris |
| Trithrinax acanthocoma | Washingtonia filifera & robusta |

TOTAL PA: 35 SPECIES

TOTAL RPW: 19 SPECIES

