



HAL
open science

Diversity and genetic structure in selfing populations in the absence of selection

Margaux Jullien, Miguel Navascués, Joelle Ronfort, Laurène Gay

► To cite this version:

Margaux Jullien, Miguel Navascués, Joelle Ronfort, Laurène Gay. Diversity and genetic structure in selfing populations in the absence of selection. Sex uncovered: the evolutionary biology of reproductive systems, Apr 2018, Roscoff, France. 1 p., 2018. hal-02786964

HAL Id: hal-02786964

<https://hal.inrae.fr/hal-02786964v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Margaux Jullien^a, Miguel Navascués^b, Joëlle Ronfort^a, Laurene Gay^a

^a INRA, UMR AGAP, GE²POP team, Montpellier, France

^b INRA, UMR CBGP, F-34988 Montferrier-sur-Lez, France

Selfing populations are characterized by a peculiar structure in which some multilocus genotypes reach a high frequency in the population while others are rare (Fig 1). Allard (1975) argued that this structure is a consequence of the combination of selfing and selection favoring locally adapted genotypes. However, genetic drift is expected to strongly affect selfing populations and it is unclear whether genetic drift alone could be responsible for the stochastic increase of one or a few genotypes in the population. Moreover, we lack analytical predictions for multilocus diversity under predominant selfing.

Can we explain the structure of predominantly selfing populations using neutral processes only? Are genotypes maintained over time? How is genetic diversity distributed in the populations?





Figure 1: Multilocus genotypes (MLG) network in a selfing population of *Medicago truncatula* sampled in two generations

Simulation framework



- Multilocus genotype (MLG) frequencies over time
- Single and multilocus diversity (H_e and Simpson's index, λ)
- Genetic distance between MLGs within generation (D , D_{max})

Results




Figure 2: MLG frequency spectra. The horizontal axis represents the frequency at which a MLG is found in the first sample (t_0), the vertical axis represents the frequency at which it is found in the second sample (t_20). The color gradient represents the \log_{10} of the frequency at which each case is observed in 1000 replicates of simulations.

- Selfing results in repeated MLGs that are maintained over time
- Only small population size or strong population bottlenecks lead to one predominant MLG
- Extinction-recolonization completely changes the MLG composition of the population over time
- For a fixed effective population size (corrected for the selfing rate⁽²⁾), monolocus diversity is similar but distributed between MLGs as shown by increasing D_{max} with selfing




Figure 3: Maximum distance between MLGs (D_{max}), single (H_e) and multilocus diversity (λ) in populations with increasing selfing rate but equivalent drift levels

Our simulations show that neutral processes combined with predominant selfing can result in populations composed of a few selfing “lineages” differentiated from one another, as observed in natural populations. Demographic events exacerbate the effects of selfing on MLG diversity and create temporal stochasticity of the population composition. This simulation framework and diversity statistics can be used to infer populations’ demographic parameters (Approximate Bayesian Computation, ongoing).

References

Allard, R. W. 1975. “The Mating System and Microevolution.” *Genetics* 79 Suppl (June): 115–26.