



Reactive transport modeling the oxalate-carbonate pathway of the Iroko tree

Hannah Gatz-Miller, Frederic Gerard, Eric Verrechia, Klaus Ulrich Mayer

► To cite this version:

Hannah Gatz-Miller, Frederic Gerard, Eric Verrechia, Klaus Ulrich Mayer. Reactive transport modeling the oxalate-carbonate pathway of the Iroko tree. Goldschmidt Conference, Aug 2019, Barcelone, Spain. hal-02787774

HAL Id: hal-02787774

<https://hal.inrae.fr/hal-02787774>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reactive transport modeling the oxalate-carbonate pathway of the Iroko tree

GATZ-MILLER, H.¹, GÉRARD, F.², VERRECCIA, E.³,
MAYER K. U.¹

¹ Department of Earth, Ocean, and Atmospheric Sciences,
University of British Columbia, Vancouver, Canada V6T 1Z4
(hgmiller@eos.ubc.ca; umayer@eos.ubc.ca)

² UMR Eco&Sols, INRA, Montpellier, France
(frederic.gerard@inra.fr)

³ Université de Lausanne, Lausanne, Switzerland
(eric.verrecchia@unil.ch)

Biologically facilitated calcite formation through the oxalate-carbonate pathway (OCP) of the Iroko tree (*Milicia excelsa*) has been previously studied as a method of natural carbon capture. Previous field and lab work in Côte d'Ivoire, Africa, has suggested that biomineralization via the OCP functions as calcium-oxalate containing plant tissue is dropped from the tree and the oxalate is oxidized by oxalotrophic bacteria, resulting in the alkalization of local tropical soil, so that calcite may precipitate. A reactive transport (RT) model of the OCP of the Iroko tree was constructed to examine and quantify these processes.

The RT model focused on quantitatively evaluating the fluxes and fate of calcium and carbon in the different components of the OCP process, to assess how much carbon could be stored as calcite in soils around the Iroko tree over and beyond its life cycle. Calcium input from dust, precipitation, bedrock weathering, and plant nutrient cycling, was combined with precipitation and transpiration rates, litter fall, decomposition and nutrient return, and the kinetics of the relevant oxidation-reduction reactions.

A plot-scale model representing a vertical soil profile was constructed using the RT code MIN3P-THCm and the input values described. Preliminary model results indicate that given sufficient oxalate influx over time, soil pH rises from an initial acidic value (5.1) to more alkaline (7-9). Even without calcite formation, this alkalinity increase has implications for sequestration of CO₂ within the soil atmosphere and solution as CO₂ and HCO₃⁻/CO₃²⁻, respectively. If sufficient Ca²⁺ is present, calcite precipitation occurs where the most calcium oxalate was present (near the surface), but does not occur where Ca²⁺ is limited, with implications for longer-term carbon sequestration processes.

Additional models are being constructed to assess the spatial extent and influence of the pH change, to be coupled with a root architecture model to investigate the overall role of the root system on biomineralization and the OCP of the Iroko tree.