



HAL
open science

Assessing the performance of landscape-based strategies to deploy major-gene resistance

Loup Rimbaud, Julien Papaix, Luke G. Barrett, J.J. Burdon, Peter H. Thrall

► **To cite this version:**

Loup Rimbaud, Julien Papaix, Luke G. Barrett, J.J. Burdon, Peter H. Thrall. Assessing the performance of landscape-based strategies to deploy major-gene resistance. Science Protecting Plant Health 2017, Australasian Plant Pathology Society.; Cooperative Research Centre in National Plant Biosecurity. AUS., Sep 2017, Brisbane, Australia. hal-02787792

HAL Id: hal-02787792

<https://hal.inrae.fr/hal-02787792>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessing the performance of landscape-based strategies to deploy major-gene resistance

Loup Rimbaud¹, Julien Papaix², Luke Barrett¹, Jeremy Burdon¹, Pete Thrall¹

¹CSIRO Agriculture & Food, Canberra, ACT 2601, Australia

²BioSP, INRA, 84914, Avignon, France.

loup.rimbaud@csiro.au

Genetically-controlled plant resistance can reduce the damage caused by pathogens. However, pathogens have the ability to evolve and overcome such resistance. This often occurs very quickly after resistance is deployed, resulting in significant crop losses and a continuing need to breed new resistant cultivars. To tackle this issue, several strategies have been proposed to constrain the evolutionary potential of pathogen populations and thus increase the durability of resistance deployment. These strategies mainly rely on using different combinations of resistance sources in time, space, or both. In time, such combination consists of crop rotations. In space, resistance sources can be deployed in the same cultivar (pyramiding), in different cultivars within the same field (cultivar mixtures) or in different fields (mosaics). However, experimental assessment of the efficiency (i.e. ability to reduce disease impact) and the durability (i.e. ability to limit pathogen evolution and delay resistance breakdown) of different deployment strategies presents a major challenge.

Therefore, we developed a spatially-explicit stochastic model to assess the epidemiological and evolutionary outcomes of the major deployment options described above when one or two major genes for resistance are present. In addition, we analysed the impact of landscape organisation (as defined by the proportion of fields cultivated with a resistant cultivar, and their spatial aggregation) and epidemiological or evolutionary parameters (e.g. dispersal abilities, mutation rate, cost of infectivity) through sensitivity analyses and polynomial regression.

The model has been parameterised for wheat qualitative resistance to rusts, caused by fungi of the genus *Puccinia*, but can be applied to many other pathosystems. Early results suggest that strategies offering the best epidemiological control of the disease are not necessarily the most durable.

Key words: deployment strategy, durable resistance, gene-for-gene resistance, plant disease, rust of wheat, spatially-explicit modelling