

Quantifying the impact of tree-diebacksand salvage logging on mountain forest biodiversity using metabarcoding

Lucas Sire, Rodolphe Rougerie, Laurent Larrieu, Christophe Bouget, Béatrice Courtial, Annie Bézier, Douglas W. Yu, Elisabeth Herniou, Carlos

Lopez-Vaamonde

► To cite this version:

Lucas Sire, Rodolphe Rougerie, Laurent Larrieu, Christophe Bouget, Béatrice Courtial, et al.. Quantifying the impact of tree-diebacksand salvage logging on mountain forest biodiversity using metabarcoding. 4. Colloque de Génomique Environnementale, Sep 2017, Marseille, France. 1 p., 2017. hal-02788069

HAL Id: hal-02788069 https://hal.inrae.fr/hal-02788069

Submitted on 5 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantifying the impact of tree-diebacks and salvage logging on mountain forest biodiversity using metabarcoding

Lucas SIRE¹, Rodolphe ROUGERIE², Christophe BOUGET³, Laurent LARRIEU⁴, Béatrice COURTIAL⁵, Annie BEZIER¹, Douglas YU⁶, Elisabeth HERNIOU¹, Carlos LOPEZ-VAAMONDE^{1,5}

: IRBLUMR CNRS 7261 – Université François Rabelais de Tours – Tours, France;²: Muséum national d'Histoire Naturelle, UMR 7205 ISYEB – Paris, France;³: Irstea, UR EFNO, Nogent-sur-Vernisson, France;⁴: INRA, UMR 1201 DYNAFOR & CNPF/CRPFOcc – Toulouse, France;⁵: INRA, UR0633 Zoologie Forestière – Orléans, France,⁶: Kunming Institute of Zoology, Yunnan, China.

> Background:

Climate change scenarii predict continued rising temperatures and an increase in the intensity and frequency of climate extremes. Extensive tree mortality (forest diebacks) are expected to become more widespread, frequent and severe. Tree diebacks is often followed by removal of dead trees by foresters (salvage logging).







> Metabarcoding & Bioinformatic pipeline:

mtDNA strand

Tagged primers per individual & PCR



> Main objective:

To measure the impact of climate induced silver fir forest diebacks and associated salvage logging on insect biodiversity using mass trapping and analysis of bulk samples with DNA (meta)barcoding.


> Methodology:

56 experimental plots have been set up in the in French Pyrenees silver firdominated moutain forests (Pays de Sault and Aure valley).

Environmental measurements:

- Local dieback assessment (ARCHI), tree defoliation index, mistletoe occurrence.
- Remote sensing data.
- Dendrometric measurements, stand structure, deadwood and tree-related microhabitats.

Biomonitoring using mass trapping:



- One Malaise trap and two interception traps per plot deployed from May to September 2017.
- Samples collected once a month (224 Malaise samples and 448 polytrapTM samples in total).



DNA barcoding reference libraries for French saproxylic beetles (PASSIFOR):


• 1571 specimens, 785 COI barcodes, 485 species, 53 families.

Metabarcoding:

- Development of a metabarcoding and standardized bioinformatical pipeline.
- Test of metabarcoding analysis from preservative ethanol

> Deliverables

- Comprehensive DNA barcode reference libraries for saproxylic beetles.
- An innovative workflow and laboratory protocol for biomonitoring forest biodiversity using metabarcoding.
- A complete list of species or species-level OTUs for each collecting event for biodiversity assessments.
- Measures of the difference in taxonomic, phylogenetic and functional diversity for invertebrates communities in forest plots differently impacted by climate change in two régions of the French Pyrenees.





Marie Lissndra Zepeda-Mendoza, Kristine Bohmann, Aldo Carmona Baez, M. Thomas P. Gilbert. 2016. « DAMe: a toolkit for the initial processing of datasets with PCR replicates of double-tagged amplicons for DNA metabarcoding analyses » *BMC Research Notes*. (9):255. DOI 10.1186/s13104-016-2064-9









