A mechanistic model to predict distribution of carbon among multiple sinks.

André Lacointe, Peter E.H. Minchin

► To cite this version:

André Lacointe, Peter E.H. Minchin. A mechanistic model to predict distribution of carbon among multiple sinks.. Phloem : Methods and Protocols, Editions Springer, 473 p., 2019, Methods in Molecular Biology, 978-1-4939-9561-5. hal-02788117

HAL Id: hal-02788117

<https://hal.inrae.fr/hal-02788117>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A mechanistic model to predict distribution of carbon among multiple sinks.

André Lacointe⁽¹⁾, Peter E.H. Minchin⁽²⁾

(1) Université Clermont Auvergne, INRA, PIAF, F-63000, Clermont-Ferrand, France; email: andre.lacointe@inra.fr

(2) New Zealand Institute for Plant and Food Research, Motueka Research Centre, Motueka, New Zealand; email: Peter.Minchin@plantandfood.co.nz

Abstract/summary.

Modelling is a fundamental part of quantitative science. It is a methodology of the holistic approach of bringing together quantitative ideas, many of which will have been developed though a reductionist approach that allows a lot of detail to be gathered on a small part of the system of interest. Phloem and xylem physiology are both descriptions of whole plant behaviour. The phloem is especially difficult to study in a reductionist way because as soon as the phloem is disturbed, even very carefully, it stops functioning by induction of blockage and other defensive mechanisms. This was the cause of a long debate on the basic structure of the phloem's long-distance transport pathway. Were the sieve-tubes 'blocked' at the sieve-plates or was there a continuous open conduit between source and sink? Developments in very rapid chilling of small pieces of phloem tissue, to obtain the required speed of cooling, was needed before reliable micrographs could be obtained and conclusively showed that the observed sieve-plate blockages were an artefact brought about by phloem damage quickly leading to blockage mechanisms, believed to be needed to prevent loss of significant phloem sap when plants are damaged

It is now generally accepted that phloem flow is the result of bulk solution flow generated by osmotic pressure generated by phloem loading. But there is still little agreement on how sink competition functions and the well documented source sink relations observed with tracer studies. More recently the importance of phloem pathway leakage (unloading) and reloading

has been recognised and the role of this is still being unravelled. Interactions between phloem and xylem flows are now thought to be important, and may have a role in carbohydrate source-sink relations through potassium recirculation.

All of these areas are extremely difficult to research by the reductionist approach, with modelling being an important tool to test the consequences of proposed mechanisms which can then be tested in whole plant experiments.

Phloem/xylem modelling has been at the limits of quantitative modelling, especially when dynamic models are needed to explain tracer studies. Huge advances in computing now enable more realistic modelling and the PiafMunch approach has extended that even further by enabling much more mechanistic detail to be incorporated. With the recent introduction of tracer dynamics now incorporated in PiafMunch it will be possible to look at the effects of specific phloem mechanisms upon the shape of evolving tracer profiles.

Keywords: Münch model, carbon allocation, sink priority, phloem, xylem, coupled water and carbon fluxes, plant architecture, functional - structural plant modelling, source sink relations

Introduction

Wardlaw (1990) reviewed a large body of experimental data on carbon partitioning in plants and found no mechanistic understanding of the data. More recently Lacoïnte (2000) reviewed the range of models used in functional-structural tree models where he reviewed the empirical methods based on allometry, sink priorities and functional equilibrium, but found no mechanistic approaches to this fundamental aspect of balanced plant growth.

Currently, the general consensus of phloem flow is that proposed by Münch (1928) with bulk flow of phloem sap driven by an osmotically generated pressure gradient created by loading of photosynthate (usually sucrose) at the source and unloading at the sink. In many plant species phloem loading is an active transport process across the sieve-tube plasmalemma resulting in a source solute concentration in the order of 0.8 M while in other species this is a passive process relying on diffusional flow from the cells associated with photosynthesis.

Detailed biophysical models of this process were first described by Christy and Ferrier (1973) and this work has had considerable complexity added resulting in the recent work of Thompson and Holbrook (2003). This work describes a single-source single-sink system with concomitant water flow in terms of parameters describing phloem loading, phloem unloading, and includes lateral water flow through an ideal (*i.e.* reflection coefficient for the solute is one) sieve-tube plasmalemma.

The first attempt to extend this approach to multiple sinks supplied by a single source was that of Minchin *et al.* (1993). Their simple 2-sink model was able to mimic several observed phenomena involving shoot-root interactions and gave the first quantitative explanation of sink priority. This model predicted that the relative sink priorities between the shoot and root

of a barley seedling could be reversed by cooling the root, and this was subsequently demonstrated (Minchin *et al.*, 1994). This preliminary model was based on a non-permeable, either by water or solute, long-distance transport pathway which is known not to be the case. This simplification greatly simplified the model equations allowing them to be solved analytically. Bidel *et al.* (2000) expanded this approach to many sinks representing a growing root and used an iterative approach to determine how carbohydrate flows were able to mimic different patterns of root growth by altering the individual sink properties.

But it is well known that the plant vasculature consists of both phloem and xylem which are physically close and readily interact through both water relations and controlled transfer of solutes. Daudet *et al.* (2002) incorporated xylem/phloem interactions through water relations which could now incorporate effects of transpiration-induced gradients of water potential. Local gradients of all water- and carbon-flux related variables could be accounted for by a spatially discretized approach, which turned partial differential to ordinary differential equations. They used P-SpiceTM software to illustrate their methods on a branching system with three source leaves, and three competing fruits. This work has been extended (Lacointe and Minchin, 2008) to allow huge flexibility in architecture and specific mechanistic detail through use of recently developed numerical methods, resulting in the model 'PiafMunch'. Recently, Hall and Minchin (2013) proposed a closed-form solution for steady-state coupled phloem/xylem using the Lambert-W function, which can handle multiple sinks. While incorporating some of the added complexities, such as variation of phloem resistance with solute concentration, and deviations in the Van't Hoff expression for osmotic pressure, the differential equation approach is still quite limited in its ability to handle a lot of detail of physiological interest (*e.g.* pathway unloading/reloading of solute, different unloading

kinetics). By contrast, a major advantage of the PiafMunch approach is its flexibility to be able to work with a huge range of local loading and unloading mechanisms.

In this chapter, we will first describe the original PiafMunch model as published in 2008 in detail. Then its capacities will be illustrated by examples of use and results. The third part will introduce recent and current developments regarding (i) a more general description of the plant architecture, and (ii) inclusion of additional, refined biophysical or metabolic processes. Finally, practical details will be given to help potential users to handle the model efficiently.

PiafMunch -- the original model (Lacointe and Minchin, 2008)

As a functional-structural model, PiafMunch includes both an architectural description of the plant structure and a mechanistic description of relevant biophysical processes at local level.

Discretisation of the plant structure

The plant skeleton is discretized into an arbitrary number of segments delimited by junction nodes¹ (Fig. 1). The plant architecture is thus represented as a collection of elements, each consisting of a topological node and an associated axial pathway segment, with the exception of the ‘collar’ node, whose physical connecting upward and downward pathway segments are conventionally assigned respectively to its upper stem and lower root elements. Most elements are connected to one upper and one lower element, except for the ends of roots where there is no ‘lower’ element, the ends of stems where there is no ‘upper’ element, and branching elements which have a single connection at one end and two at the other. Thus, a total of seven different element types (denoted by different colours in Fig. 1) are required and sufficient to describe any branched architecture.

¹the term ‘node’ here is being used in the topological sense, without reference to botanical nodes which bear the leaves on a plant shoot. In future we will simply use the term node.

Local hydraulic architecture

Each axial pathway segment includes one phloem and one xylem pathway, which are connected to each other by a transverse pathway (Fig.2) allowing for lateral water exchanges between the sieve tube and local apoplasm. At end nodes, water exchanges with external environment are represented, either as imposed local fluxes (*e.g.* measured transpiration rates), or constrained by outside (*e.g.* soil) local water potential. Those represent the system boundary conditions, which are allowed to fluctuate.

Hydraulic fluxes

According to the accepted Münch theory (1928), viscous flow of phloem sap is driven by an axial hydraulic pressure gradient generated by active loading of solutes at the source and unloading in the sink. Lateral solute leakage with reloading occurs along the long-distance pathway, as does lateral water flow determined by water potential gradients and sieve-tube membrane water permeability.

Xylem flow is driven by the axial apoplastic pressure gradient generated by leaf transpiration.

These basic principles are expressed for each element as a set of equations involving its own local variables and parameters. The volume fluxes of xylem and phloem water are respectively

$$JW_{xyl} = \Delta P_{xyl}/r_{xyl} \quad \text{xylem flow between connected elements} \quad (1)$$

$$JW_{st} = \Delta P_{st}/r_{st} \quad \text{phloem flow between connected elements} \quad (2)$$

The phloem sieve tube resistance r_{st} can be either entered as a local parameter or estimated from sieve tube geometry and sap viscosity (Thompson and Holbrook 2003 – see chapters 22 and 26 in this book). Sap viscosity is dependent on temperature and solute concentration,

which is empirically described by an exponential to within 1% of experimental values over a wide range of temperatures and concentrations (Gilli 1997, after Mathlouthi and Génotelle 1995).

Lateral water flow from xylem to phloem sieve tube is driven by the difference in water potential:

$$JW_{lat} = (\Psi_{xyl} - \Psi_{ST})/r_{lat} \quad (3)$$

where r_{lat} is the sum of the apoplastic pathway resistance between xylem and phloem and the sieve-tube cross-membrane resistance, which is inversely proportional to the membrane permeability.

Taking into account the non-zero partial molal volume of sucrose $\bar{V}$ adds an extra lateral component NZS to the volume flow into the sieve tubes:

$$JW_{lat} = NZS + (\Psi_{xyl} - \Psi_{ST})/r_{lat} \quad (3')$$

$$NZS = \bar{V} \cdot JS_{lat} \quad (3'')$$

where JS_{lat} is the lateral solute flow (see next section).

Hydrostatic pressure P_{ST} within the sieve tubes is given by the difference between total phloem water potential and osmotic potential inside sieve tubes:

$$P_{ST} = \Psi_{ST} - \Pi_{ST} \quad (4)$$

Xylem sap has a very low solute concentration which we shall ignore, so there is no osmotic component to its total water potential:

$$P_{xyl} = \Psi_{xyl} \quad (4')$$

For a single phloem solute, Π_{ST} is determined by its concentration C_{ST} . For a dilute solution, Π_{ST} is given by the Van't Hoff relation:

$$\Pi_{ST} = -R T C_{ST} \quad (5)$$

where R is the universal gas constant and T the absolute temperature. For a non-dilute solution we use the empirical equation stated by Thompson and Holbrook (2003):

$$\Pi_{ST} = -\rho_w R T (0.998 m + 0.089 m^2) \quad (5')$$

with ρ_w the density of water and m the molality given by

$$m = C_{ST} / [\rho_w (1 - C_{ST} \cdot \bar{V})] \quad (5'')$$

If the partial molal volume of sucrose $\bar{V}$ is taken as zero then equations 5' and 5'' reduces to the Van't Hoff relation (5), and equation (3') reduces to the Ohm's law analog (3). When the solute is sucrose and the concentration is 1 mol L⁻¹ (typical at the site of phloem loading) then using the Van't Hoff relation for a dilute solution results in about an 8% error in Π_{ST} , while at 0.5 mol L⁻¹ sucrose this reduces to a 2% error. As this is low enough in most situations (compared to other error sources *e.g.* in the model parameter values), this refinement can be deactivated by the user to reduce computation time.

The set of water-flow equations for each element is completed by a flow conservation statement for each of the two hydraulic pathways within an element, one for the xylem and one for the phloem (Fig. 2):

$$\sum_k JW_k = 0 \quad (6)$$

where JW_k (with appropriate sign) represents the lateral and all longitudinal liquid flows to/from the node k , the number of which depends on the element type as defined above. In particular, xylem flow for terminal elements includes transpiration at stem ends ('leaves') and water uptake from soil at the root tips (Fig. 2). Note that eq. (6) assumes unchanging sieve tube volume (rigid sieve-tube cell walls), which Thompson and Holbrook (2003) showed to be an acceptable approximation in many situations.

Solute flows

Longitudinal phloem solute flow between two connected elements is given by:

$$JS_{ST} = JW_{ST} C_{ST} \quad (7)$$

where C_{ST} is the sieve-tube solute concentration in the upflow element.

Variation of sieve-tube solute content $Q_{ST} (= C_{ST} \cdot V_{ST})$ is:

$$\frac{dQ_{ST}}{dt} = JS_{lat} + \sum_k JS_{_k} \quad (8)$$

where $JS_{_k}$ (with appropriate sign) represents all longitudinal solute flows from/to the connected element(s), and JS_{lat} is the lateral solute flow into the sieve-tube.

The lateral solute flow rate JS_{lat} , *i.e.* the local unloading/reloading, can be either set directly as an independent equation or derived from local metabolism (*e.g.* respiration, photosynthesis or starch $\leftrightarrow$ soluble sugar conversion) occurring in an attached parenchyma compartment. This is up to users who can write their own set of equations for JS_{lat} which can have any form, including ordinary differential equations. However, a predefined set of classical equations is proposed for convenience :

$$JS_{lat} = k_1 \cdot (C_{Par} - C_{ST}) \cdot V_{ST} + (k_2 C_{Par} + k_3) V_{Par} \quad (9)$$

where C_{Par} is the parenchyma solute concentration, V_{ST} the sieve-tube volume and V_{Par} the parenchyma volume. This allows a number of different dynamics by assigning specific values to local parameters k_1, k_2, k_3 , *e.g.*:

- diffusion-like kinetics ($k_2 = k_3 = 0$);
- constant loading/unloading ($k_1 = k_2 = 0$);
- concentration-dependent loading with a target concentration C_{targ}

$$(k_2 = -k_1 V/V_{Par}, \quad k_3 = k_1 C_{targ} V_{ST}/V_{Par});$$

- concentration-dependent unloading as in Thompson and Holbrook (2003)

$$(k_3 = 0, k_2 = -k_1 V_{ST}/V_{Par})$$

such that simple cases of symplastic loading/unloading are currently built into equation (9).

The default equation for parenchyma solute content $Q_{Par} (= C_{Par} \cdot V_{Par})$ change rate simulates

the result of sucrose exchange with local sieve tubes (JS_{lat} , eq. 9), exchange with

environment (maintenance respiration and/or photosynthesis) and starch/sucrose

interconversion:

$$\frac{dQ_{Par}}{dt} = -JS_{lat} - R_M + Ph - \frac{dS}{dt} \quad (10)$$

where respiration R_M , photosynthesis Ph , and starch S are all expressed in sucrose equivalents.

The dynamics of photosynthesis Ph may be either read from an external file or modelled by

the user *e.g.* as a periodic function (Daudet *et al.*, 2002). For maintenance respiration R_M , the

proposed formalism is that of Thornley (1970; see review by Le Roux *et al.*, 2001), with a

concentration-dependent maintenance coefficient to account for phloem sucrose leakage /

active reloading:

$$R_M = (k_4 + k_5 C_{ST}) Sr \quad (11)$$

where Sr is the structural carbon content of the element biomass, expressed in sucrose

equivalents.

The default representation of starch metabolism uses a general equation derived from Daudet

et al. (2002):

$$\frac{dS}{dt} = \frac{v_{max} \cdot C_{Par}}{k_M + C_{Par}} \cdot V_{Par} - k_{hyd} \cdot S + k_6 \cdot (C_{Par} - C_{targ}) \cdot V_{Par} \quad (12)$$

allowing simulation of a number of dynamics, *e.g.* Michaelis-Menten kinetics for synthesis

from a sucrose substrate (through parameters v_{max} and k_M), starch content-dependent

1 hydrolysis back to sucrose (parameter k_{hyd}) or sucrose concentration-dependent
2 interconversion with a target concentration C_{targ} .

3
4 All solute-fluxes equations are fully editable, with a possible redefinition of all predefined
5 parameters or definition of new ones, allowing *e.g.* very simple configurations like Minchin *et*
6 *al.* (1993) or Thompson and Holbrook (2003) which were used to test the model (Table 2). If
7 edited, it is up to the user to make sure that the equations make sense. By contrast, all water-
8 fluxes equations, which are the heart of the model, are hard coded except for parameters. The
9 full system is coded in C++ and solved by a combination of LAPACK linear algebra package
10 (Anderson *et al.*, 1999) with the sparse extension TAUCS (Toledo, 2003) and SUNDIALS
11 algebraic/differential equation solver package (Hindmarsh *et al.*, 2005). The software includes
12 a graphic user interface to specify the architecture, parameters, initial values and other
13 settings. This allows use of the software, with the default equations, without having to
14 recompile it. More flexibility can be achieved by editing the solute fluxes equations and
15 recompiling.

17 **Examples of what the PiafMunch model has been used on**

18 The first application of the PiafMunch software was to a single-source single-sink linked by a
19 5m long distance pathway consisting of a tube of 7.5 μm diameter conduit with membrane
20 permeable to water but not to the solute (sucrose), *i.e.* a ‘perfect’ semipermeable membrane,
21 with uniform loading along the first 0.5 m and unloading along the last 0.5 m. This example
22 was chosen for direct comparison with the work of Thompson and Holbrook (2003) using a
23 continuous differential equation framework. With the PiafMunch approach we started with
24 $N=3160$ elements and then looked at the effect of reducing N to a much smaller number.
25 With the large value of N the phloem water flux was very similar to that calculated by the

continuum approach of Thompson and Holbrook (2003), with the greatest differences being about 2% when the fluxes were changing at the greatest rate (see Lacoïnte and Minchin 2008, Fig.3). That low 2% discrepancy can be ascribed to the variation of lumen diameter with pressure, which was included in Thompson and Holbrook (2003) but ignored in PiafMunch. With much lower values of the element number N (*e.g.* $N = 30$), the two approaches differed most at the sites of most rapid flux change with differences ca. 10%, and were very close in the regions along the long-distance pathway where the water flux was not rapidly changing. When these comparisons were made 24 hr into the simulation when the flows had reached equilibrium, there was only a small difference between the two approaches, even with N as low as 10, with a maximum deviation below 10%. From this it was concluded that the discrete approach of the PiafMunch model resulted in similar results to the continuous method of Thompson and Holbrook (2003), and that the number of discrete elements required for a good approximation of the continuous system does not need to be very high.

The variation of sap viscosity with sugar concentration was first introduced by Bancal and Soltani (2002) in a simplified Münch model, assuming uniform concentration along the pathway and ignoring membrane permeability. A few years later, Hölttä *et al.* (2006) introduced the viscosity change in a more realistic model but ignoring the specific partial molal volume of sugar. In both studies, the authors concluded that high sugar loading rates could block phloem transport due to high sap viscosity. PiafMunch incorporates that variation in sap viscosity with solute concentration as well as the partial molal volume of the solute; these refinements can be set on/off by a simple click, either individually or both together. It was confirmed that both of these had significant effect, both on the equilibrium and non-equilibrium flux. Work is needed to determine if these effects have any physiological significance. Hölttä *et al.* (2006) also showed that transpiration rate can be expected to

1 interact with phloem flow, *via* water relations described by equation 3 above. This was also
2 demonstrated in the initial PiafMunch work (Lacointe and Minchin 2008, Fig. 8).

3
4 The PiafMunch model is meant to handle more complex source-sink configurations. Working
5 with a considerably simplified model with a solute molal volume of zero, constant sap
6 viscosity with changes in solute content, and a non-permeable long-distance transport conduit,
7 Minchin *et al.* (1993) developed a Munch model describing flow between 1-source and 2-
8 sinks which predicted changes in the proportion of total solute flow delivered to each sink
9 when the source supply changed, which in experimental work has been described in terms of
10 sink priority. This was the first mechanistic description of sink priority. Working with the
11 same source sink configuration, and incorporating non-zero pathway permeability the
12 PiafMunch model showed the similar priority behaviour. The main purpose of this work was
13 to determine if the PiafMunch model gave results consistent with previous work, and it passed
14 this test with flying colours so we can have confidence in this new approach and now
15 investigate examples the previous methods cannot handle.

16
17 Thorpe *et al.* (2011) went on to model a 2-source 3-sink configuration generated in a heavily
18 pruned dwarf bean plant and test the predictions using ^{11}C tracer. Several observed treatment
19 responses were successfully predicted, but the observations could not be completely explained
20 when the modelled common pathway, comprising the stem, contained just one phloem
21 pathway. Bidirectional flow within the stem was necessary to explain the observed flows.
22 It is now accepted that the long-distance phloem transport pathway is leaky, and also takes up
23 solute from the immediate apoplast. This manifests itself in tracer studies through the
24 observed buffering of phloem flow when the phloem pathway is disturbed or there are sudden
25 changes in source or sink function. The PiafMunch model has been used to determine if this

leakage/reloading alters source sink dynamics (Minchin and Lacointe, 2017). This modelling indicated that the phloem flow does not follow Poiseuille dynamics, i.e. the water flux was not proportional to ΔP , due to there always being water flow across the membrane, even without pathway unloading and or reloading of solute. At equilibrium, the presence of unloading altered the solute concentration and hydrostatic pressure profiles. With adequate reloading along the pathway the effects of pathway unloading were completely compensated for, making the equilibrium system look like one with no pathway unloading. Further work is needed here to look at the non-equilibrium flows. To do this even more model parameters are needed, though this might be a means of estimating these parameter values through optimising the parameters to produce behaviour similar to that seen in plant experiments.

Current developments

A new version has been developed (PiafMunch v.2) which features a few major improvements:

(1) The extension of the architectural pattern from branched to any network

architecture, including loops and nodes of any connectivity order (that was limited to 3 in v.1, *i.e.* each node could be connected to 3 other nodes at most). That significantly extends the scope of the model, to *e.g.* the looped nervation pattern of an isolated leaf, or non-binary, verticillate branching patterns as exhibited by conifers.

(2) A significant refinement of the local tissue model at the node level (Fig. 3 and 4),

with explicit apoplastic, possibly solute-containing, compartments attached to both phloem (in addition to the sieve tube) and lateral parenchyma (in addition to symplasm). In particular, the lateral pathway from xylem vessel to phloem sieve tube (r_{lat} , JW_{lat} in eq. 3) is now explicitly segmented into an apoplastic pathway (r_{Trsv} , JW_{Trsv}) and the cross-membrane pathway from phloem apoplasm to sieve tube (r_{PhIMb} ,

JW_{PhlMb}). Water and solute fluxes between adjacent compartments are described by equations similar to (1-9) above, allowing more realistic simulation of *e.g.* apoplastic (un)loading and related cross-membrane processes, or a convective component of symplasmic (un)loading.. Again, all parameters and solute flux equations can be redefined or edited to reduce the extended model to v.1 (Lacointe and Minchin 2008), or the simpler Minchin *et al.* (1993).

(3) **User-defined sharp parameter changes** can be implemented at specific time points, in addition to continuous parameter changes which were already possible in v.1 (with concentration-dependent phloem viscosity as a built-in example). This allows simulation of *e.g.* cold blocking / unblocking of the phloem pathway or aquaporin function - related changes in cross-membrane resistances (Steppe *et al.*, 2012).

(4) **Lateral parenchyma symplasmic volumes are now considered variables** that can be driven by differential equations involving any other variables like local pressure, allowing simulation of *e.g.* reversible, elastic volume changes, *i.e.* water capacitance, involved in reversible stem diameter changes (Steppe *et al.*, 2012), or plastic, irreversible growth (Lockhart, 1965; Daudet *et al.*, 2002).

(5) **Tracer analysis facilities** have been included as a helper to design and analyse results from tracer experiments like that of Thorpe *et al.* (2011). They involve equations similar to the solute-related equations with additional terms for radioactive decay, in particular for ^{11}C which has a very short half-life of *ca.* 20 min.

Concluding remarks

PiafMunch has proved highly efficient and reliable on simple systems -- even though they could be more complex than those modelled by other approaches in literature. However, it is meant to handle truly complex architectures and detailed local processes, using highly

efficient, state-of-the-art numerical methods. It can be used to simulate and test effects of any known or hypothetical mechanism, both at the local and at the global, plant-wide scale level. It can be used also to simulate coupled solute/water relations of a single, isolated organ, *e.g.* an isolated leaf, given appropriate boundary conditions dynamics. Furthermore, it can be easily extended, as shown by the development of v.2. This could be readily further extended to *e.g.* a vacuolar compartment with specific aquaporins mediating tonoplast resistances; another possible extension, though slightly more difficult to implement, would be to introduce multiple phloem pathways as suggested by results from Thorpe *et al* (2011). It should be emphasized, however, that any extension involves additional parameters (theoretically N additional values for each additional local parameter, although it is often reasonable to assume a common, single value), which can result in unreasonable, and confusing, complexity. This can be handled by setting unnecessary parameters to zero or infinite values to keep focus on those relevant to the issue of interest. Because of the multiplicity of parameters, it is unrealistic to use the model as a means to optimise parameters to fit experimental data; however, this could be made possible by setting a significant subset of parameters to known, reasonable values and optimising only another, limited, focus subset of parameters. This emphasizes the importance of membrane-focused and other experimental studies at the tissue or cell level to provide such required parameter values. As for all modelling, its use is in testing ideas and never a substitute to experiment.

Potential users interested in implementing PiafMunch in their work are very welcome to contact the authors. We will be pleased to help, either for specific applications or for practical installation details.

Table 1. Variables involved in the model PiafMunch v.1 (2008)

Symbol	Meaning	unit
C_{ST}, C_{Par}	sucrose molar concentration (sieve tube, parenchyma)	mmol mL ⁻¹
JS_{lat}	lateral sucrose flux	mmol h ⁻¹
$JS_{ST}^{(1)}$	longitudinal phloem sucrose flow between adjacent elements	mmol h ⁻¹
JS_k	longitudinal sucrose flow between current node and node #k	mmol h ⁻¹
JW_{lat}	transverse water flux	mL h ⁻¹
JW_k	lateral or longitudinal solution flow	
	between current node and node #k	mL h ⁻¹
$JW_{Xyl}, JW_{ST}^{(1)}$	longitudinal water flux (xylem, sieve tube)	mL h ⁻¹
m	solution molality	mol(kg of water) ⁻¹
P_{ST}, P_{Xyl}	hydrostatic pressure (sieve tube, xylem)	MPa
Q_{ST}, Q_{Par}	quantity of sucrose (sieve tube, parenchyma)	mmol
R_M	maintenance respiration	mmol _{sucrose} h ⁻¹
Sr	structural carbon content	mmol _{sucrose} equivalent
S	starch content	mmol _{sucrose} equivalent
V_{ST}, V_{Par}	volume (sieve tube, parenchyma)	mL
NZS	non-zero sugar volume flow accompanying JS_{lat}	mL h ⁻¹
Π_{ST}	osmotic potential	MPa
Ψ_{Xyl}, Ψ_{ST}	water potential (xylem, sieve tube)	MPa

⁽¹⁾ JS_{ST}, JW_{ST} were called resp. JS_{Phl}, JW_{Phl} in Lacointe and Minchin 2008

1 **Table 2.** Model parameters

<u>Symbol</u>	<u>Meaning (equation. involved)</u>	<u>value as used to simulate Thompson & Holbrook 2003</u>
C_{targ}	target sucrose concentration for starch metabolism (eq. 12)	$0.1 \text{ mmol mL}^{-1} \text{ }^{(1)}$
k_1	lateral carbon flow rate parameters (eq. 9)	3.23994 h^{-1} in unloading zone ; 0 elsewhere
k_2		-3.23994 h^{-1} in unloading zone ; 0 elsewhere
k_3		$0 \text{ mmol mL}^{-1} \text{ h}^{-1}$
k_4	Maintenance respiration – related parameters (eq. 11)	0 h^{-1}
k_5		$0 \text{ mL mmol}^{-1} \text{ h}^{-1}$
k_6	Starch metabolism – related parameter (eq. 12)	0 h^{-1}
k_{hyd}	relative rate of starch hydrolysis (eq. 12)	0 h^{-1}
k_M	Michaelis-Menten constant for starch synthesis (eq. 12)	$0 \text{ mmol}_{sucrose} \text{ mL}^{-1} \text{ }^{(1)}$
Ph	rate of photosynthesis (eq. 10)	$0 \text{ mmol}_{sucrose} \text{ h}^{-1} \text{ }^{(2)}$
R	universal gas constant (eqs. 5, 5')	$0.0083143 \text{ MPa mL K}^{-1} \text{ mmol}^{-1}$
r_{lat}	lateral hydraulic flow resistance (eq. 3, 3')	$23.5785 \times N^{(3)} \text{ MPa h mL}^{-1}$
$r_{ST}^{(4)}$	axial phloem sieve tube hydraulic flow resistance (eq. 2)	$14050.3 / N^{(3)} \text{ MPa h mL}^{-1} \text{ for } C_{ST} = 0.5 \text{ mmol mL}^{-1} \text{ }^{(4)}$
r_{Xyl}	axial xylem hydraulic flow resistance (eq. 1)	$10^{-200} / N^{(3)} \text{ MPa h mL}^{-1}$

T	absolute temperature (eqs. 5, 5')	293 K
$\bar{V}$	partial molal volume of sucrose (eqs. 3', 5'')	0.2155 mL mmol ⁻¹
v_{max}	kinetic parameter for starch synthesis (eq. 12)	0 mmol _{sucrose equ.} mL ⁻¹ h ⁻¹
ρ_w	density of pure water (eqs. 5', 5'')	0.99803 kg L ⁻¹ at $T = 293$ K

- 1 ⁽¹⁾ software default value, without effect in simulating either Thompson & Holbrook (2003) or Minchin *et al* (1993).
- 2 ⁽²⁾ user-defined input variable (does not have to be constant).
- 3 ⁽³⁾ axial (resp. lateral) hydraulic resistances are proportional (resp. inverse proportional) to element length, hence the scaling by N (see text).
- 4 ⁽⁴⁾ r_{ST} (called r_{Phl} in Lacointe and Minchin 2008) changes with C_{ST} , in proportion to the solution viscosity.

References

- Anderson E, Bai Z, Bischof C, Blackford S, Demmel J, et al. (1999). 'LAPACK users' guide. 3rd edn. <http://www.netlib.org/lapack/lug/index.html> (accessed 28 March 2018).
- Bancal P, Soltani F (2002). Source-sink partitioning. Do we need Münch? *Journal of Experimental Botany* **53**, 1919-1928.
- Bidel LPR, Pages L, Riviere LM, Pelloux G Lorendeau JY (2000). Mass Flow Dyn 1: A carbon transport and partitioning model for root system architecture. *Annals of Botany* **85**, 869-886.
- Christy A.L, Ferrier JM (1973). A mathematical treatment of Münch's pressure-flow hypothesis of phloem translocation. *Plant Physiology* **52**, 531-538.
- Daudet FA, Lacoite A, Gaudillère JP, Cruiziat P (2002). Generalized Münch coupling between sugar and water fluxes for modelling carbon allocation as affected by water status. *Journal of Theoretical Biology* **214**, 481-498.
- Gilli R (1997). Evaluation de différentes données physico-chimiques relatives aux solutions sucrées. <http://www.associationavh.com/fr/feuilles.html> (accessed 28 March 2018).
- Hall AJ, Minchin PEH (2013). A closed-form solution for steady-state coupled phloem/xylem flow using the Lambert-W function. *Plant, Cell and Environment* **36**, 2150-2162.

- 1 Hindmarsh AC, Brown PN, Grant KE, Lee SL, Serban R, Shumaker DE, Woodward CS
2 (2005). SUNDIALS: Suite of Nonlinear and Differential/Algebraic Equation Solvers.
3 *ACM Transactions on Mathematical Software* **31**, 363-396.
4
- 5 Hölltä T, Vesala T, Sevanto S, Perämäki M, Nikinmaa E (2006). Modeling xylem and phloem
6 water flows in trees according to cohesion theory and Münch hypothesis. *Trees* **20**,
7 67-78.
8
- 9 Lacoite A (2000). Carbon allocation among tree organs. A review of basic processes and
10 representation in functional tree-models. *Annals of Forest Science* **57**, 521-533.
11
- 12 Lacoite A, Minchin PEH (2008). Modelling phloem and xylem transport within a complex
13 architecture. *Functional Plant Biology* **35**, 772-780. <https://doi.org/10.1071/FP08085>
14
- 15 Le Roux X, Lacoite A, Escobar-Gutiérrez A, Le Dizès S (2001). Carbon-based models of
16 individual tree growth: a critical appraisal. *Annals of Forest Science* **58**, 469-506.
17
- 18 Mathlouthi M, Génotelle J (1995). Rheological properties of sucrose solutions and
19 suspensions. *In: 'Sucrose. Properties and Applications'*. (Ed. M.Mathlouthi and P.
20 Reiser) pp. 126-154. (Blackie Academic & Professional)
21
- 22 Minchin PEH, Thorpe MR, Farrar JF (1993). A simple mechanistic model of phloem
23 transport which explains sink priority. *Journal of Experimental Botany* **44**, 947-955.
24

- 1 Minchin PEH, Farrar JF, Thorpe MR (1994). Partitioning of carbon in split roots of barley:
2 effect of temperature of the root. *Journal of Experimental Botany* **45**, 1103-1109.
3
- 4 Minchin PEH, Lacointe A (2017). Consequences of phloem pathway unloading/reloading on
5 equilibrium flows between source and sink: a modelling approach. *Functional Plant*
6 *Biology* **44**, 507-514.
7
- 8 Münch E (1928). Versuche über den Saftkreislauf. *Deutsche botanische Gesellschaft* **45**, 340-
9 356.
10
- 11 Steppe K, Cochard H, Lacointe A, Ameglio T (2012). Could rapid diameter changes be
12 facilitated by a variable hydraulic conductance? *Plant, Cell and Environment*, **35** (1),
13 150-157.
14
- 15 Thompson MV, Holbrook NM (2003). Application of a single-solute non-steady-state phloem
16 model to the study of long-distance assimilate transport. *Journal of Theoretical*
17 *Biology* **220**, 419-455.
18
- 19 Thornley JMH (1970). Respiration, growth and maintenance in plants. *Nature* **227**, 304–305.
20
- 21 Thorpe MR, Lacointe A, Minchin PEH (2011).Modelling phloem transport with a pruned
22 dwarf bean: a 2-source- 3-sink system. *Functional Plant Biology* **38**, 127-138.
23
- 24 Toledo S (2003). TAUCS, a library of sparse linear solvers.
25 <http://www.tau.ac.il/~stoledo/taucs> (accessed on 28 March 2018)

1

2 Wardlaw IF (1990). The control of carbon partitioning in plants. *New Phytologist* **116**, 341-

3 381.

Figure captions

Figure 1. Representation of plant architecture. In this example, a plant (A) with 2 roots and 3 branches each terminated by a leaf, is represented by 18 elements (B) : one collar element (blue, No. 1), six non-branched stem elements (brown; No. 2, 3, 4, 5, 6, 7), three shoot terminations (green; No. 8, 9, 10), two branched stem elements (orange; No. 11, 12), 1 branched root element (black; No. 13), three non-branched root elements (dark grey; No. 14, 15, 16) and two root tips (light grey; No. 17, 18). Reproduced from Lacoïnte *et al.* (2008) with permission from CSIRO Publishing.

Figure 2. Network of hydraulic pathways for the example presented in Fig.1. The colour coding is the same as in Fig. 1. Reproduced from Lacoïnte *et al.* (2008) with permission from CSIRO Publishing.

Figure 3. The PiafMunch v.2. extended volume flow model. $r_{...}$, $JW_{...}$, $P_{...}$, $\Psi_{...}$, $\Pi_{...}$, $V_{...}$: resp., hydraulic resistance, volume flux, turgor pressure, water potential, osmotic potential, volume; for resp. (subscripts): ST , Xyl , $Trsv$, Apo , $PhlMb$, $ParMb$, $ParApo$, $Sympl$: sieve tube, xylem, transverse (xylem to phloem) apoplasm, pathway, phloem to lateral parenchyma apoplasm, pathway, sieve tube plasmalemma, lateral parenchyma plasmalemma, lateral parenchyma apoplasm, lateral parenchyma symplasm, NZS , NZS_{Par} : non-zero sugar volume flow, resp. into sieve tube and parenchyma symplasm. Operator Δ means '[in] – [out]' when applied to node-to-node-connector variables (JW_{ST} , JW_{Xyl}), or '[upflow] – [downflow]' when applied to node variables (P_{ST} , P_{Xyl}).

Figure 4. The PiafMunch v.2. extended solute flow model. $JS_{...}$, $Q_{...}$, $C_{...}$, $V_{...}$: resp., solute

- 1 flux, total solute content, solute concentration, volume; for resp. (subscripts): *ST, Apo, PhlMb,*
- 2 *ParMb, ParApo, Sympl* : sieve tube, phloem to lateral parenchyma apoplasm, pathway, sieve tube
- 3 plamalemma, lateral parenchyma plasmalemma, lateral parenchyma apoplasm, lateral
- 4 parenchyma symplasm. Operator Δ means '[in] – [out]'.

1 Figure 1

2

3

4

5

1 Figure 2.

1 Figure 3

Phloem
sieve tube

Lateral tissues

Xylem
vessel

$$\Delta P_{ST} = r_{ST} \cdot JW_{ST}$$

$$\Psi_{ST} = P_{ST} + \Pi_{ST}$$

$$\Psi_{PhlApo} = P_{PhlApo} + \Pi_{PhlApo}$$

$$\Delta JW_{ST} = -NZS - JW_{PhlMb} - JW_{SympL}$$

$$JW_{Trsv} = JW_{PhlMb} + JW_{Apo}$$

$$P_{Xyl} - P_{PhlApo} = r_{Trsv} \cdot JW_{Trsv}$$

$$JW_{Apo} = JW_{ParMb}$$

$$\Psi_{ParApo} = P_{ParApo} + \Pi_{ParApo}$$

$$\Psi_{SympL} = P_{SympL} + \Pi_{SympL}$$

$$P_{SympL} - P_{ST} = r_{SympL} \cdot JW_{SympL}$$

$$\Psi_{PhlApo} - \Psi_{ST} = r_{PhlMb} \cdot JW_{PhlMb}$$

$$P_{PhlApo} - P_{ParApo} = r_{Apo} \cdot JW_{Apo}$$

$$\Psi_{ParApo} - \Psi_{SympL} = r_{ParMb} \cdot JW_{ParMb}$$

$$dV_{SympL}/dt = JW_{ParMb} - JW_{SympL} + NZS_{Par}$$

$$r_{Xyl} \cdot JW_{Xyl} = \Delta P_{Xyl}$$

$$JW_{Trsv} + [Transpirat./-Absorb.] = \Delta JW_{Xyl}$$

—— Biological membrane

- - - - Non-membr. boundary

1 Figure 4

Phloem
sieve tube

Lateral tissues

—— Biological membrane
- - - Non-membr. boundary

$$JS_{PhlMb} = (user-defined)$$

$$JS_{ParMb} = (user-defined)$$

$$JS_{Apo} = (user-defined)$$

$$JS_{SympI} = C_{sympI/ST} \cdot JW_{SympI}$$

$$dQ_{ST}/dt = \Delta JS_{ST} + JS_{PhlMb} + JS_{SympI}$$

$$dQ_{SympI}/dt = JS_{ParMb} - JS_{SympI} + (user-defined)$$

$$dQ_{PhlApo}/dt = -JS_{PhlMb} - JS_{Apo} + (user-defined)$$

$$dQ_{ParApo}/dt = JS_{Apo} - JS_{ParMb} + (user-defined)$$

$$C_{ST} = Q_{ST}/V_{ST}$$

$$C_{SympI} = Q_{SympI}/V_{SympI}$$

$$C_{PhlApo} = Q_{PhlApo}/V_{PhlApo}$$

$$C_{ParApo} = Q_{ParApo}/V_{ParApo}$$