



HAL
open science

Two lysine motif receptor-like kinases (VvLYKs) participate in chitin-triggered immunity in grapevine

Daphnée Brulé, Clizia Villano, Laura J. Davies, Lucie Trdá, Justine Claverie, Marie-Claire Heloir, Annick Chiltz, Marielle Adrian, Benoît Darblade, Lena Stransfeld, et al.

► To cite this version:

Daphnée Brulé, Clizia Villano, Laura J. Davies, Lucie Trdá, Justine Claverie, et al.. Two lysine motif receptor-like kinases (VvLYKs) participate in chitin-triggered immunity in grapevine. Future IPM 3.0 towards a sustainable agriculture, IOBC/WPRS., Oct 2017, Riva del Garda, Italy. 380 p. hal-02788249

HAL Id: hal-02788249

<https://hal.inrae.fr/hal-02788249v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



IOBC-WPRS



Future IPM 3.0 towards a sustainable agriculture

IOBC-WPRS general assembly
Meeting of the WGs Integrated protection in viticulture,
Induced resistance in plants against insects and diseases and
Multitrophic interactions in soil

15-20 October 2017, Riva del Garda, Italy



Future IPM 3.0

BOOK OF ABSTRACTS



Two lysine motif receptor-like kinases (VvLYKs) participate in chitin-triggered immunity in grapevine

Daphnée Brulé, Clizia Villano, Laura J. Davies, Lucie Trdá, Justine Claverie, Marie-Claire Héloir, Annick Chiltz, Marielle Adrian, Benoit Darblade, Lena Stransfeld, Freddy Boutrot, Cyril Zipfel, Ian B. Dry, Benoit Poinssot

First, fourth, fifth, sixth, seventh, eighth and fourteenth authors: Agroécologie, Agrosup Dijon, INRA, Université Bourgogne Franche-Comté, CNRS ERL 6003, Dijon, France; second author: University of Naples Federico II, Portici, Naples, Italy; third and previous last author: Commonwealth Scientific and Industrial Research Organisation (CSIRO), Adelaide, South Australia, Australia; ninth author: Elicityl, 746 avenue Ambroise Croizat, F-38920 Crolles, France; tenth, eleventh and twelfth author: The Sainsbury Laboratory, Norwich Research Park, Norwich, NR4 7UH, United Kingdom.
E-mail address: benoit.poinssot@inra.fr

Highlights

- Two Pattern Recognition Receptors (PRRs) VvLYK1-1 and VvLYK1-2 participate in the signaling of chito-oligosaccharides in grapevine
- VvLYK1-1 is involved in powdery mildew resistance

Introduction

In nature, plants are constantly exposed to potentially pathogenic microbes such as bacteria, fungi, oomycetes or viruses. However, plants have developed effective immune systems triggering various defence reactions against invading pathogens upon the perception of pathogen-associated molecular patterns (PAMPs; Dodds and Rathjen, 2010). The recognition of these conserved microbial signatures is ensured by Pattern Recognition Receptors (PRRs) which also detect plant endogenous molecules released during pathogen invasion, called damage-associated molecular patterns (DAMPs; Boller and Felix, 2009).

Chitin, a fungal cell wall component, is a well-known PAMP that triggers defence responses in many mammal and plant species. The aim of the study was to determine the effects of chito-oligosaccharides on grapevine's immunity and identify the receptor(s) involved in the perception of chito-oligosaccharides in grapevine.

Material and methods

Grapevine cells (*Vitis vinifera* cv Gamay) were cultivated as described in Gauthier et al. (2014). *Arabidopsis thaliana* plants from wild-type (WT) Columbia (Col-0), mutant and transgenic lines were grown *in vitro* for two weeks in controlled conditions for defence responses or in jiffy peat pellets in a controlled growth chamber for four weeks for protection assays. Grapevine cells or *Arabidopsis* plants were treated with water, chitin, chitosan (Elicityl, 0.1 g/l for cells and 1 g/l for plants) or flagelline (10 μ M) taken as a positive control. ROS production and cytosolic Ca^{2+} variations ($[\text{Ca}^{2+}]_{\text{cyt}}$) in grapevine cells were performed according to Dubreuil-Maurizi et al. (2010) after elicitor treatments, by measuring the chemiluminescence of luminol for H_2O_2 production and using apoaequorin expressing cells to detect variations of $[\text{Ca}^{2+}]_{\text{cyt}}$. Protein extraction, SDS-PAGE and western blotting for MAPK phosphorylation analysis were carried out as previously described (Trdá et al., 2014). RNA extraction and quantitative real-time PCR were performed using primers for the