

Connecting the dots between computational tools to analyse soil-root water relations

Sixtine Passot, Valentin Couvreur, Félicien Meunier, Xavier Draye, Mathieu Javaux, Daniel Leitner, Loïc Pagès, Andrea Schnepf, Jan Vanderborght, Guillaume Lobet

▶ To cite this version:

Sixtine Passot, Valentin Couvreur, Félicien Meunier, Xavier Draye, Mathieu Javaux, et al.. Connecting the dots between computational tools to analyse soil-root water relations. 2018. hal-02788325

HAL Id: hal-02788325 https://hal.inrae.fr/hal-02788325

Preprint submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

bioRxiv preprint first posted online May. 9, 2018; doi: http://dx.doi.org/10.1101/312918. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. It is made available under a CC-BY 4.0 International license.

Preprint article

Connecting the dots between computational tools to analyse soil-root water relations

Sixtine Passot⁺¹, Valentin Couvreur⁺¹, Félicien Meunier^{+1,2}, Xavier Draye¹, Mathieu Javaux^{1,3}, Daniel Leitner⁴, Loïc Pagès⁵, Andrea Schnepf³, Jan Vanderborght³ and Guillaume Lobet^{1,3,0}

⁺ These authors contributed equally to this work.

- ° Corresponding author. +32 10 47 92 87
- ¹ Earth and Life Institute, Université catholique de Louvain, Louvain-Ia-Neuve, Belgium
- ² Computational and Applied Vegetation Ecology lab, Ghent University, Gent, Belgium

³ Agrosphere, IBG3, Forschungszentrum Jülich, Jülich, Germany

- ⁴ Simulationswerkstatt, Ortmayrstrasse 20, A-4060 Leonding, Austria
- ⁵ INRA, Centre PACA, UR 1115 PSH, 84914 Avignon cedex 9, France

Abstract

In the recent years, many computational tools, such as image analysis, data management, process-based simulation and upscaling tools, were developed to help quantify and understand water flow in the soil-root system, at multiple scales (tissue, organ, plant and population). Several of these tools work together or, at least, are compatible. However, for the un-informed researcher, they might seem disconnected, forming a unclear and disorganised succession of tools.

> In this article, we present how different pieces of work can be further developed by connecting them to analyse soil-root-water relations in a comprehensive and structured network. This "explicit network of soil-root computational tools" informs the reader about existing tools and help them understand how their data (past and future) might fit within the network. We also demonstrate the novel possibilities of scale-consistent parameterizations made possible by the network with a set of case studies from the literature. Finally, we discuss existing gaps in the network and how we can move forward to fill them.

10

15

20

25

Highlights

Many computational tools exist to quantify water flow in the soil-root system. These tools can be arranged in a comprehensive network that can be leveraged to better interpret experimental data.

35

Keywords

Computational tools, image analysis, simulation, network, root, soil, water

Manuscript information

- Words: 5568
- Figures: 6 colors
- Table: 1
- Web application: https://plantmodelling.shinyapps.io/water_network/.

Sixtine Passot	sixtine.passot@uclouvain.be	0000-0002-6191-723X
Valentin Couvreur	valentin.couvreur@uclouvain.be	0000-0002-1087-3978
Félicien Meunier	felicien.meunier@uclouvain.be	0000-0003-2486-309X
Xavier Draye	xavier.draye@uclouvain.be	0000-0002-3637-3330
Mathieu Javaux	mathieu.javaux@uclouvain.be	0000-0002-6168-5467
Daniel Leitner	daniel.leitner@simwerk.at	
Loïc Pagès	loic.pages@inra.fr	0000-0002-2476-6401
Andrea Schnepf	a.schnepf@fz-juelich.de	0000-0003-2203-4466
Jan Vanderborght	j.vanderborght@fz-juelich.de	0000-0001-7381-3211
Guillaume Lobet	g.lobet@fz-juelich.de	0000-0002-5883-4572

Glossary

Term	Definition	Reference
Standard Uptake Fraction	Relative distribution of root water uptake between root segments when water is equally available in space (units: %)	[1]
High pressure flow meter	Device designed to measure the root system conductance by perfusing pressurized water into a root system opposite from the natural direction of the transpiration stream	[2]
Root pressure probe	Device designed to measure the hydraulic conductance of a single root through variations of water pressure and flow at the cut end of a root	[3]
Cell pressure probe	Device designed to measure the hydraulic conductivity of the membranes of a single plant cell by observing the relaxation time of water pressure pulses applied to the cell	[4]
RSML	Root System Markup Language: File format for the storage of root system data	[5]
MTG	Multi-Tree Graph: File format for the storage of multidimensional tree information, designed for plant models.	[6]
FSPM	Functional-Structural Plant Model: computer model of plant that combine a detailed representation of the plant 3D architecture with functional properties (radial hydraulic conductivity, solute permeability, etc.)	[7]
Parameter	Fixed input of the models that characterizes a specific property of the system, within the scope of study. For instance, branching rates are parameters in root architectural models	
State variable	Variable that characterizes the state of the system at any moment of the simulation. For instance, the water potential within the plant is a state variable in water FSPM	
Boundary conditions	Variables constraining the model at its external boundaries for the entire duration of the simulation. For instance, in model of water flow within the root system, evaporative demand or soil water potentials are the plant boundary conditions.	
Upscaled property	System property that is an output of the model, at a higher scale than the input parameters. For instance, the root radial conductivity is an upscaled property of models of root organ water flow.	

Water flow in the soil-root system

130 135	Water deficit is one of the most dramatic abiotic stresses in agriculture [8]. It occurs when leaf water supply is limited by either the low potential of soil water, and/or by the high hydraulic resistance of the soil-plant system [9]. At this point, the atmospheric demand for water is hardly met and stomata close, reducing the plant transpiration and photosynthesis. To investigate when such limitation occurs, the complex plant-soil-atmosphere system is often conceptualised as a multidimensional hydraulic network, in which both soil and root hydraulic properties may substantially control shoot water supply [10–12].
140	The structural properties of the roots compose the first dimension of the soil-root hydraulic network. Structural properties refer to the physical position and arrangement of the objects of interest. They can be conceptualised at the tissue/organ (transversal anatomy, fig. 1A), plant (root architecture, fig. 1B), or population scale (rooting density profile, fig. 1C).
145	A second dimension, overlaying structural properties, encompasses the system functional properties . When studying water movement, functional properties often refer to hydraulic conductivities or reflection coefficients. Like in the structural layer, these properties can be defined at different scales. Local radial and axial hydraulic conductivities can be defined at the
150	organ scale (fig. 1D) while the entire root system of a single plant can be characterized by its conductance (fig. 1E) and would relate to plant water status [13]. An extension of this property to the population scale is the plant population hydraulic conductance per unit horizontal area (fig. 1F), common in canopy models [14], and recently integrated in root models [15].
155	Finally, a third dimension describes the plant environment . In this contribution, we focus on the soil compartment, which includes the rhizosphere and the bulk soil (fig. 1G). Their respective spatial domains are concentric around individual roots, and their properties differ substantially, so that the rhizosphere is often considered to critically affect plant water availability under water deficit [16,17]. The bulk soil and rhizosphere
160	hydraulic properties may be described by their water retention and hydraulic conductivity curves. The former defines the pressure needed to extract water from the porous media, and the latter the relation between water flux and water potential gradient in space [18]. The water potential that defines the energy level of water is a critical environmental variable, driving the flow of water in the soil-plant system. Similarly to the plant, the soil could be divided

165	into functional and structural components and described according to the studied scale. However, we did not explicit this separation in the following as we rather focus on the plant property description in this study.
170	Each element of the network is dynamic and heterogeneous. Root systems grow, develop and take up water, while soil water content continuously changes in response to root water uptake and climatic conditions, potentially resulting in complex system behaviour. In addition, some key variables and parameters are hard (if not impossible) to quantify experimentally. As a result, the whole system is difficult to apprehend, and novel approaches might prove useful to study it.
175	In the recent years, many computational tools (image analysis, data analysis, process based modelling and upscaling tools) were developed to help quantify and understand water dynamics in the soil-plant system. Some of these tools were developed to work together, or at least be compatible. However, for the uninformed researcher, they might seem disconnected
180	forming a collection of tools with, at best, a common target (plant-water relation exploration) but unrelated to each other.
	The overall objective of this paper is to draw and discuss the role of a functional landscape of interconnected experiments and models for the study of soil-plant water relations. It is articulated as 3 sub-objectives: (i) to inform
185	readers about existing procedures and tools used for the quantification of water flow at the organ, root system and plant population scales, as well as their interconnections forming a comprehensive, though non-exhaustive, network, (ii) to provide examples of studies combining experiments, analytical and modelling tools in this network, motivating the use of such
190	approaches to enhance interpretations of available and future data, and (iii) to identify gaps in the network and argue for a better integration of future tools in this workflow with appropriate experiment and model design. A web interface was developed to help researchers use the network: it is available at <u>https://plantmodelling.shinyapps.io/water_network/</u> .

bioRxiv preprint first posted online May. 9, 2018; doi: http://dx.doi.org/10.1101/312918. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. It is made available under a CC-BY 4.0 International license.

Passot, Couvreur, Meunier et al. 2018 - Tools for the analysis of water relations - Preprint



Figure 1: Quantifying water relations in the soil plant system. Tools, properties and state variables used to quantify: (A) the structure of root organ; (B) the structure of root system; (C) the structure of root profiles; (D) the water flow at the organ scale (root section); (E) the water flow at the plant scale (root system); (F) the water flow at the population scale; (G) the water flow in the soil. Without appropriate tools, variables of interest, scales and even plants and their environment seem disconnected.

200

Connecting the dots between research tools

Through four examples, we illustrate how the dots, consisting in apparently scattered data and tools, can be connected together in a comprehensive network. These examples span over the different scales mentioned above (organ, root system and population) and for all of them, we present data that can be obtained experimentally, technical limitations that need to be overcome and computational tools readily available. In all these examples, we focus on the soil-root water relation specificities at different scales, except for one where an architectural root growth model is introduced.

Water flow at the root cross section scale

210	Different tools and techniques exist to quantify root structural properties at the organ scale. Histology and microscopy techniques enable precise observation of root anatomical structures (the interconnected network of cells). For instance, staining or fluorescence microscopy can be used to
215	acquire images of the organization of different cell types within roots and the nature of cell walls [19]. Different image analysis tools are then available to extract quantitative information out of these images. On the one hand, CellSet [20] is currently the only tool that enables a complete digitization of the entire cell network. As an output, each single cell is represented by a set of connected edges and nodes. Unfortunately, depending on the image
220	quality, the unautomated part of the procedure can be time-consuming. On the other hand, RootScan [21], PHIV-RootCell [22] and RootAnalyzer [23] are fully automated tools that can quantify anatomical properties (such as the number of cells or the mean size of each cell type) but do not provide a digitized cell network.
225	As a part of the functional layer, cell hydraulic properties are hard to estimate as water fluxes are difficult to measure at this scale. The cell pressure probe enables this estimation from measurements of water pressure relaxation times of individual cells, at a high time cost [4,24]. Osmotic pressure can be measured using papoliter osmometer [25] or scapping electron microscopy
230	[26]. However, the latter is expensive and generally not part of the standard equipment of a plant physiology laboratory. At the organ scale, the root pressure probe enables the measurement of axial and radial conductivities of root segments [3] and junctions to the stem [27]. Some properties of the system can hardly be determined experimentally such as the partitioning of
235	water pathways across cell layers (apoplastic or cell-to-cell) (Barzana et al, 2012).
	Detailed root cross-section anatomical descriptions and a minimal set of empirical cell hydraulic properties (e.g. permeability of cell walls and membranes) enables the simulation of water flow across root cross sections.
240	Like at other scales, water flow in the system is solved using transfer equation with boundary water pressures and conductances as input parameters. Such a model can estimate the equivalent hydraulic conductivity of the root cylinder as well as the partitioning of water flow between apoplastic and symplastic compartments of the system. For instance, by
245	combining measurements of cell and root permeability with a hydraulic model, [28]demonstrated that water flow is primarily apoplastic in lupin roots.

A recent study took advantage of these computational tools to estimate the contribution of pearl millet root types to water uptake. Five types were identified based on cross-section anatomy: primary roots, crown roots and 3 types of lateral roots [29]. A cross-section was thoroughly digitized for each type of root using CellSet [20] (fig. 2A). Root axial hydraulic conductances were estimated using the simplified model of Hagen-Poiseuille (fig. 2B), based on measured xylem vessels dimensions. The digitized root anatomical network served as input for a mechanistic model of radial water flow in roots, namely MECHA [30] (fig. 2C). The model was used to estimate the radial conductivity of a typical segment of each root type. In this example, different tools (image analysis and modelling) were combined to estimate radial and axial conductivities, based on easy-to-acquire experimental data (cross section images). While complementary measurements of root hydraulic properties will always remain an asset (e.g. in order to cross-validate the estimated properties), this method opens the way to high-throughput estimations of root hydraulic properties.



Figure 2: Details of the connected dots to compute the hydraulic properties of the different root types of pearl millet. Colored parts are the tools, models, properties and state variables used in the approach. Specific tools names were added where relevant. See text for details.A: CellSet, B: Hagen-Poiseuille, C: MECHA, D: Output of the different models

250

255

260

Root system architecture

	Unlike plant cells, the root system of annual crops has a convenient macroscopic scale and all elements (roots) are visible to the human eye.
270	However, the main difficulties faced when retrieving the root system architecture are the hidden nature of this part of the plant, the large number of elements that can possibly overlay and the fragility of the smallest roots,
	making the full excavation of a complete intact root system particularly difficult. Direct manual methods exist to measure single root architectural
275	traits such as the angle of crown roots with a protractor [31] or with the
	basket method [32,33], the length of individual roots with a ruler [31,34] or a
	combination of several root architectural traits [31]. However, these manual methods do not give access to the full root architecture.
	Several digital tools have been developed and are now widely used to
280	access root architectural traits, mostly from images of root system grown in
	specific experimental setups (see [35] for a review of existing root
	www.plant-image-analysis org and will not be detailed here (Lobet et al
	2013. Lobet 2017). The only point to underline is that each tool generally
285	corresponds to a specific growth medium and image capture technique (eg:
	RooTrak applies to root system growing in 3D and imaged with X-ray Micro
	Computed Tomography [36]. While some of these tools have been designed
	to retrace a full root system architecture (often with an important manual
	input), many of them only extract some root architectural traits (eg. mean
290	lateral root length, number of seminal roots, crown root emergence angle).
	Furthermore, even with the use of specifically designed image analysis tools,
	whole root system digitization becomes time consuming as soon as the
	reconstruct full root system architectures from extracted root traits.
295	Root architecture models, such as SimRoot [37], RootBox [38], RootTyp [39],
	ArchiSimple [40], OpenSimRoot [41] or CRootBox [42], are designed to
	simulate root systems from a limited number of traits, given as input
	parameters. The major interest of root architectural models is to generate a
200	large number of contrasted root system architectures. Root system modeling
300	simulation of contrasted architectures, even from synthetic datasets. These
	contrasting architectures can then be tested in different scenarios to identify
	traits that would be beneficial in challenging environments.
	5 5

305

310

315

320

An example that illustrates how root architecture models can be applied to interpret experimental data of other root zone processes is given by Schnepf et al. (2016). Those authors developed a 3D model of the development of mycorrhizal root systems. The model was designed to simulate primary and secondary root infection with arbuscular mycorrhizal (AM) fungi as well as growth of external fungal hyphae in soil. It was calibrated using root architectural data obtained from pot experiments of *Medicago truncatula*, with and without mycorrhizal inoculum of the AM fungus Rhizophagus irregularis BEG 158. In those pots, AM root colonization was determined under a compound microscope and the abundance of *R. irregularis* hyphal biomass was determined using real-time PCR. The root system architecture, however, could not be parameterised from those pot experiments. The authors re-used published images from a previous study [43] (fig. 3A) and re-analysed them with the image analysis tool RootSystemAnalyzer [44] (fig. 3B). The traits extracted with RootSystemAnalyzer served as parameters for the RootBox model (fig. 3C) [38] which was used to simulate the root system development of mature plants, together with the arbuscular mycorrhizal fungi. This example highlights how published data can be reused to obtain input parameters for modelling. The current literature is filled with similar resources, opening up numerous opportunities. It also highlights the importance of sharing raw experimental data (in this case the images).



325 Figure 3: Details of the connected dots for root system architecture generation. Colored parts are the tools, models and properties used in the approach. Specific tools names were added where relevant. See text for details. A: Published data, B: Root System Analyzer, C: RootBox

Water flow in the root system

At the root system scale, understanding which root traits positively influence plant water uptake dynamics for a given pedo-climatic situation remains an 330 important research question. Ideotypes have been proposed, but are always tied to a specific environment [45–49]. Different traits, either functional or structural, have been found to maximize the final crop yield depending on the environment [50]. Ultimately, we need more than single traits or final yield to have a better understanding of plant-environment interactions. We 335 need to understand how water flow within the plant is dynamically regulated, both spatially and temporally. Unfortunately, accurately measuring water flow is often the limiting step of the experimental pipeline. Several techniques exist to dynamically measure changes in soil water content, such as X-ray computed tomography [51], Electrical Resistivity Tomography [52], neutron 340 tomography [53–56], light transmission imaging [57] or Magnetic Resonance Imaging [58–60]. These techniques can be deployed for a relatively high number of plants. However, due to water capillary flow within the soil domain, observed changes in soil water content are rarely (if never) a direct indication of the location of root water uptake. Water uptake rate itself can be 345 estimated using more advanced but time-consuming lab techniques that use tracers, such as deuterated water that is monitored using neutron radiography [61,62]. Functional-structural plant models (FSPM) are often used to decipher plant-environment relationships [7]. FSPMs couple a complete 350 representation of the root system architecture (or whole plant or shoot) with functional properties. Their input parameters are both functional and structural. For FSPMs simulating soil-root interactions, hydraulic parameters can be obtained using a root pressure probe [63] or the outputs of organ scale models but, as stated earlier, are generally difficult to acquire. Thus 355 they are frequently adapted from the available literature. Rhizosphere hydraulic properties can also be coupled to FSPM [64] and would constitute a central component of plant water availability [16]. Rhizosphere properties are however difficult to parametrize, and would display complex temporal dynamics [65]. The FSPM structural input consists of an explicit 360

representation of the root architecture (see Root system architecture section and the related previous case study). Together with the root system geometry, hydraulic properties define the root system hydraulic architecture [11] and are critical for water stress determination [66,67].

365 Water-related FSPMs provide a exhaustive description of the root water relations (uptake rates, water potentials, etc.) in both space and time. Thus,

370

395

they constitute an important way to integrate different types of information about properties of the root system and soil state variables in the root zone, which can be obtained experimentally, and to translate this information into a distributed pattern of water flows and local state variables (e.g. water potentials at the soil-root interface) within the root zone. The latter type of information is, as of today, hardly accessible experimentally. An exhaustive review of FSPMs related to water flow can be found in Ndour et al. [68].

FSPMs can also be used in so-called *inverse modelling* studies. In such 375 case, the output of the model is known and the model is used to estimate one of the input parameters. For instance, the most likely distribution of root hydraulic properties (that are usually assumed to be age- and order-dependent) can be estimated using a soil-root water flow model and laboratory measurements [69]. In this study, measurements of local water 380 fluxes were obtained from neutron radiography at different locations in the root system [56]. As the experiment took place in a rhizotron, the root system could be fully digitized using an appropriate image analysis tool, which provided accurate information on the root system topology and positions in space (fig. 4A and see also Root system architecture section). Water uptake 385 patterns and axial flows within the root system could then be modelled by applying existing water flow equation resolution algorithms [70,71] to the segmented root system (fig. 4B). The water flow model requires boundary conditions that need to be estimated or measured. In this case, the water potential at the root collar was measured using a pressure probe and 390 root-soil interface water potentials were estimated from water content distribution.

> Such a coupling allowed the authors to estimate the parameters of the root hydraulic conductivity function that best fitted the water flow measurements (fig. 4C and D). These parameters then allowed for novel predictions including water uptake and axial flow distributions everywhere in the root system and not only at observed segments, in homogeneous and heterogeneous soil conditions or under various evaporative demands.



Figure 4: Details of the connected dots for estimating root conductivities from experimental observations through inverse modelling. Colored parts are the tools, models and inputs used in the approach. Specific tools names were added where relevant. The red arrow indicate the optimisation step used in the inverse modelling. The red boxes highlight the variables evaluated using the inverse modelling. See text for details. A: SmartRoot, B: Hybrid water flow model, C: Optimisation, D: Root hydraulic properties

Water flow at the population scale

405

400

410

The population level is a pivotal scale. It interfaces with general circulation models that represent, among others, the circulation of the atmosphere and its interaction with land surface for climate forecasting [72]. It also introduces variables of critical agronomic interest like crop yield per acre [73]. Like at other scales, robust predictions of the system behaviour require the ability to quantify system properties and a proper validation, here involving field scale observations of water fluxes. These fluxes can be estimated with heavy instrumentation and data analytics, for instance using eddy covariance flux towers [74] or soil moisture sensors grids [75].

Structural root information can be obtained using either destructive sampling, such as core sampling [76], monolith excavation [77], trenches [78] or root 415 crown excavation [79], or non-destructive ones, such as minirhizotrons [80]. None of these techniques allows for direct reconstruction of the root system, but rather extract synthetic metrics such as a root length density profile, or root crown data (angles, numbers etc.). Some root architectural traits can be derived from data obtained with these techniques using root architecture 420 models and inverse modeling, as stated above [81,82]. Functional plant properties, such as root system and stomatal conductances, which coordinate shoot water supply [13] and underground water uptake distribution [83], can be characterized on individual plants with low-throughput instruments, such as the high-pressure flowmeter [84] or the 425 porometer [85], then scaled to the population level using the planting density. The main limitation of plant measurement in the field is often the limited sample size, that might not reflect the general behaviour of the system. The same critique can be made about soil hydraulic properties estimated on small and (un)disturbed samples as they may not be representative of the 430 hydrological behaviour at the population level [86,87]. These limitations motivate the use of effective descriptions of population water relations, tailored for this specific scale, such as the transpiration correction for "soil water stress" [88] or one-dimensional soil water and nutrient transfer principles [89]. Two major methodologies address the parametrization of effective field water relations. First, the artificial neural 435 network approach takes advantage of the availability of large amounts of data to train a model. It was used to predict canopy water fluxes from state variables such as the vapor pressure deficit and soil moisture [90,91]. Second, the inverse modelling approach (as described previously) builds on state-of-the-art models to simulate spatio-temporal series of the system 440 state. The model parameter values producing simulated series that best match field observations are considered optimal and representative of the system behaviour. This approach was used to connect models of soil and plant water flow to observations of soil moisture and transpiration in an 445 almond orchard, in order to estimate soil and plant properties, as well as the hardly measurable leaching of water below the root zone [92]. Numerous variables can be used for inverse modeling, such as soil water content, isotopes distributions or root length density profiles.

Going one step further, simplistic macroscale models can be derived from equations of water flow at a lower scale, offering an interesting trade-off between functional simplicity and realism. This type of model involves scale-consistent properties and processes. A cross-validation is thus possible between parameter values estimated directly at the macroscale of

interest (e.g. plant population hydraulic conductance per surface area) and derived from the lower scale (e.g. upscaled values derived from root 455 architectural and hydraulic properties). In order to parametrize such a macroscale model of water dynamics in the soil-wheat system. Cai et al. [15] combined one-dimensional process-based models of water flow (i) in soil (Hydrus-1D, [93]) (fig. 5A), (ii) in roots [94], and (iii) in leaves with an isohydric constraint on transpiration [95] (fig. 5B). Regarding soil properties, 460 soil water retention curves were fitted on simultaneous soil water content and pressure head measurements [96] with the software RETC [97] (fig. 5C). However, the parameters of the soil hydraulic conductivity curve were not experimentally determined (fig. 5D). The vertical distribution of roots in the 465 field (root length density profiles over time) was extracted from in situ rhizotube pictures, with the software Rootfly [98] (fig. 5E). Its relative distribution is typically used as proxy for the water uptake distribution in uniformly wet conditions [99,100] (fig. 5F). Plant hydraulic properties could not be observed in situ for the wheat population (fig. 5F-G). 470 An inverse modelling strategy was therefore used to find the "optimal" soil and plant hydraulic properties (fig. 5H) that best fitted the observed soil water status dynamics. The optimized plant hydraulic parameters were cross-validated with properties at the individual plant scale. Conductance parameters obtained for winter wheat at the same stage of maturity using the hydraulic architecture approach turned out to be consistent with the inversely 475 modeled properties at the population scale [15].

In order to limit the number of parameters, this approach requires the assumption that system properties are time invariant (e.g. soil hydraulic conductivity curve). Because root system conductance tends to scale with root length, the root conductance per unit root length was assumed invariant in order to accommodate for root growth. Such a constraint also matters when accounting for the spatial heterogeneity of root development under different soil/microclimate environments in macroscale simulations [101].



485

Figure 5: Details of the connected dots for estimate soil and plant-scale conductivities through inverse modelling. Colored parts are the tools, models and inputs used in the approach. Specific tools names were added where relevant. The red arrow indicate the optimisation step used in the inverse modelling. The red boxes highlight the variables evaluated using the inverse modelling. See text for references to letters A-H.

Discussions and perspectives

490

495

Many computational tools exist to better understand water dynamics in soil-plant systems. These tools span different scales (organ, plant and population), types (image analysis, data storage, simulation models) and computational languages (Python, Fortran, C++, C#, Java, ...). For the average user, this multitude might seem overly complex and hard to understand. Yet, most of the tools could work together and form a continuous network. Using this network, experimental data can be transferred from scale to scale and generate new insights (fig. 6, and case studies developed above). Modelling tools currently present in the network are listed in table 1. This list is non-exhaustive as the objective of this paper is less to review

500

505

510

515

520

525

530

existing tools than to encourage their integration in order to enhance our understanding soil-plant water relations. For image analysis tools, we refer the reader to the <u>www.plant-image-analysis.org</u> database [102,103].

We created an interactive online visualization of the network, which contains links to the different tools. We also added a submission form, such as the community could update the network with new (or missing) tools. The online visualisation tool (open-source) is available at https://plantmodelling.shinyapps.io/water_network/

Identifying gaps in the tool network

Analogies between tool connectivity patterns at each scale in figure 6 reveal the existence of network gaps (represented by the dashed lines in fig.6). Yet, these gaps should not be filled just for the sake of symmetry. Here, we analyze what the function of filling these gaps could be.

The plant "structure" row in Fig. 6 has the most striking pattern, with imaging techniques systematically feeding image-analysis tools. These tools extract two types of data: (i) explicit spatial structures (e.g. RSML, MTG), and (ii) structural pattern properties (e.g. growth rates, branching rates). At the population and plant scales, root development models [39,104] offer the possibility to convert root pattern properties into predicted root structures. While root anatomical patterns can be automatically characterized by image analysis tools such as PHIV-RootCell [22], no root anatomy development model exists at that scale. In the perspective of generating a mechanistic model of a whole plant from the cell scale [105], a root development model would become essential. It would fulfill two main functions: (i) conducting predictions and test hypotheses related to root anatomical development, and (ii) allowing the spatial and temporal interpolation of root anatomies between experimental observations.

Models using explicit root anatomical structures to test hypotheses about hormone signalling [106], tropisms [107] or radial water flow [30] have emerged lately. However, models of axial water flow remain largely underexplored. In the broadly used Poiseuille-Hagen model, only the quantity and diameter of xylem vessels are accounted for. Yet, it is known for a long time that xylem porous plates, pit membranes and persistent primary cross-walls limit root axial conductivity [108–113] and affect the partitioning of water uptake among root types [27].

bioRxiv preprint first posted online May. 9, 2018; doi: http://dx.doi.org/10.1101/312918. The copyright holder for this preprint (which was not peer-reviewed) is the author/funder. It is made available under a CC-BY 4.0 International license.

Passot, Couvreur, Meunier et al. 2018 - Tools for the analysis of water relations - Preprint



Figure 6: Full network of tools and data used to quantify water flow in the soil plant system. The network connects experimental procedures, computational tools and data related to water flow in the soil-plant system. It is 535 organised by scales (organ, plant and population) and by the types of information (structural or functional, see text for details).A. Tools to quantify the water flow at the organ scale (root section). B. Tools to quantify the structure of root organ. C. Tools to quantify the water flow at the plant scale (root system). D. Tools to quantify the structure of root system. E. Tools to quantify the water flow at the population scale. F. Tools to quantify the structure of root profiles. G. Tools to quantify water flow in the soil. 540

> Experimental methods and numerical tools to represent numerically xylem anatomy and hydraulic properties are missing and may reveal a complexity that is neglected so far. Such numerical representations of xylem vessel structure and hydraulic properties in the axial direction would allow the use of alternatives to Hagen-Poiseuille law [114,115]. Lewis and Boose point out that "Ideally, the exact solutions should be used to calculate [volume flow rate] in xylem conduits, but the equations are difficult to solve without the aid of computer" [115]. Computer availability is no longer an issue and we expect that explicit models of xylem flow will soon emerge. We expect that filling this gap will shed light on the role of cross-walls in the generation of root hydraulic types, and in root-leaf preferential connectivity [116].

Similarly, at the soil-root interface, imaging tools are now available to precisely observe processes at the scale of the soil particle and root hair

545

[117]. Connecting such soil-root interface geometrical descriptions to root hydraulic anatomical models would open new avenues to understand how root hairs enhance plant water availability in dry soils [118].

Table 1: List of modelling tools fitting into the network. For image analysis tools, we refer the reader to the <u>www.plant-image-analysis.org</u> website

Туре	Scale	Name	Reference
Functional	Radial water flow	MECHA	[30]
Functional	Root system water flow	R-SWMS	[119]
Functional - structural	Root system water flow	PlaNet-Maize	[120]
Functional - structural	Root system water flow	OpenSimRoot	[121]
Functional	Root system water flow		[122]
Functional	Soil water flow	HYDRUS	[93]
Functional	Soil water flow	RSWMS	[119]
Structural	Root system architecture	CRootBox	[123]
Structural	Root system architecture	ArchiSimple	[39]
Structural	Root system architecture	RootTyp	[124]
Structural	Root system architecture	DigR	[125]
Structural	Root system density		[126]

Soil water fluxes were only explicitly considered in the last case study (population scale). In other case studies, soil was either neglected (organ and plant scales) or included as static boundary conditions. However, in all cases, a model of water flow in the soil domain can be coupled to the plant water flow. Such analyses were for instance carried out at the plant level [41,119,127,128] or the population level [129,130]. Such models may incorporate multiple soil characteristics such as macropores [131], solute convection-dispersion [132] or specific rhizosphere properties [133,134]. For an extensive review of existing soil models, we refer the reader to Vereecken et al. [135].

555

615

Limitations and future developments

	Simulating water fluxes in roots with this collection of tools can either help understanding plant water relations as a main goal or be a tool for further application. These tools could also be used as a side usage of a dataset
625	obtained for other purposes. The advent of imaging in plant sciences and the huge progress made in image analysis allow generating high quality quantitative data of plant structure, suitable for model parameterization. Many models exist at different scales and we highlighted the many possibilities to combine these tools. This set of tools greatly increases the
630	potential of interpretation of experimental data. Yet many authors still publish rich datasets without using modelling tools to interpret them. Using models is not trivial for a large part of the plant science community. Coupling several tools and models together is still rarely applied. Several requirements seem essential to facilitate the use of this pipeline whenever it may add value to
635	the data.
	A lot of image analysis tools and models flourish in the water transport domain. Thanks to the wide breadth of scientific literature available (scientific papers, reviews, websites), developers are usually aware of already existing tools and keep in mind to justify the interest of their tool in this
640	landscape. However, we suggest that further efforts should be made to render the tools compatible with existing ones. In this context, the existence of several tools at the same place of the network (eg. RootTyp and CRootBox for root architecture simulation) is not conflicting. Each one can best suit one scale or specific situation. In our opinion, special attention
645	should be paid to the data format. Indeed, the output data format of upstream tools must be compatible with the input format required by downstream ones. If this is not the case, easy-to-handle tools must exist to convert these data. The multiplication of formats and the need to convert data from one type to another may discourage the use of some of the
650	models. In this respect, the existence of standard formats, such as the Root System Markup Language [5] for root architecture, smooths the interconnection between tools.
655	When a new tool is created, the documentation of its potential connections with existing ones (e.g. in the user guide) would benefit the whole network. It is indeed expected that the knowledge and the use of all modelling tools will increase. It also underlines the need to keep the models and their documentation updated. Pioneering tools sometimes get outdated by new ones that do similar tasks but that are more user-friendly, faster, use the latest formalisms, or are better connected with newly existing tools.

660 Therefore, either the interconnection between tools needs to be part of a huge maintenance effort for already existing tools, or the acceptance that pioneering tools are doomed to sink into oblivion.

Making the different tools freely available to the community is also a key aspect in their long term maintenance [103]. Many different repositories and licences exist so that everyone should be able to find a combination that suits their (and their institution's) needs. Free access to the tools' source codes would indeed greatly facilitate their evolution, reproducibility of the in silico experiments and allow future developers to interconnect them more easily.

670

665

Acknowledgments

The authors would like to thank Jennifer Brophy, Heike Lindner and Thérèse LaRue (Carnegie Institution for Science, Stanford, CA) for their useful comment on the first version of manuscript. This work was also supported by the Belgian French community ARC 16/21-075 project. VC was also supported by the Interuniversity Attraction Poles Programme-Belgian Science Policy (grant IAP7/29). During the preparation of this manuscript, FM was supported by the "Fonds National de la Recherche Scientifique" (FNRS) of Belgium as a Research Fellow and is grateful to this organization for its financial support.

680

675

Author contributions

	SP	VC	FM	XD	MJ	DL	LP	AS	JV	GL
Conceptualization										
Visualization										
Writing - draft										
Writing - review / editing										

695

References

	1.	Couvreur V, Vanderborght J, Javaux M. A simple three-dimensional macroscopic root water uptake model based on the hydraulic architecture approach. Hydrol Earth Syst Sci. Copernicus GmbH; 2012;16: 2957–2971.
700	2.	Tsuda M, Tyree MT. Plant hydraulic conductance measured by the high pressure flow meter in crop plants. J Exp Bot. Soc Experiment Biol; 2000;51: 823.
	3.	Steudle E, Jeschke WD. Water transport in barley roots : Measurements of root pressure and hydraulic conductivity of roots in parallel with turgor and hydraulic conductivity of root cells. Planta. 1983;158: 237–248.
705	4.	Steudle E. Water-relation Parameters of Individual Mesophyll Cells of the Crassulacean Acid Metabolism Plant Kalanchoë daigremontiana. Plant Physiol. 1980;66: 1155–1163.
	5.	Lobet G, Pound MP, Diener J, Pradal C, Draye X, Godin C, et al. Root System Markup Language: Toward a Unified Root Architecture Description Language. Plant Physiol. American Society of Plant Biologists; 2015;167: 617–627.
710	6.	Godin C, Costes E, Sinoquet H. A Method for Describing Plant Architecture which Integrates Topology and Geometry. Ann Bot. 1999;84: 343–357.
	7.	Godin C, Sinoquet H. Functional-structural plant modelling. New Phytol. Blackwell Science Ltd; 2005;166: 705–708.
715	8.	Cattivelli L, Rizza F, Badeck F-W, Mazzucotelli E, Mastrangelo AM, Francia E, et al. Drought tolerance improvement in crop plants: An integrated view from breeding to genomics. Field Crops Res. 2008;105: 1–14.
	9.	Sperry JS, Hacke UG, Oren R, Comstock JP. Water deficits and hydraulic limits to leaf water supply. Plant Cell Environ. 2002;25: 251–263.
720	10.	Draye X, Kim Y, Lobet G, Javaux M. Model-assisted integration of physiological and environmental constraints affecting the dynamic and spatial patterns of root water uptake from soils. J Exp Bot. 2010;61: 2145–2155.
	11.	Lobet G, Couvreur V, Meunier F, Javaux M, Draye X. Plant water uptake in drying soils. Plant Physiol. 2014;164: 1619–1627.
725	12.	Schoppach R my, Wauthelet D, Jeanguenin L, Sadok W. Conservative water use under high evaporative demand associated with smaller root metaxylem and limited trans-membrane water transport in wheat. Funct Plant Biol. CSIRO PUBLISHING; 2013; Available: http://www.publish.csiro.au/?paper=FP13211
	13.	Alsina MM, Smart DR, Bauerle T, de Herralde F, Biel C, Stockert C, et al. Seasonal changes of whole root system conductance by a drought-tolerant grape root system. J Exp Bot. 2011;62: 99–109.
730	14.	Cox PM, Huntingford C, Harding RJ. A canopy conductance and photosynthesis model for use in a GCM land surface scheme. J Hydrol. 1998;212-213: 79–94.
	15.	Cai G, Vanderborght J, Couvreur V, Mboh CM, Vereecken H. Parameterization of Root Water Uptake Models Considering Dynamic Root Distributions and Water Uptake Compensation. Vadose Zone J. 2017;0: 0.
735	16.	Carminati A, Schneider CL, Moradi AB, Zarebanadkouki M, Vetterlein D, Vogel H-J, et al. How

 Ahmed MA, Kroener E, Holz M, Zarebanadkouki M, Carminati A. Mucilage exudation facilitates root water uptake in dry soils. Funct Plant Biol. CSIRO PUBLISHING; 2014;41: 1129–1137. van Genuchten MT. A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. Soil Sci Sco Am J. 1980;44: 891–898. Costa G, Plazanet I. Plant Cell Wall, a Challenge for Its Characterisation. Adv Biol Chem. 2016;06: 70–105. Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP. CellSeT: Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. Plant Cell. American Society of Plant Biologists; 2012; Available: http://www.plantcell.org/content/early/2012/03/30/tpc.112.096289 Burton AL, Williams M, Lynch JP, Brown KM. RootScan: Software for high-throughput anal of root anatomical traits. Plant Soil. Kluwer Academic Publishers; 2012;357: 199–203. Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plan Sci. 2014;5: 770. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Imag Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of a from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1999;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Boi. 1999;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine			the rhizosphere may favor water availability to roots. Vadose Zone J. Soil Science Society; 2011;10: 988–998.
 van Geruchten MT. A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. Soil Sci Soc Am J. 1980;44: 891–898. Costa G, Plazanet I. Plant Cell Wall, a Challenge for Its Characterisation. Adv Biol Chem. 2016;06: 70–105. Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP. CellSeT. Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. Plant Cell. American Society of Plant Biologists, 2012; Available: http://www.plantcell.org/content/early/2012/03/30/tpc.112.096289 Burton AL, Williams M, Lynch JP. Brown KM. RootScam: Software for high-throughput anal of root anatomical traits. Plant Soil. Kluwer Academic Publishers; 2012;37: 189–203. Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plan Sci. 2014;6: 790. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Image Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Erns L, Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of sol-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2016; jpinp.2017.12.019 Bramley H, Turrer NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and a	740	17.	Ahmed MA, Kroener E, Holz M, Zarebanadkouki M, Carminati A. Mucilage exudation facilitates root water uptake in dry soils. Funct Plant Biol. CSIRO PUBLISHING; 2014;41: 1129–1137.
 Costa G, Plazanet I. Plant Cell Wall, a Challenge for Its Characterisation. Adv Biol Chem. 2016;06: 70–105. Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP. CellSeT: Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. Plant Cell. American Society of Plant Biologists; 2012. Available: http://www.plantcell.org/content/Vearly/2012/03/30/tpc.112.096289 Burton AL, Williams M, Lynch JP, Brown KM. RootScan: Software for high-throughput anal of root anatomical traits. Plant Soli. Kluwer Academic Publishers; 2012;357: 189–203. Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plan Sci. 2014;5: 790. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Imag Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137656. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis for fasp from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J. Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarch S, Ontega BM, et al. Characterization of Pear Millel Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont		18.	van Genuchten MT. A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. Soil Sci Soc Am J. 1980;44: 891–898.
 Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP. CellSeT: Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. Plant Cell. American Society of Plant Biologists; 2012; Available: http://www.plantcell.org/content/early/2012/03/30/tpc.112.096289 Burton AL, Williams M, Lynch JP, Brown KM. RootScan: Software for high-throughput anal of root anatomical traits. Plant Soil. Kluwer Academic Publishers; 2012;357: 189–203. Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plan Sci. 2014;5: 790. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Imag Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctiti J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turrer NC, Turrer DW, Tyerman SD, Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Th		19.	Costa G, Plazanet I. Plant Cell Wall, a Challenge for Its Characterisation. Adv Biol Chem. 2016;06: 70–105.
 Burton AL, Williams M, Lynch JP, Brown KM. RootScan: Software for high-throughput anal of root anatomical traits. Plant Soil. Kluwer Academic Publishers; 2012;357: 189–203. Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plant Sci. 2014;5: 790. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Image Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Image Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Java	745	20.	Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP. CellSeT: Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. Plant Cell. American Society of Plant Biologists; 2012; Available: http://www.plantcell.org/content/early/2012/03/30/tpc.112.096289
 Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell supervised image analysis tool for rice root anatomical parameter quantification. Front Plan Sci. 2014;5: 790. Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Imag Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Erns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. 	750	21.	Burton AL, Williams M, Lynch JP, Brown KM. RootScan: Software for high-throughput analysis of root anatomical traits. Plant Soil. Kluwer Academic Publishers; 2012;357: 189–203.
 Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Imag Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015; e0137655. Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ro pressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic roots and the shaping of the device optic of roots and the shaping of the process of the device of roots and the shaping of the device optic of roots and the shaping of the 		22.	Lartaud M, Perin C, Courtois B, Thomas E, Henry S, Bettembourg M, et al. PHIV-RootCell: a supervised image analysis tool for rice root anatomical parameter quantification. Front Plant Sci. 2014;5: 790.
 Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882. Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ropressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitoric response of roots and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determined of the context and the shaping of the determ	755	23.	Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ. RootAnalyzer: A Cross-Section Image Analysis Tool for Automated Characterization of Root Cells and Tissues. PLoS One. 2015;10: e0137655.
 Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926. Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ropressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctio J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Scil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitopic response of roots and the shaping of the desired of the field of the field. Plant Scil. 2010;341: 75–87. 		24.	Hose E, Steudle E, Hartung W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. Planta. Springer-Verlag; 2000;211: 874–882.
 Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high ropressure. J Exp Bot. 1998;49: 1539–1544. Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junction J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the composite of the More. 	760	25.	Malone M, Leigh RA, Tomos AD. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. Plant Cell Environ. 1989;12: 919–926.
 Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junction J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the second second se		26.	Enns L. Solute concentrations in xylem sap along vessels of maize primary roots at high root pressure. J Exp Bot. 1998;49: 1539–1544.
 Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150 348–364. Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the 	765	27.	Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctions. J Plant Physiol. 2018; doi:10.1016/j.jplph.2017.12.019
 Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829. Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the 		28.	Bramley H, Turner NC, Turner DW, Tyerman SD. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. Plant Physiol. 2009;150: 348–364.
 Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the shaping of the shape of the shaping of the shape of the s	770	29.	Passot S, Gnacko F, Moukouanga D, Lucas M, Guyomarc'h S, Ortega BM, et al. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. Front Plant Sci. 2016;7: 829.
 Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the shaping	775	30.	Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X. Novel multiscale insights into the composite nature of water transport in roots [Internet]. bioRxiv. 2017. p. 147314. doi:10.1101/147314
32. Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the		31.	Trachsel S, Kaeppler SM, Brown KM, Lynch JP. Shovelomics: high throughput phenotyping of maize (Zea mays L.) root architecture in the field. Plant Soil. 2010;341: 75–87.
root system in cereal plants. Environ Exp Bot. 1993;33: 141–158.		32.	Oyanagi A, Nakamoto T, Morita S. The gravitropic response of roots and the shaping of the root system in cereal plants. Environ Exp Bot. 1993;33: 141–158.

780	33.	Uga Y, Okuno K, Yano M. Dro1, a major QTL involved in deep rooting of rice under upland field conditions. J Exp Bot. 2011;62: 2485–2494.
	34.	Pritchard J, Barlow PW, Adam JS, Tomos AD. Biophysics of the inhibition of the growth of maize roots by lowered temperature. Plant Physiol. 1990;93: 222–230.
785	35.	Paez-Garcia A, Motes CM, Scheible W-R, Chen R, Blancaflor EB, Monteros MJ. Root Traits and Phenotyping Strategies for Plant Improvement. Plants. 2015;4: 334–355.
	36.	Mairhofer S, Zappala S, Tracy SR, Sturrock C, Bennett M, Mooney SJ, et al. RooTrak: automated recovery of three-dimensional plant root architecture in soil from x-ray microcomputed tomography images using visual tracking. Plant Physiol. 2012;158: 561–569.
790	37.	Lynch JP, Nielsen KL, Davis RD, Jablokow AG. SimRoot: Modelling and visualization of root systems. Plant Soil. Kluwer Academic Publishers; 1997;188: 139–151.
	38.	Leitner D, Klepsch S, Bodner G, Schnepf A. A dynamic root system growth model based on L-Systems. Plant Soil. Kluwer Academic Publishers; 2010;332: 117–192.
795	39.	Pagès L, Bécel C, Boukcim H, Moreau D, Nguyen C, Voisin A-S. Calibration and evaluation of ArchiSimple, a simple model of root system architecture. Ecol Modell. Elsevier B.V.; 2013;290: 76–84.
	40.	Pagès L, Bécel C, Boukcim H, Moreau D, Nguyen C, Voisin A-S. Calibration and evaluation of ArchiSimple, a simple model of root system architecture. Ecol Modell. 2014;290: 76–84.
800	41.	Postma JA, Kuppe C, Owen MR, Mellor N, Griffiths M, Bennett MJ, et al. OpenSimRoot: widening the scope and application of root architectural models. New Phytol. 2017;215: 1274–1286.
	42.	Schnepf A, Leitner D, Landl M, Lobet G, Mai TH, Morandage S, et al. CRootBox: a structural–functional modelling framework for root systems. Ann Bot. 2018; doi:10.1093/aob/mcx221
805	43.	Bourion V, Martin C, de Larambergue H, Jacquin F, Aubert G, Martin-Magniette M-L, et al. Unexpectedly low nitrogen acquisition and absence of root architecture adaptation to nitrate supply in a Medicago truncatula highly branched root mutant. J Exp Bot. 2014;65: 2365–2380.
	44.	Leitner D, Felderer B, Vontobel P, Schnepf A. Recovering root system traits using image analysis exemplified by two-dimensional neutron radiography images of lupine. Plant Physiol. 2014;164: 24–35.
810	45.	Pagès L. Links between root developmental traits and foraging performance. Plant Cell Environ. 2011;34: 1749–1760.
	46.	Lynch JP. Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root systems. Ann Bot. 2013;112: 347–357.
815	47.	Dencic S. Designing a Wheat Ideotype with Increased Sink Capacity. Plant Breed. 1994;112: 311–317.
	48.	Wasson AP, Richards RA, Chatrath R, Misra SC, Prasad SVS, Rebetzke GJ, et al. Traits and selection strategies to improve root systems and water uptake in water-limited wheat crops. J Exp Bot. 2012;63: 3485–3498.
820	49.	Guswa AJ. Effect of plant uptake strategy on the water-optimal root depth. Water Resour Res. 2010;46. doi:10.1029/2010wr009122
	50.	Tardieu F. Any trait or trait-related allele can confer drought tolerance: just design the right drought scenario. J Exp Bot. 2012;63: 25–31.

	51.	Hainsworth JM, Aylmore L. The use of computer assisted tomography to determine spatial distribution of soil water content. Aust J Soil Res. 1983;21: 435.
825	52.	Garré S, Coteur I, Wongleecharoen C, Hussain K, Omsunrarn W, Kongkaew T, et al. Can We Use Electrical Resistivity Tomography to Measure Root Zone Dynamics in Fields with Multiple Crops? Procedia Environmental Sciences. 2013;19: 403–410.
	53.	Carminati A, Moradi AB, Vetterlein D, Vontobel P, Lehmann E, Weller U, et al. Dynamics of soil water content in the rhizosphere. Plant Soil. 2010;332: 163–176.
830	54.	Esser HG, Carminati A, Vontobel P, Lehmann EH, Oswald SE. Neutron radiography and tomography of water distribution in the root zone. J Plant Nutr Soil Sci. 2010;173: 757–764.
	55.	Tötzke C, Kardjilov N, Manke I, Oswald SE. Capturing 3D Water Flow in Rooted Soil by Ultra-fast Neutron Tomography. Sci Rep. 2017;7: 6192.
835	56.	Zarebanadkouki M, Kroener E, Kaestner A, Carminati A. Visualization of root water uptake: quantification of deuterated water transport in roots using neutron radiography and numerical modeling. Plant Physiol. 2014;166: 487–499.
	57.	Garrigues E, Doussan C, Pierret A. Water Uptake by Plant Roots: I – Formation and Propagation of a Water Extraction Front in Mature Root Systems as Evidenced by 2D Light Transmission Imaging. Plant Soil. 2006;283: 83–98.
840	58.	Jahnke S, Menzel MI, van Dusschoten D, Roeb GW, Bühler J, Minwuyelet S, et al. Combined MRI-PET dissects dynamic changes in plant structures and functions. Plant J. 2009;59: 634–644.
845	59.	Pohlmeier A. MRI in Soils: Determination of Water Content Changes Due to Root Water Uptake by Means of a Multi-Slice-Multi-Echo Sequence (MSME). The Open Magnetic Resonance Journal. 2010;3: 69–74.
	60.	Rascher U, Blossfeld S, Fiorani F, Jahnke S, Jansen M, Kuhn AJ, et al. Non-invasive approaches for phenotyping of enhanced performance traits in bean. Funct Plant Biol. 2011;38: 968.
850	61.	Zarebanadkouki M, Kroener E, Kaestner A, Carminati A. Visualization of root water uptake: quantification of deuterated water transport in roots using neutron radiography and numerical modeling. Plant Physiol. 2014;166: 487–499.
	62.	Warren JM, Bilheux H, Kang M, Voisin S, Cheng C-L, Horita J, et al. Neutron imaging reveals internal plant water dynamics. Plant Soil. 2013;366: 683–693.
855	63.	Steudle E, Rr O, Schulze E. Water transport in maize roots. Measurement of hydraulic conductivity, solute permeability and of 1987;
	64.	Schneider CL, Attinger S, Delfs J-O, Hildebrandt A. Implementing small scale processes at the soil-plant interface – the role of root architectures for calculating root water uptake profiles. Hydrol Earth Syst Sci Discuss. 2009;6: 4233–4264.
860	65.	Carminati A, Vetterlein D. Plasticity of rhizosphere hydraulic properties as a key for efficient utilization of scarce resources. Ann Bot. 2013;112: 277–290.
	66.	Vadez V. Root hydraulics: The forgotten side of roots in drought adaptation. Field Crops Res. Elsevier B.V.; 2014; 1–10.
865	67.	Leitner D, Meunier F, Bodner G, Javaux M, Schnepf A. Impact of contrasted maize root traits at flowering on water stress tolerance – A simulation study. Field Crops Res. 2014;165: 125–137.
	68.	Ndour A, Vadez V, Pradal C, Lucas M. Virtual Plants Need Water Too: Functional-Structural

		Root System Models in the Context of Drought Tolerance Breeding. Front Plant Sci. 2017;8: 1577.
870	69.	Zarebanadkouki M, Meunier F, Couvreur V, Cesar J, Javaux M, Carminati A. Estimation of the hydraulic conductivities of lupine roots by inverse modelling of high-resolution measurements of root water uptake. Ann Bot. 2016; doi:10.1093/aob/mcw154
	70.	Doussan C, Pagès L, Vercambre G. Modelling of the hydraulic architecture of root systems: An integrated approach to water absorption - Model description. Ann Bot. Annals Botany Co; 1998;81: 213–223.
875	71.	Meunier F, Draye X, Vanderborght J, Javaux M, Couvreur V. A hybrid analytical-numerical method for solving water flow equations in root hydraulic architectures. Appl Math Model. 2017;52: 648–663.
880	72.	Shrestha P, Sulis M, Masbou M, Kollet S, Simmer C. A Scale-Consistent Terrestrial Systems Modeling Platform Based on COSMO, CLM, and ParFlow. Mon Weather Rev. 2014;142: 3466–3483.
	73.	Holzworth D, Huth NI, Fainges J, Brown H, Zurcher E, Cichota R, et al. APSIM Next Generation: Overcoming challenges in modernising a farming systems model. Environmental Modelling & Software. 2018;103: 43–51.
885	74.	McElrone AJ, Shapland TM, Calderon A, Fitzmaurice L, Paw U KT, Snyder RL. Surface renewal: an advanced micrometeorological method for measuring and processing field-scale energy flux density data. J Vis Exp. 2013; e50666.
	75.	Famiglietti JS, Ryu D, Berg AA, Rodell M, Jackson TJ. Field observations of soil moisture variability across scales. Water Resour Res. 2008;44. doi:10.1029/2006wr005804
890	76.	Wasson AP, Rebetzke GJ, Kirkegaard JA, Christopher J, Richards RA, Watt M. Soil coring at multiple field environments can directly quantify variation in deep root traits to select wheat genotypes for breeding. J Exp Bot. 2014;65: 6231–6249.
	77.	Kuchenbuch RO, Horst GH, Buczko U. Spatial distribution of maize roots by complete 3D soil monoltih sampling. Plant Soil. Kluwer Academic Publishers; 2009;315: 297–314.
895	78.	Vepraskas MJ, Hoyt GD. Comparison of the trench-profile and core methods for evaluating root distributions in tillage studies. Agron J. American Society of Agronomy; 1988;80: 166–172.
	79.	Colombi T, Kirchgessner N, Le Marié CA, York LM, Lynch JP, Hund A. Next generation shovelomics: set up a tent and REST. Plant Soil. Kluwer Academic Publishers; 2015; Available: http://link.springer.com/10.1007/s11104-015-2379-7
900	80.	Rewald B, Ephrath J. Minirhizotron Techniques. In: Beeckman T, editor. Plant Roots. CRC Press; 2013. pp. 42–1–42–16.
	81.	Garré S, Pagès L, Laloy E, Javaux M, Vanderborght* J, Vereecken H. Parameterizing a Dynamic Architectural Model of the Root System of Spring Barley from Minirhizotron Data. Vadose Zone J. 2012;11: 0.
905	82.	Vansteenkiste J, Van Loon J, Garré S, Pagès L, Schrevens E, Diels J. Estimating the parameters of a 3-D root distribution function from root observations with the trench profile method: case study with simulated and field-observed root data. Plant Soil. 2013;375: 75–88.
	83.	Meunier F, Rothfuss Y, Bariac T, Biron P, Richard P, Durand J-L, et al. Measuring and Modeling Hydraulic Lift of Using Stable Water Isotopes. Vadose Zone J. 2018;17: 0.
910	84.	Tsuda M, Tyree MT. Plant hydraulic conductance measured by the high pressure flow meter in

crop plants. J Exp Bot. 2000;51: 823-828.

	85.	Rodrigues ML, Lucília Rodrigues M, Santos TP, Rodrigues AP, de Souza CR, Lopes CM, et al. Hydraulic and chemical signalling in the regulation of stomatal conductance and plant water use in field grapevines growing under deficit irrigation. Funct Plant Biol. 2008;35: 565.
915	86.	Hopmans JW, Nielsen DR, Bristow KL. How useful are small-scale soil hydraulic property measurements for large-scale vadose zone modeling? Geophysical Monograph Series. 2002. pp. 247–258.
920	87.	Vrugt JA, Schoups G, Hopmans JW, Young C, Wallender WW, Harter T, et al. Inverse modeling of large-scale spatially distributed vadose zone properties using global optimization. Water Resour Res. 2004;40. doi:10.1029/2003wr002706
	88.	Verhoef A, Egea G. Modeling plant transpiration under limited soil water: Comparison of different plant and soil hydraulic parameterizations and preliminary implications for their use in land surface models. Agric For Meteorol. 2014;191: 22–32.
925	89.	Baram S, Couvreur V, Harter T, Read M, Brown PH, Kandelous M, et al. Estimating Nitrate Leaching to Groundwater from Orchards: Comparing Crop Nitrogen Excess, Deep Vadose Zone Data-Driven Estimates, and HYDRUS Modeling. Vadose Zone J. 2016;15: 0.
	90.	Couvreur V, Kandelous MM, Sanden BL, Lampinen BD, Hopmans JW. Downscaling transpiration rate from field to tree scale. Agric For Meteorol. 2016;221: 71–77.
930	91.	Whitley R, Taylor D, Macinnis-Ng C, Zeppel M, Yunusa I, O'Grady A, et al. Developing an empirical model of canopy water flux describing the common response of transpiration to solar radiation and VPD across five contrasting woodlands and forests. Hydrol Process. 2012;27: 1133–1146.
	92.	Vrugt JA, van Wijk MT, Hopmans JW, Šimunek J. One-, two-, and three-dimensional root water uptake functions for transient modeling. Water Resour Res. 2001;37: 2457–2470.
935	93.	Simunek J, Van Genuchten M. Numerical model for simulating multiple solute transport in variably-saturated soils. Ground Water. ars.usda.gov; 1995; Available: https://www.ars.usda.gov/research/publications/publication/?seqNo115=64408
940	94.	Couvreur V, Vanderborght J, Beff L, Javaux M. Horizontal soil water potential heterogeneity: simplifying approaches for crop water dynamics models. Hydrol Earth Syst Sci Discuss. 2014;11: 1203–1252.
	95.	Couvreur V, Vanderborght J, Draye X, Javaux M. Dynamic aspects of soil water availability for isohydric plants: Focus on root hydraulic resistances. Water Resour Res. 2014;50: 8891–8906.
945	96.	Cai G, Vanderborght J, Klotzsche A, van der Kruk J, Neumann J, Hermes N, et al. Construction of Minirhizotron Facilities for Investigating Root Zone Processes. Vadose Zone J. Madison, WI: The Soil Science Society of America, Inc.; 2016;15: 1–13.
	97.	van Genuchten M, Leij FJ, Yates. MS. The RETC Code for Quantifying the Hydraulic Functions of Unsaturated Soils. EPA Report 600/2-91/065, U.S. Salinity Laboratory, USDA, ARS, Riverside, California.; 1991.
950	98.	Zeng G, Birchfield ST, Wells CE. Automatic discrimination of fine roots in minirhizotron images. New Phytol. Blackwell Science Ltd; 2008;117: 549–557.
	99.	Feddes RA, Kowalik PJ, Zaradny H. Simulation of Field Water Use and Crop Yield. Wiley; 1978.
	100	Šimůnek J, Hopmans JW. Modeling compensated root water and nutrient uptake. Ecol

955	Modell. Elsevier B.V.; 2009;220: 505–521.
	101. Cai G, Vanderborght J, Langensiepen M, Schnepf A, Hüging H, Vereecken H. Root growth, water uptake, and sap flow of winter wheat in response to different soil water conditions. Hydrol Earth Syst Sci. Copernicus GmbH; 2018;22: 2449.
960	102. Lobet G, Draye X, Périlleux C. An online database for plant image analysis software tools. Plant Methods. 2013;9: 38.
	103. Lobet G. Image Analysis in Plant Sciences: Publish Then Perish. Trends Plant Sci. 2017;22: 559–566.
	104. Dupuy L, Gregory PJ, Bengough AG. Root growth models: towards a new generation of continuous approaches. J Exp Bot. Soc Experiment Biol; 2010;61: 2131–2143.
965	105. Band LR, Wells DM, Fozard JA, Ghetiu T, French AP, Pound MP, et al. Systems Analysis of Auxin Transport in the Arabidopsis Root Apex. Plant Cell. American Society of Plant Biologists; 2014; Available: http://www.plantcell.org/cgi/doi/10.1105/tpc.113.119495
970	106. Stoma S, Lucas M, Chopard J, Schaedel M, Traas J, Godin C. Flux-based transport enhancement as a plausible unifying mechanism for auxin transport in meristem development. PLoS Comput Biol. 2008;4: e1000207.
	107. Robbins NE 2nd, Dinneny JR. Growth is required for perception of water availability to pattern root branches in plants. Proc Natl Acad Sci U S A. 2018; doi:10.1073/pnas.1710709115
	108. Choat B, Jansen S, Brodribb TJ, Cochard H, Delzon S, Bhaskar R, et al. Global convergence in the vulnerability of forests to drought. Nature. 2012;491: 752–755.
975	109. Choat B, Cobb AR, Jansen S. Structure and function of bordered pits: new discoveries and impacts on whole-plant hydraulic function. New Phytol. 2008;177: 608–625.
980	 Sanderson J, Whitbread FC, Clarkson DT. Persistent xylem cross-walls reduce the axial hydraulic conductivity in the apical 20 cm of barley seminal root axes: implications for the driving force for water movement. Plant Cell Environ. Blackwell Publishing Ltd; 1988;11: 247–256.
	 Shane MW, McCully ME, Canny MJ. Architecture of Branch-root Junctions in Maize: Structure of the Connecting Xylem and the Porosity of Pit Membranes. Ann Bot. Oxford University Press; 2000;85: 613–624.
985	112. Tixier A, Herbette S, Jansen S, Capron M, Tordjeman P, Cochard H, et al. Modelling the mechanical behaviour of pit membranes in bordered pits with respect to cavitation resistance in angiosperms. Ann Bot. 2014;114: 325–334.
	 Brodersen CR, McElrone AJ, Choat B, Lee EF, Shackel KA, Matthews MA. In vivo visualizations of drought-induced embolism spread in Vitis vinifera. Plant Physiol. 2013;161: 1820–1829.
990	114. Giordano R, Salleo A, Salleo S, Wanderlingh F. Flow in xylem vessels and Poiseuille's law. Can J Bot. 1978;56: 333–338.
	 Lewis AM, Boose ER. Estimating Volume Flow Rates Through Xylem Conduits. Am J Bot. 1995;82: 1112.
995	116. David TS, David JS, Pinto CA, Cermak J, Nadezhdin V, Nadezhdina N. Hydraulic connectivity from roots to branches depicted through sap flow: analysis on a Quercus suber tree. Funct Plant Biol. CSIRO PUBLISHING; 2012;39: 103–115.
	117. Keyes SD, Daly KR, Gostling NJ, Jones DL, Talboys P, Pinzer BR, et al. High resolution synchrotron imaging of wheat root hairs growing in soil and image based modelling of

1000	phosphate uptake. New Phytol. Blackwell Science Ltd; 2013; Available: http://doi.wiley.com/10.1111/nph.12294
	118. Carminati A, Passioura JB, Zarebanadkouki M, Ahmed MA, Ryan PR, Watt M, et al. Root hairs enable high transpiration rates in drying soils. New Phytol. 2017; doi:10.1111/nph.14715
	119. Javaux M, Schroeder T, Vanderborght J, Vereecken H. Use of a three-dimensional detailed modeling approach for predicting root water uptake. Vadose Zone J. 2008;7: 1079–1088.
1005	120. Lobet G, Pagès L, Draye X. A modeling approach to determine the importance of dynamic regulation of plant hydraulic conductivities on the water uptake dynamics in the soil-plant-atmosphere system. Ecol Modell. Elsevier B.V.; 2014;290: 65–75.
1010	 Postma JA, Kuppe C, Owen MR, Mellor N, Griffiths M, Bennett MJ, et al. OpenSimRoot: widening the scope and application of root architectural models. New Phytol. 2017; doi:10.1111/nph.14641
	122. Doussan C, Pierret A, Garrigues E, Pagès L. Water Uptake by Plant Roots: II – Modelling of Water Transfer in the Soil Root-system with Explicit Account of Flow within the Root System – Comparison with Experiments. Plant Soil. Kluwer Academic Publishers; 2006;283: 99–117.
1015	123. Schnepf A, Leitner D, Landl M, Lobet G, Mai TH, Morandage S, et al. CRootBox: A Structural-Functional Modelling Framework For Root Systems [Internet]. bioRxiv. 2017. p. 139980. doi:10.1101/139980
	124. Pagès L, Vercambre G, Drouet J-L, Lecompte F, Collet C, LeBot J. RootTyp: a generic model to depict and analyse the root system architecture. Plant Soil. Kluwer Academic Publishers; 2004;258: 103–119.
1020	125. Barczi J-F, Rey H, Griffon S, Jourdan C. DigR: a generic model and its open source simulation software to mimic three-dimensional root-system architecture diversity. Ann Bot. 2018; doi:10.1093/aob/mcy018
1025	126. Dupuy L, Fourcaud T, Stokes A, Danjon F. A density-based approach for the modelling of root architecture: application to Maritime pine (Pinus pinaster Ait.) root systems. J Theor Biol. Elsevier; 2005;236: 323–334.
	127. Huber K, Vanderborght J, Javaux M, Schröder N, Dodd IC, Vereecken H. Modelling the impact of heterogeneous rootzone water distribution on the regulation of transpiration by hormone transport and/or hydraulic pressures. Plant Soil. Kluwer Academic Publishers; 2014; Available: http://link.springer.com/10.1007/s11104-014-2188-4
1030	 Huber K, Vanderborght J, Javaux M, Vereecken H. Simulating transpiration and leaf water relations in response to heterogeneous soil moisture and different stomatal control mechanisms. Plant Soil. 2015;394: 109–126.
1035	129. Hack-ten Broeke MJD, Kroes JG, Bartholomeus RP, van Dam JC, de Wit AJW, Supit I, et al. Quantification of the impact of hydrology on agricultural production as a result of too dry, too wet or too saline conditions [Internet]. 2016. doi:10.5194/soil-2016-24
	 Gijsman AJ, Jagtap SS, Jones JW. Wading through a swamp of complete confusion: how to choose a method for estimating soil water retention parameters for crop models. Eur J Agron. 2002;18: 77–106.
1040	 Landl M, Huber K, Schnepf A, Vanderborght J, Javaux M, Glyn Bengough A, et al. A new model for root growth in soil with macropores. Plant Soil. Springer International Publishing; 2016; 1–18.
	132. Schroeder N, Javaux M, Vanderborght J, Steffen B, Vereecken H. Effect of Root Water and Solute Uptake on Apparent Soil Dispersivity: A Simulation Study. Vadoze Zone Journal. 2012;

Available: http://vzj.geoscienceworld.org/content/11/3/vzj2012.0009.abstract

- Roose T, Keyes SD, Daly KR, Carminati A, Otten W, Vetterlein D, et al. Challenges in imaging and predictive modeling of rhizosphere processes. Plant Soil. Kluwer Academic Publishers; 2016; 1–30.
 - 134. Schwartz N, Carminati A, Javaux M. The impact of mucilage on root water uptake—A numerical study. Water Resour Res. 2016;52: 264–277.

1050

- 135. Vereecken H, Schnepf A, Hopmans JW, Javaux M, Or D, Roose T, et al. Modeling Soil Processes: Review, Key Challenges, and New Perspectives. Vadose Zone J. 2016;15: 0.
 - Zeng, G., Birchfield, S. T., & Wells, C. E. (2008). Automatic discrimination of fine roots in minirhizotron images. *The New Phytologist*, *117*(2), 549–557. Retrieved from http://doi.wiley.com/10.1111/j.1469-8137.2007.02271.x